

Regular Article

Autistic traits, alexithymia, and emotion recognition of human and anime faces

Bridger J. Standiford and Kevin J. Hsu 

Department of Psychological and Social Sciences, Pennsylvania State University, Abington, PA, USA

Abstract

Individuals on the autism spectrum or with elevated autistic traits have shown difficulty in recognizing people's facial emotions. They also tend to gravitate toward anime, a highly visual medium featuring animated characters whose facial emotions may be easier to distinguish. Because autistic traits overlap with alexithymia, or difficulty in identifying and describing feelings, alexithymia might explain the association between elevated autistic traits and difficulty with facial emotion recognition. The present study used a computerized task to first examine whether elevated autistic traits in a community sample of 247 adults were associated with less accurate emotion recognition of human but not anime faces. Results showed that individuals higher in autistic traits performed significantly worse on the human facial emotion recognition task, but no better or worse on the anime version. After controlling for alexithymia and other potentially confounding variables, autistic traits were no longer associated with performance on the facial emotion recognition tasks. However, alexithymia remained a significant predictor and fully mediated the relationship between autistic traits and emotion recognition of both human and anime faces. Findings suggest that interventions designed to help individuals on the autism spectrum with facial emotion recognition might benefit from targeting alexithymia and employing anime characters.

Keywords: alexithymia; anime; autism spectrum; autistic traits; emotion; facial emotion recognition

(Received 14 December 2023; revised 20 January 2025; accepted 29 January 2025)

Individuals with autism spectrum conditions (ASC) experience clinically significant difficulty in their social or emotional communication, which can include marked impairment in the appropriate use of non-verbal signals (e.g., facial expressions), recognition or sharing of emotions, and responses to the emotions of others (American Psychiatric Association, 2013). Unusually narrow, rigid, or atypical interests are another characteristic of individuals with ASC. Although they describe a neurodevelopmental condition and a psychiatric diagnosis, ASC are the clinical extreme in a broad spectrum of neurodiversity. Indeed, autistic traits are commonly understood to occur along a continuum not only within clinical populations, but also among the general population (Allison et al., 2012; Baron-Cohen et al., 1997; Ruzich et al., 2016).

Because emotion recognition is key to effective social interactions, research has focused on understanding the characteristics and limitations of this skill in individuals with ASC, or elevated autistic traits. On the one hand, many studies have demonstrated that individuals with ASC, or elevated autistic traits, have more difficulty interpreting the facial emotions of others than do those without ASC, or those who are low in autistic traits (Actis-Grosso et al., 2015; Ashwin et al., 2006; Celani et al., 1999; Clark et al., 2008; Golan et al., 2006; Hobson et al., 1989; Ingersoll, 2010;

Macdonald et al., 1989; McKenzie et al., 2018; Philip et al., 2010; Poljac et al., 2013; Sucksmith et al., 2013; Wallace et al., 2008). On the other hand, some research has failed to demonstrate a relationship between ASC or autistic traits and facial emotion recognition (Baron-Cohen et al., 1997; Harms et al., 2010; Jones et al., 2011; Miyahara et al., 2007; Piggot et al., 2004; Tracy et al., 2011; see Yeung, 2022 for a recent review). Uljarevic and Hamilton (2013) conducted a meta-analysis of 48 studies comparing those formally diagnosed with ASC and those without ASC on their ability to accurately identify the emotions of human faces in behavioral tasks. Although the behavioral tasks used to assess emotion recognition varied from study to study, they most commonly involved asking participants to identify the emotions displayed in a series of faces, or to select the faces matching a series of emotions. After correcting for publication bias, the results of this meta-analysis supported the evidence that individuals with ASC show more difficulty with facial emotion recognition (Hedges' $g = 0.41$).

Despite previous research converging on the general finding that ASC and elevated autistic traits are associated with deficits in facial emotion recognition, it remains unclear why only some individuals on the autism spectrum experience difficulty with recognizing facial emotions, and why those individuals developed this difficulty. In their meta-analysis, Uljarevic and Hamilton (2013) found that factors such as age, intelligence (IQ), and type of behavioral task used in a study did not explain the deficits in facial emotion recognition found among individuals with ASC. They suggested instead that a lack of attending to some important

Corresponding author: Kevin J Hsu; Email: khsu@psu.edu

Cite this article: Standiford, B. J., & Hsu, K. J. (2025). Autistic traits, alexithymia, and emotion recognition of human and anime faces. *Development and Psychopathology*, 1–14, <https://doi.org/10.1017/S0954579425000100>



features of human faces, such as the eyes, might explain these deficits. Indeed, a different line of research has revealed that individuals with ASC or elevated autistic traits may be less oriented to human faces (Black et al., 2017; Dawson et al., 2005; Guillon et al., 2016; Klin et al., 2002; Pelphrey et al., 2002; Vettori et al., 2020) and find viewing them less rewarding (Chevallier et al., 2012; Cuve et al., 2018; Silva et al., 2015; Stuart et al., 2023; Tottenham et al., 2014).

Autism spectrum, anime, and facial emotion recognition

Atherton and Cross (2018) suggested that the reduced saliency of human faces for individuals with ASC or elevated autistic traits, combined with their propensity to develop strong circumscribed interests, results in their tendency to gravitate toward non-human social interests, such as cartoons. In line with this reasoning, Silva et al. (2015) found that children diagnosed with ASC showed a preference for viewing cartoon faces over human faces and were, in fact, avoidant of human faces. Jones et al. (2011) reported survey data indicating that adolescents with ASC spend a large amount of time viewing screen-based media. Mazurek et al. (2012) and Shane and Albert (2008) similarly noted that parents of children with ASC reported electronic screen engagement, and especially animated television shows and movies, as their child's most common leisure activity. Finally, individuals with ASC or elevated autistic traits have specifically shown a strong preference for *anime* (Atherton et al., 2023), a style of Japanese animation that has rapidly grown in popularity since the introduction of *Pokémon* in the 1990s (Allison, 2006; Napier, 2006; Otmazgin, 2014). Anime is a specific type of cartoon that has a distinct art style originating in Japan and features vibrant characters with exaggerated facial expressions. Cartoons include anime and a range of other art styles, but the term usually refers to Western animation. One study analyzed the constricted interests that tend to characterize individuals on the autism spectrum, revealing that anime was the most frequently reported topic of interest (South et al., 2005). Another study examining the media use of adolescents with ASC found that websites focused on anime were the second most popular among all the websites that they visited, after websites discussing video games (Kuo et al., 2014).

The well-documented interest in viewing animated media and the specific preference for anime among individuals on the autism spectrum have been attributed to the use of distinct characters in these media, where facial expressions are exaggerated to the point of caricature (Atherton et al., 2023; Liu et al., 2019). When anime characters show surprise, excitement, or happiness, for example, their eyes widen to comically unrealistic proportions. Similarly, sadness in anime characters is often expressed through a waterfall of tears literally splashing down their faces. These exaggerated displays of emotional expression in the faces of anime characters might make them especially clear and salient even to individuals with ASC or elevated autistic traits. Consistent with this idea, Berthoz et al. (2013), Rump et al. (2009), and Song and Hakoda (2018) found that individuals with ASC were less impaired in their facial emotion recognition when presented with exaggerated, overt emotional expressions, but had more difficulty accurately interpreting subtle facial emotions. Rozema (2015) suggested that the visual tropes used in media like anime to communicate emotions provide clear and repetitive visual cues, allowing individuals on the autism spectrum to recognize and remember underlying patterns of emotional expressions more easily. As a

result, individuals with ASC or elevated autistic traits may develop a particular affinity for anime relative to other interests.

While it has been suggested that individuals on the autism spectrum prefer anime because they can more easily understand the often-exaggerated facial emotions of the characters, no previous studies of facial emotion recognition have used anime characters to test this possibility. Furthermore, there have not been many previous studies that directly examined facial emotion recognition for animated or cartoon faces in individuals with ASC or elevated autistic traits. In one study, Rosset et al. (2008) found that children with ASC processed cartoon faces like typically developing children processed human faces, and their facial emotion recognition when viewing cartoon faces was not impaired to the same degree as it was when viewing human faces. Brosnan et al.'s (2015) study supported and expanded these findings, showing that adolescents with ASC not only performed better when viewing cartoon faces compared with human faces, but they also outperformed those without ASC in accurately recognizing emotions expressed in static cartoon faces. Similarly, Cross et al. (2019) found that adolescents with ASC showed improvement in facial emotion recognition when human faces were put through an animal filter and thus appeared like anthropomorphic lion or gorilla faces. In another study, Atherton and Cross (2019) showed that the use of animal versus human characters in social stories was related to improved recognition of social norm violations among those high in autistic traits. Finally, two recent studies examined the extent to which ASC (Cross et al., 2022) or elevated autistic traits (Atherton & Cross, 2022) were related to facial emotion recognition accuracy using both a human and a cartoon version of a test for reading emotions in the eyes. They found that adults with ASC were no different, and adults with elevated autistic traits were significantly worse than those without ASC or with low autistic traits on the human test. However, the group with ASC performed better, and the group with elevated autistic traits no differently, when the cartoon version of the test was used. These findings suggest that facial emotion recognition deficits in individuals with ASC or high in autistic traits are not global, but specific to the evaluation of human faces.

Autism spectrum, alexithymia, and facial emotion recognition

Alexithymia is a subclinical trait characterized by difficulties in identifying and describing feelings, associating bodily sensations with specific feeling states, and using words to express emotions (Berthoz et al., 2011; Nemiah et al., 1976). Like autistic traits, alexithymia appears to exist along a continuum, with higher levels reflecting more difficulty in the cognitive processing and regulation of emotions (Taylor et al., 1997). Studies have revealed a much higher rate of alexithymia among individuals on the autism spectrum relative to the general population, with estimates for autistic populations ranging from 50%–85% (Berthoz & Hill, 2005; Berthoz et al., 2013; Hill et al., 2004). For comparison, the prevalence rate of alexithymia in the general population is estimated to be closer to 10% (Linden et al., 1995).

Some researchers have found evidence that the difficulties in emotion recognition observed among individuals on the autism spectrum might not be the result of social deficits characteristic of ASC or autistic traits, but rather the frequently co-existing condition of alexithymia (Bernhardt et al., 2014; Bird & Cook, 2013; Cook et al., 2013; Heaton et al., 2012; Milosavljevic et al., 2016; Oakley et al., 2016; Ola & Gullon-Scott, 2020; Santiesteban

et al., 2021; Trevisan et al., 2016). Indeed, one study found that reduced eye gaze to facial emotional expressions (e.g., lower fixation count and duration) was associated more with alexithymia than with either ASC or autistic traits in a sample of adults with and without a formal diagnosis (Cuve et al., 2021). In another study examining the role of alexithymia and anxiety in the relationship between empathic ability and autistic traits, Brett and Maybery (2022) found that autistic traits predicted empathy through alexithymia and anxiety.

Based on their systematic review and meta-analysis of 15 studies, Kinnaird et al. (2019) concluded that it was common but not universal for individuals on the autism spectrum to experience co-occurring alexithymia, finding a mean prevalence rate of 50%. They further suggested that emotional recognition difficulties traditionally associated with ASC or autistic traits might be better explained by alexithymia, following previous research (e.g., Bird & Cook, 2013). Most recently, results from factor analysis of measures widely used to assess alexithymia and autistic traits indicated that these two constructs were distinct, because items from each measure loaded on separate factors (Cuve et al., 2022). Network analyses similarly produced separate clusters comprising only of items representing either alexithymic or autistic traits.

Considering that alexithymia might characterize approximately half or more of the autistic population, other researchers have argued that alexithymia is a core feature or consequence of ASC or autistic traits rather than a distinct condition (Gaigg, 2012; Quattrocki & Friston, 2014). They have also pointed to the problems of relying on self-report measures to assess alexithymic and autistic traits, noting that self-report could bias the conclusions drawn from this body of research (Gaigg et al., 2018). The significant overlap between alexithymia and ASC or autistic traits has led to the theory that they share the same underlying developmental process leading to their observed social and emotional deficits. Several neuroimaging studies have contributed evidence to support this theory (Grynberg et al., 2012; Moriguchi et al., 2006; Silani et al., 2008), showing that there are some neural components potentially shared between individuals high in alexithymia and individuals on the autism spectrum, specifically those brain regions responsible for perspective taking and mentalizing (i.e., understanding the mental states of the self and others).

Discussing the complex relationship between alexithymia and ASC, Poquérousse et al. (2018) acknowledged that there is no clear answer as to whether they are co-occurring conditions or whether alexithymia is a common component of ASC. In the absence of consensus on whether ASC and alexithymia (or autistic and alexithymic traits more broadly) may be considered distinct but commonly co-occurring conditions relevant to facial emotion recognition deficits, our hypotheses and study design were guided by an understanding that it would be important to evaluate the role of both autistic and alexithymic traits on facial emotion recognition.

The present research

To address gaps in the previous literature, the present study used a computerized task to examine not only whether individuals higher in autistic traits are less accurate in recognizing the facial emotional expressions of human characters, consistent with past studies, but also whether they do not experience this same difficulty in recognizing emotions when presented with anime faces. In other words, we hypothesized that elevated autistic traits would be

associated with lower emotion recognition scores when human faces are used as targets, but not when anime faces are used as targets. Anime faces were chosen because they are drawn with exaggerated facial expressions that may help to reduce some of the visual processing deficits associated with ASC and autistic traits (Atherton et al., 2023; Liu et al., 2019; Rozema, 2015), and because there is a demonstrated preference for anime among individuals in the autistic community (Kuo et al., 2014; South et al., 2005).

A measure of alexithymia was included to examine whether higher scores on this subclinical trait are also related to more difficulty recognizing human and anime facial emotions, and whether alexithymia can fully or in part explain the associations between autistic traits and facial emotion recognition. Although we hypothesized that elevated alexithymia would be associated with lower human facial emotion recognition scores, we did not have a strong prediction about the relationship between alexithymia and anime facial emotion recognition scores. However, we expected that when autistic traits and alexithymia control for one another in predicting human and anime facial emotion recognition scores, alexithymia would explain more of the variance in those facial emotion recognition scores than autistic traits.

If individuals higher in autistic traits do not show the same difficulty recognizing the facial emotional expressions of anime characters as they do the expressions of human characters, these findings could inform the development of more effective clinical interventions aimed at improving facial emotion recognition and positive social behaviors among individuals on the autism spectrum. If alexithymia is found to be more relevant than autistic traits in explaining facial emotion recognition, interventions could more specifically target alexithymia.

Method

Participants

Participants were 247 adults (129 male, 115 female, 2 non-binary, and 1 other) ranging in age from 18 to 63 years old ($M = 27.65$ years, $SD = 10.63$). They included 125 who were recruited from the psychology participant pool at Pennsylvania State University Abington, and another 122 from the crowdsourcing website Amazon Mechanical Turk (<https://www.mturk.com/mturk/welcome>). Amazon Mechanical Turk connects paid volunteers with researchers who are looking for participants to complete their studies. Participants engaged in this online study on their own computer and at their own convenience by first choosing from a list of available studies either through Amazon Mechanical Turk or the psychology participant pool, and then by being directed to a website to complete the study. After completing the study, participants from the psychology participant pool received credits toward a research requirement in their psychology course, and those recruited from Amazon Mechanical Turk were compensated \$1.00. A wage analysis of 4,500 studies hosted on Amazon Mechanical Turk between 2015 and 2019 estimated that participants were completing studies for an average of \$6.00–\$7.00 an hour (Moss et al., 2023). Because our study had a median completion time of about 10 minutes, payment of \$1.00 closely approximated the average hourly earnings for participants on Amazon Mechanical Turk. Furthermore, we were limited by funding constraints. This study and its procedures (including compensation) were approved by Pennsylvania State University's Institutional Review Board.

Participants from Amazon Mechanical Turk were significantly older ($M = 35.45$ years, $SD = 8.90$) than those from the psychology

participant pool ($M = 20.03$ years, $SD = 5.33$), $t(197) = 16.46$, $p < .0001$. This age difference was expected, because we intentionally recruited participants from Amazon Mechanical Turk to obtain a more representative sample, knowing that the psychology participant pool is only open to undergraduate students at our campus. While the reliability and validity of data collected on Amazon Mechanical Turk have been supported by some researchers (Buhrmester et al., 2018; Paolacci & Chandler, 2014), others have recently raised concerns about data collection on this platform (Chmielewski & Kucker, 2020; Newman et al., 2021; Webb & Tangney, 2024).

Participants from either recruitment source were excluded from all data analyses if they did not reach the end of the study ($n = 18$) or completed it in less than five minutes ($n = 8$). We further excluded those participants who provided the same rating to all 10 items on the Short Autism Spectrum Quotient ($n = 5$) or to all 23 items on the Revised Toronto Alexithymia Scale ($n = 8$), because these response patterns suggest that they were not paying close attention to the questions (see next section for descriptions of these measures). The numbers of participants excluded for these reasons were not mutually exclusive, and participants excluded for one reason often qualified to be excluded for another reason. Interestingly, all participants excluded for not reaching the end of the study or completing it in less than five minutes were from Amazon Mechanical Turk, while all but two participants excluded for providing the same rating to all items on at least one of the measures were from the psychology participation pool.

Measures

Short Autism Spectrum Quotient

A short form of the Autism Spectrum Quotient (AQ-10; Allison et al., 2012) was previously developed and validated for measuring autistic traits with 10 items. Example items include statements intended to assess social deficits: "I find it difficult to work out people's intentions," or "I find it easy to work out what someone is thinking or feeling just by looking at their face." Example items also include statements intended to assess unusually narrow, rigid, or atypical interests, among other autistic traits: "I like to collect information about categories of things (e.g. types of car, types of bird, types of train, types of plant)" and "I often notice small sounds when others do not." For each of the 10 items, participants were asked whether they *definitely agree*, *slightly agree*, *slightly disagree*, or *definitely disagree* with the statement. On normally coded items, responses of *definitely agree* or *slightly agree* earned one point. On reverse coded items, responses of *definitely disagree* or *slightly disagree* earned one point. Thus, scores on this measure range from 0–10. Higher scores on the AQ-10 indicate higher levels of autistic traits. The internal consistency of the AQ-10 in this sample was low ($\alpha = 0.45$) but consistent with previous findings (Jia et al., 2019; Sizoo et al., 2015; Taylor et al., 2020).

Revised Toronto Alexithymia Scale

The Revised Toronto Alexithymia Scale (TAS-R; Taylor et al., 1992) is a 23-item scale used to measure alexithymia, which comprises the inability (or limited ability) of a person to experience, identify, or describe emotions, or to relate them to the stimulus that caused them. Example items include statements such as: "I have feelings that I can't quite identify," or "I often don't know why I am angry." Participants were asked to indicate on a 5-point Likert scale the extent to which they agree or disagree with each item (1 = *strongly disagree*, 2 = *moderately disagree*,

3 = *neither agree nor disagree*, 4 = *moderately agree*, or 5 = *strongly agree*). Of the 23 items, six are reverse coded, such that responses of *strongly agree* or *moderately agree* are scored as 1 and 2, respectively, and responses of *moderately disagree* or *strongly disagree* are scored as 4 and 5, respectively. A total score on the TAS-R is obtained by taking the sum of responses to all 23 items, resulting in a range of 23 to 115, with higher scores indicating higher levels of alexithymia. Taylor et al. (1992) found that the mean score for individuals clinically judged to meet criteria for alexithymia was 66.4 ($SD = 13.4$), while the mean score for individuals who were not judged to meet criteria was 56.7 ($SD = 12.2$). The internal consistency of the TAS-R in this sample was high ($\alpha = 0.88$).

Facial emotion recognition test

Participants completed a facial emotion recognition test on their computer. First, they were randomly presented with two human and two anime faces to become familiar with the task. This brief practice was then followed by a series of 12 human and 12 anime faces presented in random order. Each face expressed one of six different emotions: happiness, sadness, anger, fear, disgust, or surprise. These emotions were chosen based on previous research, which found that the facial expressions of these six emotions are universally recognizable across all cultures (Ekman & Friesen, 1971; Izard et al., 1984; Izard, 1994; Matsumoto, 2001). During the test, each face was displayed for three seconds before disappearing, at which point the participant was asked to choose which of the six emotions was expressed by the face they were just shown. Participants earned one point for each facial expression they identified accurately, for a total of 12 possible points on the human section of the facial emotion recognition test, and 12 possible points on the anime section. The timing of three seconds was chosen after reviewing a previous study that analyzed multiple facial emotion recognition tests, finding that two to three seconds was ideal for preventing ceiling effects, wherein neurotypical individuals can too easily receive perfect scores (Wilhelm et al., 2014).

The human faces in the facial emotion recognition test were sourced from a previously validated set of faces with differing expressions, called the Warsaw Set of Emotional Facial Expression Pictures (Olszanowski et al., 2015). This collection includes front-facing images of 30 White European individuals with acting experience, expressing a variety of emotions. These images were rated by over 1,300 independent judges to determine how well the facial expressions matched the intended emotion. For the present study, we chose the six male and six female human faces that were determined by judges to most accurately depict each of the six universally recognizable emotions. Thus, the final stimulus set included 12 different human faces: a male and a female face expressing happiness, a male and a female face expressing sadness, a male and a female face expressing anger, a male and a female face expressing fear, a male and a female face expressing disgust, and a male and a female face expressing surprise. For the practice section of the facial emotion recognition test, we used a male human face expressing sadness and a female human face expressing happiness that were different from those included in the final stimulus set.

Because no previously validated set of anime facial expressions existed, images were collected by the first author using Google Images for the sole purposes of this study. Using the search terms *anime*, *manga*, *facial expressions*, *emotion*, and *facial emotional expressions* in combination with one another, a preliminary compilation of 55 images was created. Due to constraints in time

Table 1. Means, standard deviations, and correlations among study variables

Measure	<i>M</i> (<i>SD</i>)	1.	2.	3.	4.	5.	6.
1. Age	27.65 (10.63)	–					
2. Frequency of social interaction	7.08 (1.29)	–.43***	–				
3. Frequency of anime or manga use	4.35 (2.35)	.13*	–.05	–			
4. AQ-10	3.79 (1.77)	–.03	–.12	.11	–		
5. TAS-R	64.37 (14.84)	.30***	–.27***	.32***	.33***	–	
6. Human facial emotion recognition score	9.50 (2.18)	–.29***	.27***	–.32***	–.20**	–.44***	–
7. Anime facial emotion recognition score	6.81 (2.37)	–.49***	.34***	–.07	–.05	–.37***	.47***

Note. $N = 247$. All correlations used Spearman's ρ . AQ-10 = Short Autism Spectrum Quotient (Allison et al., 2012); TAS-R = Revised Toronto Alexithymia Scale (Taylor et al., 1992). * $p < .05$, ** $p < .005$, *** $p < .0001$.

and funding, we could not recruit a separate sample for a pilot study in which these images of anime faces could be systematically rated on how much they represented the emotions that they were supposed to depict. Because we could not validate the anime stimuli with a pilot study, we reviewed and discussed the images among ourselves before choosing the 12 images that were determined to best express each of the six universally recognizable emotions. The final stimulus set included one male and one female anime face expressing happiness, one male and one female anime face expressing sadness, two male and one female anime face expressing anger, two female anime faces expressing fear, one male and one female anime face expressing disgust, and one male anime face expressing surprise. The same male anime character was used to express three of the six emotions (surprise, sadness, and happiness), while the other nine anime characters were unique. For the practice section of the facial emotion recognition test, we used a male anime face expressing happiness and the same male anime face expressing anger. This anime character was also different from the others. Because anime faces are less familiar to the general population, we provide the complete set of images (both human and anime faces) used in the facial emotion recognition test at the following link: <https://osf.io/qyzgd>.

Potentially confounding variables

To address potential confounds, participants answered the following two additional questions about their frequency of social interaction and their frequency of anime or manga use, respectively: “Approximately how often do you have face-to-face social interactions with others?” and “Approximately how often do you watch anime or read manga?” Responses to both questions were on an 8-point scale (1 = *never*, 2 = *less than once per year*, 3 = *once per year*, 4 = *once per month*, 5 = *once per week*, 6 = *2–3 times per week*, 7 = *once per day*, or 8 = *every day, multiple times per day*).

Results

Preliminary analyses

Table 1 presents the means, standard deviations, and correlations among the study variables, including age, frequency of social interaction, frequency of anime or manga use, AQ-10, TAS-R, human facial emotion recognition score, and anime facial emotion recognition score. Due to the non-normal distributions for most of the study variables, Spearman's rank order correlation coefficients

were used for tests of associations instead of Pearson's correlation coefficients.

As shown in Table 1, participants reported high frequencies of social interaction (i.e., the relevant distribution was negatively skewed), but moderate and more varied frequencies of anime or manga use. Scores on the AQ-10 also varied, and their distribution was positively skewed. Interestingly, 40 participants (16% of the sample) scored 6 or higher on this scale, which is the cutoff that has been used to refer individuals for the assessment of clinically diagnosed ASC (Allison et al., 2012). Scores on the TAS-R showed considerable variation and a slight negative skew in their distribution. Finally, the distribution of scores on the human section of the facial emotion recognition test was negatively skewed, with most participants correctly identifying the emotions of more than 9 out of 12 faces, while scores on the anime section were more normally distributed.

There was a strong positive correlation between scores on the human section and scores on the anime section of the facial emotion recognition test, $r_s(245) = .47, p < .0001$. Furthermore, the potentially confounding variables of age, frequency of social interaction, and frequency of anime or manga use were significantly correlated not only with TAS-R, but also with scores on the human and anime facial emotion recognition test (see Table 1). The one exception was that frequency of anime or manga use was negatively, but not significantly correlated with anime facial emotion recognition scores, $r_s(245) = -.07, p = .2967$.

Correlations of AQ-10 and TAS-R with human and anime facial emotion recognition

Table 1 shows that in accordance with our hypothesis, higher scores on the AQ-10 were negatively and significantly correlated with performance on the human facial emotion recognition test, $r_s(245) = -.20, p = .0015$, but not with performance on the anime facial emotion recognition test, $r_s(245) = -.05, p = .3937$. Thus, participants with more autistic traits tended to have more difficulty in accurately recognizing emotional expressions in human faces. When anime faces were presented, however, those higher in autistic traits did not show significantly more difficulty with emotion recognition than did those lower in autistic traits. Figure 1 visually depicts the correlations between scores on the AQ-10 and scores on the human and anime facial emotion recognition tests in the form of two scatterplots.

Table 1 shows that higher scores on the TAS-R were negatively and significantly correlated with performance on both the human facial emotion recognition test, $r_s(245) = -.44, p < .0001$, and the




Figure 1. Correlations of autistic traits with performance on the human (left) and anime (right) facial emotion recognition tests. Note. $N = 247$. Points represent individual participants. Shaded regions represent 95% confidence intervals. AQ-10 = Short Autism Spectrum Quotient (Allison et al., 2012).




Figure 2. Correlations of alexithymia with performance on the human (left) and anime (right) facial emotion recognition tests. Note. $N = 247$. Points represent individual participants. Shaded regions represent 95% confidence intervals. TAS-R = Revised Toronto Alexithymia Scale (Taylor et al., 1992).

anime facial emotion recognition test, $r_s(245) = -.37, p < .0001$. In contrast to the AQ-10, TAS-R showed much stronger and more significant associations with both human and anime versions of the facial emotion recognition test. Participants with more alexithymia tended to have more difficulty with recognizing the emotions in faces regardless of whether they were human or anime, although the correlation was smaller for anime faces. Figure 2 visually depicts the correlations between scores on the TAS-R and scores on the human and anime facial emotion recognition tests in the form of two scatterplots.

Multiple regression analyses

Two hierarchical multiple regression analyses were conducted to determine the extent to which autistic traits and alexithymia were unique predictors of performance on both the human and anime facial emotion recognition tests. In the first model of the

hierarchical multiple regression, we entered age, frequency of social interaction, and frequency of anime or manga use as predictor variables, because these variables were significantly associated with performance on either or both of the facial emotion recognition tests. In the second model of the hierarchical multiple regression, we entered the two competing variables of interest, AQ-10 and TAS-R, as predictor variables in addition to those included in the first model. The dependent variables for these two hierarchical multiple regression analyses were scores on the human and anime facial emotion recognition tests. Before conducting multiple regression analyses, the variance inflation factors of predictor variables were examined to ensure a lack of multicollinearity among them. Variance inflation factors did not exceed 1.30, which indicated that each predictor variable contained sufficiently unique information over and above that provided by the others.

Table 2. Hierarchical multiple regression results for age, frequency of social interaction, frequency of anime or manga use, autistic traits, and alexithymia predicting human and anime facial emotion recognition scores

Model	Predictor	Human facial emotion recognition score	Anime facial emotion recognition score
1	Age	-0.18** (0.01)	-0.37*** (0.01)
	Frequency of social interaction	0.19** (0.05)	0.15* (0.05)
	Frequency of anime or manga use	-0.30*** (0.03)	-0.05 (0.03)
	<i>F</i>	19.92***	20.48***
	<i>R</i> ²	0.20	0.20
	<i>R</i> ² _{adj}	0.19	0.19
2	Age	-0.15* (0.01)	-0.33*** (0.01)
	Frequency of social interaction	0.14* (0.05)	0.11 (0.05)
	Frequency of anime or manga use	-0.22** (0.03)	0.02 (0.03)
	AQ-10	-0.04 (0.04)	0.01 (0.04)
	TAS-R	-0.28*** (0.00)	-0.27*** (0.00)
	<i>F</i>	18.03***	17.09***
	<i>R</i> ²	0.27	0.26
	<i>R</i> ² _{adj}	0.26	0.25

Note. *N* = 247. All values are partial regression coefficients (standardized beta weights) or standard errors (as indicated in parentheses) except for those under *F*, *R*², and *R*²_{adj}, which are the *F* ratio, coefficient of determination, and adjusted coefficient of determination, respectively. AQ-10 = Short Autism Spectrum Quotient (Allison et al., 2012); TAS-R = Revised Toronto Alexithymia Scale (Taylor et al., 1992). **p* < .05, ***p* < .005, ****p* < .0001.

Table 2 presents the results of the two hierarchical multiple regression analyses among participants. In the first model, it was revealed that age ($\beta = -0.18$, $p = .0043$), frequency of social interaction ($\beta = 0.19$, $p = .0022$), and frequency of anime or manga use ($\beta = -0.30$, $p < .0001$) were unique and significant predictors of human facial emotion recognition scores, even with each predictor variable controlling for one another. Thus, participants who were older and watched more anime or read more manga scored lower on the human facial emotion recognition test, while those who had more frequent social interactions scored higher. For anime facial emotion recognition scores, only age ($\beta = -0.37$, $p < .0001$) and frequency of social interaction ($\beta = 0.15$, $p = .0160$) were unique and significant predictors controlling for one another. Participants who were older scored lower on the anime facial emotion recognition test, while those who had more frequent social interactions scored higher. Frequency of anime or manga use ($\beta = -0.05$, $p = .3627$), however, was not significantly associated with performance on the anime facial emotion recognition test.

In the second model, with the addition of AQ-10 and TAS-R as predictor variables, it was revealed that age ($\beta = -0.15$, $p = .0140$), frequency of social interaction ($\beta = 0.14$, $p = .0198$), frequency of anime or manga use ($\beta = -0.22$, $p = .0002$), and TAS-R ($\beta = -0.28$, $p < .0001$) were unique and significant predictors of human facial emotion recognition scores, even with each variable controlling for one another. Thus, participants who were older, watched more anime or read more manga, and experienced more alexithymia scored lower on the human facial emotion recognition test, while those who had more frequent social interactions scored higher. For

anime facial emotion recognition scores, only age ($\beta = -0.33$, $p < .0001$) and TAS-R ($\beta = -0.27$, $p < .0001$) were unique and significant predictors controlling for one another. Participants who were older and experienced more alexithymia scored lower on the anime facial emotion recognition test, and their frequency of social interaction ($\beta = 0.11$, $p = .0786$) no longer significantly predicted their performance. Regardless of whether the multiple regression model was predicting performance on the human or anime facial emotion recognition test, autistic traits ($\beta = -0.04$, $p = .4523$ and $\beta = 0.01$, $p = .8356$, respectively) were not significantly predictive after controlling for alexithymia and the other three variables, in contrast to the zero-order correlations.

The first model explained about 20% of the variance in both human [$F(3, 243) = 19.92$, $p < .0001$] and anime facial emotion recognition scores [$F(3, 243) = 20.48$, $p < .0001$] among participants. The second and final model improved slightly on the first model, explaining about 27% of the variance in human facial emotion recognition scores [$F(5, 241) = 18.03$, $p < .0001$] and about 26% of the variance in anime facial emotion recognition scores [$F(5, 241) = 17.09$, $p < .0001$] among participants. For human facial emotion recognition scores, TAS-R was the strongest contributor, followed by frequency of anime or manga use. For anime facial emotion recognition scores, age was the strongest contributor to the variance, followed by TAS-R.

Mediation analyses

The zero-order correlation between AQ-10 and human (but not anime) facial emotion recognition scores was significant, as was the zero-order correlation between AQ-10 and TAS-R. After controlling for TAS-R in the second model of the hierarchical multiple regression analyses, however, AQ-10 was no longer significantly associated with performance on the human facial emotion recognition test. These results suggest that TAS-R fully mediated the relationship between AQ-10 and human facial emotion recognition scores, based on the procedures from Baron and Kenny (1986). To further investigate this possibility, mediation analyses were conducted to determine the extent to which alexithymia mediated the relationship between autistic traits and performance on both the human and anime facial emotion recognition tests. We tested for mediation of autistic traits and anime facial emotion recognition scores through alexithymia, despite finding no significant zero-order correlation between AQ-10 and anime emotion scores, because many researchers have suggested that mediation is still possible under such conditions (Collins et al., 1998; Preacher & Hayes, 2004; Rucker et al., 2011; Shrout & Bolger, 2002; Zhao et al., 2010).

Figure 3 presents a path diagram for the mediation analysis in which AQ-10 predicted human facial emotion recognition scores through TAS-R. The standardized regression coefficient between AQ-10 and TAS-R was statistically significant ($\beta = 0.31$, $p < .0001$), as was the standardized regression coefficient between TAS-R and human facial emotion recognition scores ($\beta = -0.41$, $p < .0001$). The standardized indirect effect of AQ-10 on human facial emotion recognition scores was $(0.31)(-0.41) = -0.13$. We tested the significance of this indirect effect using bootstrapping procedures. Unstandardized indirect effects were computed for each of 1,000 bootstrapped samples, and the 95% confidence interval (CI) was computed with the indirect effects at the 2.5 and 97.5 percentiles. The bootstrapped unstandardized indirect effect of AQ-10 on human facial emotion recognition scores was -0.16 , 95% CI = $[-0.24, -0.09]$, $p < .0001$. These results show that despite

Figure 3. Path diagram for the mediation analysis in which autistic traits predicted human facial emotion recognition scores through alexithymia. Note. $N = 247$. Standardized regression coefficients are depicted for the relationship between autistic traits and performance on the human facial emotion recognition test as mediated by alexithymia. The standardized regression coefficient between autistic traits and human facial emotion recognition scores, controlling for alexithymia, is in parentheses. Brackets indicate 95% confidence intervals for each standardized regression coefficient. AQ-10 = Short Autism Spectrum Quotient (Allison et al., 2012); TAS-R = Revised Toronto Alexithymia Scale (Taylor et al., 1992). * $p < .0001$.





Figure 4. Path diagram for the mediation analysis in which autistic traits predicted anime facial emotion recognition scores through alexithymia. Note. $N = 247$. Standardized regression coefficients are depicted for the relationship between autistic traits and performance on the anime facial emotion recognition test as mediated by alexithymia. The standardized regression coefficient between autistic traits and anime facial emotion recognition scores, controlling for alexithymia, is in parentheses. Brackets indicate 95% confidence intervals for each standardized regression coefficient. AQ-10 = Short Autism Spectrum Quotient (Allison et al., 2012); TAS-R = Revised Toronto Alexithymia Scale (Taylor et al., 1992). * $p < .0001$.



the lack of a direct effect of AQ-10 on human facial emotion recognition scores ($\beta = -0.02$, $p = .6913$), there was a significant indirect effect of AQ-10 through TAS-R.

Figure 4 presents a path diagram for the mediation analysis in which AQ-10 predicted anime facial emotion recognition scores through TAS-R. The standardized regression coefficient between AQ-10 and TAS-R was statistically significant ($\beta = 0.31$, $p < .0001$), as was the standardized regression coefficient between TAS-R and anime facial emotion recognition scores ($\beta = -0.37$, $p < .0001$). The standardized indirect effect of AQ-10 on anime facial emotion recognition scores was $(0.31)(-0.37) = -0.12$. We tested the significance of this indirect effect using bootstrapping procedures. Similar to the previous mediation analysis, unstandardized indirect effects were computed for each of 1,000 bootstrapped samples, and the 95% CI was computed with the indirect effects at the 2.5 and 97.5 percentiles. The bootstrapped unstandardized indirect effect of AQ-10 on anime facial emotion recognition scores was -0.16 , 95% CI = $[-0.25, -0.08]$, $p < .0001$. These results show that despite the lack of a direct effect of AQ-10 on anime facial emotion recognition scores ($\beta = 0.07$, $p = .2971$), there was a significant indirect effect of AQ-10 through TAS-R.

Comparing analyses between participants from the psychology participant pool and Amazon Mechanical Turk

Supplementary Tables 1–6 present the results of analyses from the previous sections, separated by whether participants were from the subsample recruited from the psychology participant pool ($n = 125$) or Amazon Mechanical Turk ($n = 122$). The same patterns emerged regardless of whether analyses used the full sample ($N = 247$) or either of the subsamples. Thus, our discussion

of findings will only focus on those involving the full sample to avoid repetition.

Discussion

The purpose of this study was to examine whether individuals higher in autistic traits experience more difficulty with recognizing and interpreting the emotional expressions of human faces, while not showing such a deficit for anime faces. Based on previous research that compared the performance of individuals with ASC or elevated autistic traits on human and cartoon versions of facial emotion recognition tests (Atherton & Cross, 2022; Brosnan et al., 2015; Cross et al., 2019, 2022; Rosset et al., 2008), we hypothesized that participants higher in autistic traits would perform worse at a facial emotion recognition test involving human stimuli but not one involving anime stimuli, due to their simplified and exaggerated expressions. We also hypothesized that participants higher in alexithymia would perform worse at the human facial emotion recognition test, and that alexithymia would be more strongly associated than autistic traits with scores on the human and anime facial emotion recognition tests, following previous studies (e.g., Bird & Cook, 2013; Cook et al., 2013; Cuve et al., 2021).

Our hypotheses were supported by the data. Participants higher in autistic traits performed significantly worse at the facial emotion recognition test featuring human stimuli. This finding supports both the validity of the human facial emotion recognition test and previous research demonstrating that individuals on the autism spectrum tend to have greater difficulty interpreting the meaning of human facial expressions (e.g., McKenzie et al., 2018; Poljac et al., 2013; Uljarevic & Hamilton, 2013).

In contrast to the significant negative correlation between scores on the AQ-10 and performance at recognizing human faces, there was no significant correlation between AQ-10 scores and performance on the anime facial emotion recognition test. This finding suggests that while it is not easier for individuals higher in autistic traits to recognize the facial expressions of anime characters compared with individuals lower in autistic traits, it is also not more difficult for them. Past work has found similar evidence that the facial emotion recognition of individuals with ASC or elevated autistic traits is less impaired when viewing cartoon as opposed to human faces (Atherton & Cross, 2022; Brosnan et al., 2015; Cross et al., 2019, 2022; Rosset et al., 2008). We extend this evidence for the first time to anime faces, using a newly developed facial emotion recognition test. The exaggerated facial expressions that are characteristic of anime characters may operate as a protective factor against the deficits in facial emotion recognition typically seen in individuals with ASC or elevated autistic traits (Atherton et al., 2023; Liu et al., 2019; Rozema, 2015). For this reason, individuals on the autism spectrum may be especially drawn to anime (Atherton & Cross, 2018), and it has been shown that they have an affinity for this type of media compared to other interests (Kuo et al., 2014; South et al., 2005). In our sample, more autistic traits were slightly correlated with more frequency of anime or manga use, although this relationship was not significant.

Consistent with our hypotheses and previous research (e.g., Bird & Cook, 2013), the multiple regression and mediation analyses revealed a potentially important caveat in the associations between autistic traits and facial emotion recognition, which is the overlapping relevance of alexithymia. In contrast to the zero-order correlation, autistic traits were no longer significantly predictive of facial emotion recognition in human faces after controlling for alexithymia and potential confounds like age, frequency of social interaction, and frequency of anime and manga use. Additionally, alexithymia was more strongly and negatively correlated with performance on the human and anime facial emotion recognition tests than autistic traits, both with respect to their zero-order correlations and the hierarchical multiple regression models in which they controlled for each other. Finally, alexithymia fully mediated the relationship between autistic traits and both human and anime facial emotion recognition scores. We found no direct effect of autistic traits on emotion recognition when viewing either human or anime faces. However, there was a significant indirect effect of autistic traits on human and anime facial emotion recognition through alexithymia. These findings suggest that while the difficulty recognizing facial emotional expressions, whether in human or anime faces, characterizes individuals high in autistic traits, the overlapping subclinical trait of alexithymia is one possible source of this difficulty. Importantly, this difficulty is attenuated when non-human anime faces are viewed. Future research and clinical interventions might consider the role of alexithymia when it comes to helping autistic individuals with facial emotion recognition.

Issues in the measurement of autism spectrum and facial emotion recognition

While individuals recruited for the current study were not given a formal clinical evaluation or diagnosis, they completed the AQ-10 (Allison et al., 2012), a short scale designed to measure autistic traits. Individuals with higher scores have more autistic traits, and those who score 6 or higher might be referred for an assessment to

formally diagnose ASC but cannot be assumed to have such a diagnosis. Thus, a strength of our sample is that it consists of individuals from the community with varying degrees of autistic traits, allowing for more statistically powerful, robust, and generalizable tests of how individuals on the autism spectrum process facial emotional expressions. Indeed, our sample provided a good distribution of autistic traits as measured by the AQ-10, consistent with the idea that ASC are best conceptualized as the extreme end of a broader continuum of autistic traits (Allison et al., 2012; Baron-Cohen et al., 1997; Ruzich et al., 2016). Further supporting this idea, 40 (16%) of the 247 participants scored 6 or higher on the scale, suggesting that they might be referred for possible clinical diagnosis. Although recruiting individuals with a formal diagnosis of ASC may be applicable for some clinical or research purposes, it cannot account for variability on the autism spectrum or those individuals who may be on the autism spectrum but were never diagnosed. With the AQ-10, autistic individuals unaware of their condition would still be identified due to their heightened autistic traits.

The distributions of participants' scores on both the human and anime facial emotion recognition tests suggest that they may be useful for future research. These two tests clearly distinguish those who can more easily recognize and identify facial emotional expressions from those who have more difficulty. Furthermore, their construct validity is evidenced by their strong and significant correlations with the measure of alexithymia and with each other. The mean score on the human version of the test was higher than the mean score on the anime version, which may reflect the fact that most individuals are less familiar with anime characters than with human characters. It may also reflect the use of human faces that were sourced from a collection of images that had been previously validated (Olszanowski et al., 2015), whereas the anime faces used were found on Google Images by the first author. Future attempts to improve on the reliability and validity of these two facial emotion recognition tests would benefit from a more detailed item analysis of how participants performed with respect to the individual faces chosen (see e.g., Passarelli et al., 2018), but that endeavor is beyond the scope of this study.

Limitations and future directions

Contrary to two previous studies that found individuals with ASC performed better than those without ASC at recognizing emotions expressed in cartoon faces (Brosnan et al., 2015; Cross et al., 2022), individuals higher in autistic traits in the present study were no better or worse at recognizing the facial expressions of anime characters. A similar lack of difference in performance between autistic and neurotypical individuals in facial emotion recognition when viewing cartoon faces was found in two other studies (Atherton & Cross, 2022; Rosset et al., 2008), one of which also recruited individuals varying in autistic traits rather than those diagnosed with ASC. Although we described it as a strength of our sample, it is also a limitation that we recruited participants from community populations rather than those who had been clinically diagnosed with ASC. A sample of individuals with ASC would have allowed more direct comparisons with much of the previous research on autism spectrum and facial emotion recognition, which has focused on clinical populations. Perhaps if the study were to have included participants clinically diagnosed with ASC instead of or in addition to participants with elevated autistic traits, our results would have been different. Importantly, findings that pertain to this sample of individuals varying in autistic traits

cannot be generalized to individuals with ASC or those from clinical populations. We cannot make claims about individuals with ASC based on the present study.

Another major limitation that requires our cautious interpretation of findings is the use of the AQ-10 to measure autistic traits in this study. Because we used the short form instead of the full form of the Autism Spectrum Quotient (AQ; Baron-Cohen et al., 2001), it is not possible to directly compare scores or results related to the AQ-10 from the present study to those using the full form AQ from some of the previous studies that helped provide the empirical basis for this work (Actis-Grosso et al., 2015; Atherton & Cross, 2022; Poljac et al., 2013). An additional concern with using the AQ-10 is that it showed low internal consistency as estimated with Cronbach's alpha, both in our sample and in previous adult samples from the general population (Jia et al., 2019; Sizoo et al., 2015; Taylor et al., 2020). To be sure, the AQ-10 has been commonly used in research since it was developed, specifically as a more efficient and less time-consuming measure of autistic traits in the general population (e.g., Bertrams & Schlegel, 2020; Forby et al., 2023; Gollwitzer et al., 2019; Mason et al., 2021; Pazhoohi et al., 2021; Rudolph et al., 2018). We thus used the AQ-10 in this study because it had been used by other researchers and allowed us to substantially reduce the time it would take for participants to complete the study, an important consideration due to the limited amount of funding we had to compensate them. However, we must emphasize the limitations of the AQ-10 and strongly encourage caution when using it in future research, especially because it demonstrates low internal consistency. Results are potentially biased when measures with low internal consistency like the AQ-10 are used to predict constructs such as facial emotion recognition and compared to measures with much higher internal consistency like the TAS-R. We recommend that researchers use the full form AQ to avoid these issues with the short form AQ-10.

Similarly, the present study was limited by its use of the TAS-R to measure alexithymia. Although the TAS-R showed good reliability and validity in its initial development (Taylor et al., 1992), it was revised and subsequently followed by the creation of the 20-item Toronto Alexithymia Scale (TAS-20; Bagby et al., 1994), which had improved psychometric properties. In contrast to the TAS-R, this newer scale has also been much more commonly used to measure alexithymia in empirical research (Bagby et al., 2020; Kooiman et al., 2002), making it more difficult to draw comparisons between the results of this study and many previous studies of alexithymia measured with the TAS-20. Unlike the reasons that motivated our use of the AQ-10, the reason that we used the TAS-R rather than the more commonly used and psychometrically improved TAS-20 was unintentional: When adding the questions for alexithymia to the survey software that was used to create this study, the Taylor et al. (1992) article was accidentally open and referenced instead of the correct Bagby et al. (1994) article. Future research should employ the full form AQ and the TAS-20, perhaps in addition to other measures of autistic traits and alexithymia, to test whether our findings can be replicated and whether they are robust.

A final limitation of the study concerns the facial emotion recognition tests. As previously mentioned, the mean accuracy score for emotion recognition of human faces was higher than the mean accuracy score for emotion recognition of anime faces, although perfect scores were earned by participants in both conditions. Anime is not an especially common interest, and participants showed moderate variation on their frequency of

anime or manga use. Thus, it is possible that participants found it more difficult on average to interpret the expressions of the anime stimuli, regardless of whether they are high on autistic traits or alexithymia, due to a lack of familiarity with this style of media. We controlled for frequency of anime or manga use in a series of hierarchical multiple regression models, however, and found no evidence that this frequency was relevant to the associations between autistic traits or alexithymia and anime facial emotion recognition. Alternatively, participants might have found the emotional expressions of anime faces more difficult on average because the anime stimuli were not previously validated like the human stimuli. While we were not able to include a pilot study to ensure that our selected images accurately depicted different facial emotional expressions, we hope that these results encourage future researchers to employ a more rigorous selection process for anime faces to be used in facial emotion recognition tasks.

To improve on future studies examining the facial emotion recognition of anime characters, it might be helpful to create a validated collection of anime faces with different emotional expressions, comparable to the Warsaw Set of Emotional Facial Expression Pictures (Olszanowski et al., 2015). This project might involve commissioning one or more sufficiently skilled artists to draw multiple original anime characters expressing a variety of facial emotions. Alternatively, one could also generate stimuli of anime faces expressing different emotions through the responsible use of artificial intelligence. These images would then need to be examined by a panel of independent judges to determine their validity and effectiveness in conveying the different emotions. Further research could also investigate whether it would be preferable to have images drawn in a consistent visual style, or in different styles to help encompass the artistic variety found in anime.

The most recent data from the Centers for Disease Control indicate that one in 54 children in the United States are diagnosed with ASC (Maenner et al., 2020). Thus, it is important for research to examine what might affect or attenuate the social deficits experienced by those on the autism spectrum or with autistic traits, such that more effective interventions can be developed. The current findings inform future interventions in at least two ways. First, our data suggest that individuals on the autism spectrum might benefit from interventions that specifically target alexithymia, because this subclinical trait was implicated as a potential mechanism through which autistic individuals experience deficits in facial emotion recognition. Second, our data suggest that interventions directed at improving social-emotional functioning in autistic populations might consider the use of anime rather than human characters to improve accessibility and effectiveness.

Supplementary material. The supplementary material for this article can be found at <https://dx.doi.org/10.1017/S0954579425000100>.

Acknowledgements. This research was supported by funding from the Abington College Undergraduate Research Activities (ACURA). We thank Meghan M. Gillen for reviewing an early version of the manuscript and providing helpful feedback. The complete set of images depicting human and anime faces used in the present study can be found at the following link: <https://osf.io/qyzgd>.

Funding statement. This research was supported by funding from the ACURA.

Competing interests. The authors declare none.

References

- Actis-Grosso, R., Bossi, F., & Ricciardelli, P. (2015). Emotion recognition through static faces and moving bodies: A comparison between typically developed adults and individuals with high level of autistic traits. *Frontiers in Psychology*, 6, 1570. <https://doi.org/10.3389/fpsyg.2015.01570>
- Allison, A. (2006). *Millennial monsters: Japanese toys and the global imagination*. University of California Press.
- Allison, C., Auyeung, B., & Baron-Cohen, S. (2012). Toward brief, red flags, for autism screening: The short autism spectrum quotient and the short quantitative checklist in 1,000 cases and 3,000 controls. *Journal of the American Academy of Child and Adolescent Psychiatry*, 51(2), 202–212. <https://doi.org/10.1016/j.jaac.2011.11.003>
- American Psychiatric Association. (2013). *Diagnostic and statistical manual of mental disorders* (5th ed.). American Psychiatric Association.
- Ashwin, E. S., Chapman, E., Colle, L., & Baron-Cohen, S. (2006). Impaired recognition of negative basic emotions in autism: A test of the amygdala theory. *Social Neuroscience*, 1(3-4), 349–363. <https://doi.org/10.1080/17470910601040772>
- Atherton, G., & Cross, L. (2018). Seeing more than human: Autism and anthropomorphic theory of mind. *Frontiers in Psychology*, 9, 528. <https://doi.org/10.3389/fpsyg.2018.00528>
- Atherton, G., & Cross, L. (2019). Animal faux pas: Two legs good four legs bad for theory of mind, but not in the broad autism spectrum. *Journal of Genetic Psychology*, 180(2-3), 81–95. <https://doi.org/10.1080/00221325.2019.1593100>
- Atherton, G., & Cross, L. (2022). Reading the mind in cartoon eyes: Comparing human versus cartoon emotion recognition in those with high and low levels of autistic traits. *Psychological Reports*, 125(3), 1380–1396. <https://doi.org/10.1177/0033294120988135>
- Atherton, G., Morimoto, Y., Nakashima, S., & Cross, L. (2023). Does the study of culture enrich our understanding of autism? A cross-cultural exploration of life on the spectrum in Japan and the West. *Journal of Cross-Cultural Psychology*, 54(5), 610–634. <https://doi.org/10.1177/00220221231169945>
- Bagby, R. M., Parker, J. D. A., & Taylor, G. J. (1994). The twenty-item Toronto Alexithymia Scale: I. Item selection and cross-validation of the factor structure. *Journal of Psychosomatic Research*, 38(1), 23–32. [https://doi.org/10.1016/0022-3999\(94\)90005-1](https://doi.org/10.1016/0022-3999(94)90005-1)
- Bagby, R. M., Parker, J. D. A., & Taylor, G. J. (2020). Twenty-five years with the 20-item Toronto Alexithymia Scale. *Journal of Psychosomatic Research*, 131, 109940. <https://doi.org/10.1016/j.jpsychores.2020.109940>
- Baron, R. M., & Kenny, D. A. (1986). The moderator-mediator variable distinction in social psychological research: Conceptual, strategic, and statistical considerations. *Journal of Personality and Social Psychology*, 51(6), 1173–1182. <https://doi.org/10.1037/0022-3514.51.6.1173>
- Baron-Cohen, S., Jolliffe, T., Mortimore, C., & Robertson, M. (1997). Another advanced test of theory of mind: Evidence from very high functioning adults with autism or Asperger syndrome. *Journal of Child Psychology and Psychiatry*, 38(7), 813–822. <https://doi.org/10.1111/j.1469-7610.1997.tb01599.x>
- Baron-Cohen, S., Wheelwright, S., Skinner, R., Martin, J., & Clubley, E. (2001). The Autism-Spectrum Quotient (AQ): Evidence from Asperger syndrome/high-functioning autism, males and females, scientists and mathematicians. *Journal of Autism and Developmental Disorders*, 31(1), 5–17. <https://doi.org/10.1023/A:1005653411471>
- Bernhardt, B. C., Valk, S. L., Silani, G., Bird, G., Frith, U., & Singer, T. (2014). Selective disruption of sociocognitive structural brain networks in autism and alexithymia. *Cerebral Cortex*, 24(12), 3258–3267. <https://doi.org/10.1093/cercor/bht182>
- Berthoz, S., & Hill, E. L. (2005). The validity of using self-reports to assess emotion regulation abilities in adults with autism spectrum disorder. *European Psychiatry*, 20(3), 291–298. <https://doi.org/10.1016/j.eurpsy.2004.06.01>
- Berthoz, S., Lalanne, C., Crane, L., & Hill, E. L. (2013). Investigating emotional impairments in adults with autism spectrum disorders and the broader autism phenotype. *Psychiatry Research*, 208(3), 257–264. <https://doi.org/10.1016/j.psychres.2013.05.014>
- Berthoz, S., Pouga, L., & Wessa, M. (2011). Alexithymia from the social neuroscience perspective. In J. Decety, & J. T. Cacioppo (Eds.), *The Oxford handbook of social neuroscience* (pp. 907–934). Oxford Library of Psychology. <https://doi.org/10.1093/oxfordhb/9780195342161.013.0060>
- Bertrams, A., & Schlegel, K. (2020). Speeded reasoning moderates the inverse relationship between autistic traits and emotion recognition. *Autism*, 24(8), 2304–2309. <https://doi.org/10.1177/1362361320937090>
- Bird, G., & Cook, R. (2013). Mixed emotions: The contribution of alexithymia to the emotional symptoms of autism. *Translational Psychiatry*, 3(7), e285. <https://doi.org/10.1038/tp.2013.61>
- Black, M. H., Chen, N. T. M., Iyer, K. K., Lipp, O. V., Bölte, S., Falkmer, M., Tan, T., & Girdler, S. (2017). Mechanisms of facial emotion recognition in autism spectrum disorders: Insights from eye tracking and electroencephalography. *Neuroscience and Biobehavioral Reviews*, 80, 488–515.
- Brett, J. D., & Maybery, M. T. (2022). Understanding oneself to understand others: The role of alexithymia and anxiety in the relationships between autistic trait dimensions and empathy. *Journal of Autism and Developmental Disorders*, 52(5), 1971–1983. <https://doi.org/10.1007/s10803-021-05086-6>
- Brosnan, M., Johnson, H., Grawmeyer, B., Chapman, E., & Benton, L. (2015). Emotion recognition in animated compared to human stimuli in adolescents with autism spectrum disorder. *Journal of Autism and Developmental Disorders*, 45(6), 1785–1796. <https://doi.org/10.1007/s10803-014-2338-9>
- Buhrmester, M. D., Talaifar, S., & Gosling, S. D. (2018). An evaluation of Amazon's Mechanical Turk, its rapid rise, and its effective use. *Perspectives on Psychological Science*, 13(2), 149–154. <https://doi.org/10.1177/1745691617706516>
- Celani, G., Battacchi, M. W., & Arcidiacono, L. (1999). The understanding of the emotional meaning of facial expressions in people with autism. *Journal of Autism and Developmental Disorders*, 29(1), 57–66. <https://doi.org/10.1023/a:1025970600181>
- Chevallier, C., Grèzes, J., Molesworth, C., Berthoz, S., & Happé, F. (2012). Brief report: Selective social anhedonia in high functioning autism. *Journal of Autism and Developmental Disorders*, 42(7), 1504–1509. <https://doi.org/10.1007/s10803-011-1364-0>
- Chmielewski, M., & Kucker, S. C. (2020). An MTurk crisis? Shifts in data quality and the impact on study results. *Social Psychological and Personality Science*, 11(4), 464–473. <https://doi.org/10.1177/1948550619875149>
- Clark, T. F., Winkielman, P., & McIntosh, D. N. (2008). Autism and the extraction of emotion from briefly presented facial expressions: Stumbling at the first step of empathy. *Emotion*, 8(6), 803–809. <https://doi.org/10.1037/a0014124>
- Collins, L. M., Graham, J. J., & Flaherty, B. P. (1998). An alternative framework for defining mediation. *Multivariate Behavioral Research*, 33(2), 295–312.
- Cook, R., Brewer, R., Shah, P., & Bird, G. (2013). Alexithymia, not autism, predicts poor recognition of emotional facial expressions. *Psychological Science*, 24(5), 723–732. <https://doi.org/10.1177/0956797612463582>
- Cross, L., Farha, M., & Atherton, G. (2019). The animal in me: Enhancing emotion recognition in adolescents with autism using animal filters. *Journal of Autism and Developmental Disorders*, 49(11), 4482–4487. <https://doi.org/10.1007/s10803-019-04179-7>
- Cross, L., Piovesan, A., & Atherton, G. (2022). Autistic people outperform neurotypicals in a cartoon version of the Reading the Mind in the Eyes. *Autism Research*, 15(9), 1603–1608. <https://doi.org/10.1002/aur.2782>
- Cuve, H. C., Castiello, S., Shiferaw, B., Ichijo, E., Catmur, C., & Bird, G. (2021). Alexithymia explains atypical spatiotemporal dynamics of eye gaze in autism. *Cognition*, 212, 104710. <https://doi.org/10.1016/j.cognition.2021.104710>
- Cuve, H. C., Gao, Y., & Fuse, A. (2018). Is it avoidance or hypoarousal? A systematic review of emotion recognition, eye-tracking, and psychophysiological studies in young adults with autism spectrum conditions. *Research in Autism Spectrum Disorders*, 55, 1–13. <https://doi.org/10.1016/j.jrasd.2018.07.002>
- Cuve, H. C., Murphy, J., Hobson, H., Ichijo, E., Catmur, C., & Bird, G. (2022). Are autistic and alexithymic traits distinct? A factor-analytic and network approach. *Journal of Autism and Developmental Disorders*, 52(5), 2019–2034. <https://doi.org/10.1007/s10803-021-05094-6>
- Dawson, G., Webb, S. J., & McPartland, J. (2005). Understanding the nature of face processing impairment in autism: Insights from behavioral and

- electrophysiological studies. *Developmental Neuropsychology*, 27(3), 403–424. https://doi.org/10.1207/s15326942dn2703_6
- Ekman, P., & Friesen, W. V.** (1971). Constants across cultures in the face and emotion. *Journal of Personality and Social Psychology*, 17(2), 124–129. <https://doi.org/10.1037/h0030377>
- Forby, L., Anderson, N. C., Cheng, J. T., Foulsham, T., Karstadt, B., Dawson, J., Pazhoohi, F., & Kingstone, A.** (2023). Reading the room: Autistic traits, gaze behaviour, and the ability to infer social relationships. *PLoS ONE*, 18(3), e0282310. <https://doi.org/10.1371/journal.pone.0282310>
- Gaigg, S. B.** (2012). The interplay between emotion and cognition in autism spectrum disorder: Implications for developmental theory. *Frontiers in Integrative Neuroscience*, 6, 113. <https://doi.org/10.3389/fnint.2012.00113>
- Gaigg, S. B., Cornell, A. S., & Bird, G.** (2018). The psychophysiological mechanisms of alexithymia in autism spectrum disorder. *Autism*, 22(2), 227–231. <https://doi.org/10.1177/1362361316667062>
- Golan, O., Baron-Cohen, S., & Hill, J.** (2006). The Cambridge Mindreading (CAM) Face-Voice Battery: Testing complex emotion recognition in adults with and without Asperger syndrome. *Journal of Autism and Developmental Disorders*, 36(2), 169–183. <https://doi.org/10.1007/s10803-005-0057-y>
- Gollwitzer, A., Martel, C., McPartland, J. C., & Bargh, J. A.** (2019). Autism spectrum traits predict higher social psychological skill. *Proceedings of the National Academy of Sciences of the United States of America*, 116(39), 19245–19247. <https://doi.org/10.1073/pnas.1911460116>
- Grynberg, D., Chang, B., Corneille, O., Maurage, P., Vermeulen, N., Berthoz, S., & Luminet, O.** (2012). Alexithymia and the processing of emotional facial expressions (EFEs): Systematic review, unanswered questions and further perspectives. *PLoS ONE*, 7(8), e42429. <https://doi.org/10.1371/journal.pone.0042429>
- Guillon, Q., Rogé, B., Afzali, M. H., Baduel, S., Kruck, J., & Hadjikhani, N.** (2016). Intact perception but abnormal orientation towards face-like objects in young children with ASD. *Scientific Reports*, 6, 22119. <https://doi.org/10.1038/srep22119>
- Harms, M. B., Martin, A., & Wallace, G. L.** (2010). Facial emotion recognition in autism spectrum disorders: A review of behavioral and neuroimaging studies. *Neuropsychology Review*, 20(3), 290–322. <https://doi.org/10.1007/s11065-010-9138-6>
- Heaton, P., Reichenbacher, L., Sauter, D., Allen, R., Scott, S., & Hill, E.** (2012). Measuring the effects of alexithymia on perception of emotional vocalizations in autistic spectrum disorder and typical development. *Psychological Medicine*, 42(11), 2453–2459. <https://doi.org/10.1017/S0033291712000621>
- Hill, E., Berthoz, S., & Frith, U.** (2004). Brief report: Cognitive processing of own emotions in individuals with autistic spectrum disorder and in their relatives. *Journal of Autism and Developmental Disorders*, 34(2), 229–235. <https://doi.org/https://doi.org>
- Hobson, R. P., Ouston, J., & Lee, A.** (1989). Naming emotion in faces and voices: Abilities and disabilities in autism and mental retardation. *British Journal of Developmental Psychology*, 7, 237–250. <https://doi.org/10.1111/j.2044-835X.1989.tb00803.x>
- Ingersoll, B.** (2010). Broader autism phenotype and nonverbal sensitivity: Evidence for an association in the general population. *Journal of Autism and Developmental Disorders*, 40(5), 590–598. <https://doi.org/10.1007/s10803-009-0907-0>
- Izard, C. E.** (1994). Innate and universal facial expressions: Evidence from developmental and cross-cultural research. *Psychological Bulletin*, 115(2), 288–299. <https://doi.org/10.1037/0033-2909.115.2.288>
- Izard, C. E., Kagan, J., & Zajonc, R. B.** (1984). *Emotions, cognition, and behavior*. Cambridge University Press.
- Jia, R., Steelman, Z. R., & Jia, H. H.** (2019). Psychometric assessments of three self-report autism scales (AQ, RBQ-2A, and SQ) for general adult populations. *Journal of Autism and Developmental Disorders*, 49(5), 1949–1965. <https://doi.org/10.1007/s10803-019-03880-x>
- Jones, C. R., Pickles, A., Falcato, M., Marsden, A. J., Happé, F., Scott, S. K., Sauter, D., Tregay, J., Phillips, R. J., Baird, G., Simonoff, E., & Charman, T.** (2011). A multimodal approach to emotion recognition ability in autism spectrum disorders. *Journal of Child Psychology and Psychiatry*, 52(3), 275–285. <https://doi.org/10.1111/j.1469-7610.2010.02328.x>
- Kinnaird, E., Stewart, C., & Tchanturia, K.** (2019). Investigating alexithymia in autism: A systematic review and meta-analysis. *European Psychiatry*, 55, 80–89. <https://doi.org/10.1016/j.eurpsy.2018.09.004>
- Klin, A., Jones, W., Schultz, R., Volkmar, F., & Cohen, D.** (2002). Visual fixation patterns during viewing of naturalistic social situations as predictors of social competence in individuals with autism. *Archives of General Psychiatry*, 59(9), 809–816. <https://doi.org/10.1001/archpsyc.59.9.809>
- Kooiman, C. G., Spinhoven, P., & Trijsburg, R. W.** (2002). The assessment of alexithymia: A critical review of the literature and a psychometric study of the Toronto Alexithymia Scale-20. *Journal of Psychosomatic Research*, 53(6), 1083–1090. [https://doi.org/10.1016/S0022-3999\(02\)00348-3](https://doi.org/10.1016/S0022-3999(02)00348-3)
- Kuo, M. H., Orsmond, G. I., Coster, W. J., & Cohn, E. S.** (2014). Media use among adolescents with autism spectrum disorder. *Autism*, 18(8), 914–923. <https://doi.org/10.1177/1362361313497832>
- Linden, W., Wen, F., & Paulhus, D. L.** (1995). Measuring alexithymia: Reliability, validity, and prevalence. In J. N. Butcher, & C. D. Spielberger (Eds.), *Advances in personality assessment* (pp. 51–95). Lawrence Erlbaum Associates.
- Liu, K., Chen, J. H., & Chang, K. M.** (2019). A study of facial features of American and Japanese cartoon characters. *Symmetry*, 11(5), 664. <https://doi.org/10.3390/sym11050664>
- Macdonald, H., Rutter, M., Howlin, P., Rios, P., Le Conteur, A., Evered, C., & Folstein, S.** (1989). Recognition and expression of emotional cues by autistic and normal adults. *Journal of Child Psychology and Psychiatry*, 30(6), 865–877. <https://doi.org/10.1111/j.1469-7610.1989.tb00288.x>
- Maenner, M. J., Shaw, K. A., Baio, J., Washington, A., Patrick, M., DiRienzo, M., Christensen, D. L., Wiggins, L. D., Pettygrove, S., Andrews, J. G., Lopez, M., Hudson, A., Baroud, T., Schwenk, Y., White, T., Rosenberg, C. R., Lee, L. C., Harrington, R. A., Huston, M., & Dietz, P. M.** (2020). Prevalence of autism spectrum disorder among children aged 8 years — Autism and developmental disabilities monitoring network, 11 sites, United States, 2016. *MMWR Surveillance Summaries*, 69(4), 1–12. <https://doi.org/10.15585/mmwr.mm6904a1>
- Mason, D., Ronald, A., Ambler, A., Caspi, A., Houts, R., Poulton, R., Ramrakha, S., Wertz, J., Moffitt, T. E., & Happé, F.** (2021). Autistic traits are associated with faster pace of aging: Evidence from the Dunedin study at age 45. *Autism Research*, 14(8), 1684–1694. <https://doi.org/10.1002/aur.2534>
- Matsumoto, D.** (2001). Culture and emotion. In D. Matsumoto (Ed.), *The handbook of culture and psychology* (pp. 171–194). Oxford University Press.
- Mazurek, M. O., Shattuck, P. T., Wagner, M., & Cooper, B. P.** (2012). Prevalence and correlates of screen-based media use among youths with autism spectrum disorders. *Journal of Autism and Developmental Disorders*, 42(8), 1757–1767. <https://doi.org/10.1007/s10803-011-1413-8>
- McKenzie, K., Murray, A. L., Wilkinson, A., Murray, G. C., Metcalfe, D., O'Donnell, M., & McCarty, K.** (2018). The relations between processing style, autistic-like traits, and emotion recognition in individuals with and without autism spectrum disorder. *Personality and Individual Differences*, 120, 1–6. <https://doi.org/10.1016/j.paid.2017.08.007>
- Milosavljevic, B., Carter Leno, V., Simonoff, E., Baird, G., Pickles, A., Jones, C. R., Erskine, C., Charman, T., & Happé, F.** (2016). Alexithymia in adolescents with autism spectrum disorder: Its relationship to internalising difficulties, sensory modulation and social cognition. *Journal of Autism and Developmental Disorders*, 46(4), 1354–1367. <https://doi.org/10.1007/s10803-015-2670-8>
- Miyahara, M., Bray, A., Tsujii, M., Fujita, C., & Sugiyama, T.** (2007). Reaction time of facial affect recognition in Asperger's disorder for cartoon and real, static and moving faces. *Child Psychiatry and Human Development*, 38(2), 121–134. <https://doi.org/10.1007/s10578-007-0048-7>
- Moriguchi, Y., Ohnishi, T., Lane, R. D., Maeda, M., Mori, T., Nemoto, K., Matsuda, H., & Komaki, G.** (2006). Impaired self-awareness and theory of mind: An fMRI study of mentalizing in alexithymia. *Neuroimage*, 32(3), 1472–1482. <https://doi.org/10.1016/j.neuroimage.2006.04.186>
- Moss, A. J., Rosenzweig, C., Robinson, J., Jaffe, S. N., & Litman, L.** (2023). Is it ethical to use Mechanical Turk for behavioral research? Relevant data from a representative survey of MTurk participants and wages. *Behavior Research Methods*, 55(8), 4048–4067. <https://doi.org/10.3758/s13428-022-02005-0>
- Napier, S.** (2006). The world of anime fandom in America. *Mechademia*, 1(1), 47–63. <https://doi.org/10.1353/mec.0.0072>

- Nemiah, J. C., Freyberger, H., & Sifneos, P. E. (1976). Alexithymia: A view of the psychosomatic process. In O. W. Hill (Eds.), *Modern trends in psychosomatic medicine* (Vol. 3, pp. 430–439). Butterworths.
- Newman, A., Bavik, Y. L., Mount, M., & Shao, B. (2021). Data collection via online platforms: Challenges and recommendations for future research. *Applied Psychology: An International Review*, 70(3), 1380–1402. <https://doi.org/10.1111/apps.12302>
- Oakley, B. F. M., Brewer, R., Bird, G., & Catmur, C. (2016). Theory of mind is not theory of emotion: A cautionary note on the Reading the Mind in the Eyes Test. *Journal of Abnormal Psychology*, 125(6), 818–823. <https://doi.org/10.1037/abn0000182>
- Ola, L., & Gullon-Scott, F. (2020). Facial emotion recognition in autistic adult females correlates with alexithymia, not autism. *Autism*, 24(8), 2021–2034. <https://doi.org/10.1177/1362361320932727>
- Olszanowski, M., Pochwatko, G., Kuklinski, K., Scibor-Rylski, M., Lewinski, P., & Ohme, R. K. (2015). Warsaw set of emotional facial expression pictures: A validation study of facial display photographs. *Frontiers in Psychology*, 5, 1516. <https://doi.org/10.3389/fpsyg.2014.01516>
- Otmazgin, N. (2014). Anime in the US: The entrepreneurial dimensions of globalized culture. *Pacific Affairs*, 87(1), 53–69. <https://doi.org/10.5509/201487153>
- Paolacci, G., & Chandler, J. (2014). Inside the Turk: Understanding Mechanical Turk as a participant pool. *Current Directions in Psychological Science*, 23(3), 184–188. <https://doi.org/10.1177/0963721414531598>
- Passarelli, M., Masini, M., Bracco, F., Petrosino, M., & Chiorri, C. (2018). Development and validation of the Facial Expression Recognition Test (FERT). *Psychological Assessment*, 30(11), 1479–1490. <https://doi.org/10.1037/pas0000595>
- Pazhoohi, F., Forby, L., & Kingstone, A. (2021). Facial masks affect emotion recognition in the general population and individuals with autistic traits. *PLoS ONE*, 16(9), e0257740. <https://doi.org/10.1371/journal.pone.0257740>
- Pelphrey, K. A., Sasson, N. J., Reznick, J. S., Paul, G., Goldman, B. D., & Piven, J. (2002). Visual scanning of faces in autism. *Journal of Autism and Developmental Disorders*, 32(4), 249–261. <https://doi.org/10.1023/a:1016374617369>
- Philip, R. C. M., Whalley, H. C., Stanfield, A. C., Sprengelmeyer, R., Santos, I. M., Young, A. W., Atkinson, A. P., Calder, A. J., Johnstone, E. C., Lawrie, S. M., & Hall, J. (2010). Deficits in facial, body movement and vocal emotional processing in autism spectrum disorders. *Psychological Medicine*, 40(11), 1919–1929. <https://doi.org/10.1017/S0033291709992364>
- Piggot, J., Kwon, H., Mobbs, D., Blasey, C., Lotspeich, L., Menon, V., Bookheimer, S., & Reiss, A. L. (2004). Emotional attribution in high-functioning individuals with autistic spectrum disorder: A functional imaging study. *Journal of the American Academy of Child and Adolescent Psychiatry*, 43(4), 473–480. <https://doi.org/10.1097/00004583-200404000-00014>
- Poljac, E., Poljac, E., & Wagemans, J. (2013). Reduced accuracy and sensitivity in the perception of emotional facial expressions in individuals with high autism spectrum traits. *Autism*, 17(6), 668–680. <https://doi.org/10.1177/1362361312455703>
- Poquérousse, J., Pastore, L., Dellantonio, S., & Esposito, G. (2018). Alexithymia and autism spectrum disorder: A complex relationship. *Frontiers in Psychology*, 9, 1196. <https://doi.org/10.3389/fpsyg.2018.01196>
- Preacher, K. J., & Hayes, A. F. (2004). SPSS and SAS procedures for estimating indirect effects in simple mediation models. *Behavior Research Methods, Instruments & Computers*, 36(4), 717–731. <https://doi.org/10.3758/BF03206553>
- Quattrocki, E., & Friston, K. (2014). Autism, oxytocin and interoception. *Neuroscience and Biobehavioral Reviews*, 47, 410–430. <https://doi.org/10.1016/j.neubiorev.2014.09.012>
- Rosset, D. B., Rondan, C., Da Fonseca, D., Santos, A., Assouline, B., & Deruelle, C. (2008). Typical emotion processing for cartoon but not for real faces in children with autistic spectrum disorders. *Journal of Autism and Developmental Disorders*, 38(5), 919–925. <https://doi.org/10.1007/s10803-007-0465-2>
- Rozema, R. (2015). Manga and the autistic mind. *English Journal*, 105(1), 60–68.
- Rucker, D. D., Preacher, K. J., Tormala, Z. L., & Petty, R. E. (2011). Mediation analysis in social psychology: Current practices and new recommendations. *Social and Personality Psychology Compass*, 5(6), 359–371. <https://doi.org/10.1111/j.1751-9004.2011.00355.x>
- Rudolph, C. E. S., Lundin, A., Åhs, J. W., Dalman, C., & Kosidou, K. (2018). Brief report: Sexual orientation in individuals with autistic traits: Population based study of 47,000 adults in Stockholm county. *Journal of Autism and Developmental Disorders*, 48(2), 619–624. <https://doi.org/10.1007/s10803-017-3369-9>
- Rump, K. M., Giovannelli, J. L., Minshew, N. J., & Strauss, M. S. (2009). The development of emotion recognition in individuals with autism. *Child Development*, 80(5), 1434–1447. <https://doi.org/10.1111/j.1467-8624.2009.01343.x>
- Ruzich, E., Allison, C., Smith, P., Watson, P., Auyeung, B., Ring, H., & Baron-Cohen, S. (2016). Subgrouping siblings of people with autism: Identifying the broader autism phenotype. *Autism Research*, 9(6), 658–665. <https://doi.org/10.1002/aur.1544>
- Santiesteban, I., Gibbard, C., Drucks, H., Clayton, N., Banissy, M. J., & Bird, G. (2021). Individuals with autism share others' emotions: Evidence from the Continuous Affective Rating and Empathic Responses (CARER) task. *Journal of Autism and Developmental Disorders*, 51(2), 391–404. <https://doi.org/10.1007/s10803-020-04535-y>
- Shane, H. C., & Albert, P. D. (2008). Electronic screen media for persons with autism spectrum disorders: Results of a survey. *Journal of Autism and Developmental Disorders*, 38(8), 1499–1508. <https://doi.org/10.1007/s10803-007-0527-5>
- Shrout, P. E., & Bolger, N. (2002). Mediation in experimental and nonexperimental studies: New procedures and recommendations. *Psychological Methods*, 7(4), 422–445. <https://doi.org/10.1037/1082-989X.7.4.422>
- Silani, G., Bird, G., Brindley, R., Singer, T., Frith, C., & Frith, U. (2008). Levels of emotional awareness and autism: An fMRI study. *Social Neuroscience*, 3(2), 97–112. <https://doi.org/10.1080/17470910701577020>
- Silva, C., Da Fonseca, D., Esteves, F., & Deruelle, C. (2015). Motivational approach and avoidance in autism spectrum disorder: A comparison between real photographs and cartoons. *Research in Autism Spectrum Disorders*, 17, 13–24. <https://doi.org/10.1016/j.rasd.2015.05.004>
- Sizoo, B. B., Horwitz, E. H., Teunisse, J. P., Kan, C. C., Vissers, C. T. W., Forceville, E. J. M., Van Voorst, A. J. P., & Geurts, H. M. (2015). Predictive validity of self-report questionnaires in the assessment of autism spectrum disorders in adults. *Autism*, 19(7), 842–849. <https://doi.org/10.1177/1362361315589869>
- Song, Y., & Hakoda, Y. (2018). Selective impairment of basic emotion recognition in people with autism: Discrimination thresholds for recognition of facial expressions of varying intensities. *Journal of Autism and Developmental Disorders*, 48(6), 1886–1894. <https://doi.org/10.1007/s10803-017-3428-2>
- South, M., Ozonoff, S., & McMahon, W. M. (2005). Repetitive behavior profiles in Asperger syndrome and high-functioning autism. *Journal of Autism and Developmental Disorders*, 35(2), 145–158. <https://doi.org/10.1007/s10803-004-1992-8>
- Stuart, N., Whitehouse, A., Palermo, R., Bothe, E., & Badcock, N. (2023). Eye gaze in autism spectrum disorder: A review of neural evidence for the eye avoidance hypothesis. *Journal of Autism and Developmental Disorders*, 53(5), 1884–1905. <https://doi.org/10.1007/s10803-022-05443-z>
- Sucksmith, E., Allison, C., Baron-Cohen, S., Chakrabarti, B., & Hoekstra, R. A. (2013). Empathy and emotion recognition in people with autism, first-degree relatives, and controls. *Neuropsychologia*, 51(1), 98–105. <https://doi.org/10.1016/j.neuropsychologia.2012.11.013>
- Taylor, E. C., Livingston, L. A., Clutterbuck, R. A., & Shah, P. (2020). Psychometric concerns with the 10-item Autism-Spectrum Quotient (AQ10) as a measure of trait autism in the general population. *Experimental Results*, 1, e3. <https://doi.org/10.1017/exp.2019.3>
- Taylor, G. J., Bagby, R. M., & Parker, J. D. (1992). The Revised Toronto Alexithymia Scale: Some reliability, validity, and normative data. *Psychotherapy and Psychosomatics*, 57(1-2), 34–41. <https://doi.org/10.1159/000288571>

- Taylor, G. J., Bagby, R. M., & Parker, J. D.** (1997). *Disorders of affect regulation: Alexithymia in medical and psychiatric illness*. Cambridge University Press. <https://doi.org/10.1017/CBO9780511526831>
- Tottenham, N., Hertzog, M. E., Gillespie-Lynch, K., Gilhooly, T., Millner, A. J., & Casey, B. J.** (2014). Elevated amygdala response to faces and gaze aversion in autism spectrum disorder. *Social Cognitive and Affective Neuroscience*, 9(1), 106–117. <https://doi.org/10.1093/scan/nst050>
- Tracy, J. L., Robins, R. W., Schriber, R. A., & Solomon, M.** (2011). Is emotion recognition impaired in individuals with autism spectrum disorders? *Journal of Autism and Developmental Disorders*, 41(1), 102–109. <https://doi.org/10.1007/s10803-010-1030-y>
- Trevisan, D. A., Bowering, M., & Birmingham, E.** (2016). Alexithymia, but not autism spectrum disorder, may be related to the production of emotional facial expressions. *Molecular Autism*, 7(1), 46. <https://doi.org/10.1186/s13229-016-0108-6>
- Uljarevic, M., & Hamilton, A.** (2013). Recognition of emotions in autism: A formal meta-analysis. *Journal of Autism and Developmental Disorders*, 43(7), 1517–1526. <https://doi.org/10.1007/s10803-012-1695-5>
- Vettori, S., Dzhelyova, M., Van der Donck, S., Jacques, C., Van Wesemael, T., Steyaert, J., Rossion, B., & Boets, B.** (2020). Combined frequency-tagging EEG and eye tracking reveal reduced social bias in boys with autism spectrum disorder. *Cortex*, 125, 135–148. <https://doi.org/10.1016/j.cortex.2019.12.013>
- Wallace, S., Coleman, M., & Bailey, A.** (2008). An investigation of basic facial expression recognition in autism spectrum disorders. *Cognition and Emotion*, 22(7), 1353–1380. <https://doi.org/10.1080/0269930701782153>
- Webb, M. A., & Tangney, J. P.** (2024). Too good to be true: Bots and bad data from Mechanical Turk. *Perspectives on Psychological Science*, 19(6), 887–890. <https://doi.org/10.1177/17456916221120027>
- Wilhelm, O., Hildebrandt, A., Manske, K., Schacht, A., & Sommer, W.** (2014). Test battery for measuring the perception and recognition of facial expressions of emotion. *Frontiers in Psychology*, 5, 404. <https://doi.org/10.3389/fpsyg.2014.0040>
- Yeung, M. K.** (2022). A systematic review and meta-analysis of facial emotion recognition in autism spectrum disorder: The specificity of deficits and the role of task characteristics. *Neuroscience and Biobehavioral Reviews*, 133, 104518. <https://doi.org/10.1016/j.neubiorev.2021.104518>
- Zhao, X., Lynch, J. G., Jr., & Chen, Q.** (2010). Reconsidering Baron and Kenny: Myths and truths about mediation analysis. *Journal of Consumer Research*, 37(2), 197–206. <https://doi.org/10.1086/651257>