INSTRUCTIONS FOR CONTRIBUTORS

SUBMISSION OF MANUSCRIPTS

Manuscripts should be submitted online via our manuscript submission and tracking site, http://www.editorialmanager.com/psm/. Full instructions for electronic submission are available directly from this site. To facilitate rapid reviewing, communications for peer review will be electronic and authors will need to supply a current e-mail address when registering to use the system.

Papers for publication from Europe (except those on genetic topics, irrespective of country), and all papers on imaging topics, should be submitted to the UK Office.

Papers from the Americas, Asia, Africa, Australasia and the Middle East (except those dealing with imaging topics), and all papers dealing with genetic topics, irrespective of country, should be sent to US Office.

Generally papers should not have text more than 4500 words in length (excluding these sections) and should not have more than a combined total of 5 tables and/or figures. Papers shorter than these limits are encouraged. For papers of unusual importance the editors may waive these requirements. A structured abstract of no more than 250 words should be given at the beginning of the article using the headings: Background; Methods; Results; Conclusions. The name of an author to whom correspondence should be sent must be indicated and a full postal address given in the footnote. Any acknowledgements should be placed at the end of the text (before the References section).

Declaration of Interest: A statement must be provided in the acknowledgements listing all financial support received for the work and, for all authors, any financial involvement (including employment, fees, share ownership) or affiliation with any organization whose financial interests may be affected by material in the manuscript, or which might potentially bias it. This applies to all papers including editorials and letters to the editor.

Contributors should also note the following:

- 1. S.I. units should be used throughout in text, figures and tables.
- 2. Authors should spell out in full any abbreviations used in their manuscripts.
- 3. Foreign quotations and phrases should be followed by a translation.
- 4. If necessary, guidelines for statistical presentation may be found in: **Altman DG, Gore SM, Gardner MJ & Pocock SJ** (1983). Statistical guidelines for contributors to medical journals. *British Medical Journal* 286, 1489–1493.

REFERENCES (1) The Harvard (author-date) system should be used in the text and a complete list of References cited given at the end of the article. In a text citation of a work by more than two authors cite the first author's name followed by et al. (but the names of all of the authors should be given in the References section). Where several references are cited together they should be listed in rising date order. (2) The References section should be supplied in alphabetical order (authors' names in **bold**, journal titles in full), following the text. Some examples follow:

Miller PM, Byrne M, Hodges A, Lawrie SM, Johnstone EC (2002). Childhood behaviour, psychotic symptoms and psychosis onset in young people at high risk of schizophrenia: early findings from the Edinburgh high risk study. *Psychological Medicine* 32, 173–179.

Cleckley HJ (1941). The Mask of Sanity, 2nd edn. Mosby: St. Louis, MO.

Brewer WJ, Wood SJ, DeLuca C, Pantelis C (2006). Models of olfaction for exploring neurodevelopment. In *Olfaction and the Brain* (ed. W. J. Brewer, D. Castle and C. Pantelis), pp. 97–121. Cambridge University Press: Cambridge.

(3) Online citations

doi (when published online prior to printed issue)

Lauritsen MB, Pedersen CB, Mortensen CB (2004). The incidence and prevalence of pervasive developmental disorders: a Danish population-based study. *Psychological Medicine*. Published online: 21 October 2004. doi:10.1017/S0033291704002387.

URL

World Bank (2003). Quantitative techniques for health equity analysis – Technical Notes (http://siteresources.worldbank. org/INTPAH/Resources/Publications/Quantitative-Techniques/health.eq tn07.pdf). Accessed 15 February 2006.

[Authors are requested to print-out and keep a copy of any online-only material, in case the URL changes or is no longer maintained.]

FIGURES AND TABLES Only essential figures and tables should be included. Further tables, figures, photographs and appendices, may be included with the online version on the journal website. To ensure that your figures are reproduced to the highest possible standards, Cambridge Journals recommends the following formats and resolutions for supplying electronic figures. Please ensure that your figures are saved at final publication size and are in our recommended file formats. Following these guidelines will result in high quality images being reproduced in both the print and the online versions of the journal. Line artwork: Format: tif or eps, Colour mode: black and white (also known as 1-bit), Resolution: 1200 dpi; Combination artwork (line/tone): Format: tif or eps, Colour mode: grayscale (also known as 8-bit), Resolution: 800 dpi; Black and white halftone artwork: Format: tif, Colour mode: grayscale (also known as 8-bit), Resolution: 300 dpi; Colour halftone artwork: Format: tif, Colour mode: CMYK colour, Resolution: 300 dpi. All photographs, graphs, and diagrams should be referred to as figures and should be numbered consecutively in Arabic numerals. Captions for figures should be typed double-spaced on separate sheets. Tables Tables should be typed above the table.

PROOFS AND OFFPRINTS Page proofs will be sent to the author designated to receive correspondence. corrections other than to printer's errors may be charged to the author. The corresponding author of each paper will receive a PDF file of their article and hard copy offprints may be purchased if they are ordered on the form supplied when the proof is returned.

PSYCHOLOGICAL MEDICINE

CONTENTS

ORIGINAL ARTICLES Disrupted resting-state brain network properties in obesity: decreased global and putaminal cortico-striatal network efficiency Baek K, Morris LS, Kundu P and Voon V	585	Longitudinal associations between late-life depression dimensions and cognitive functioning: a cross-domain latent growth curve analysis Brailean A, Aartsen MJ, Muniz-Terrera G, Prince M, Prina AM, Comijs HC, Huisman M and Beekman A	690
A novel resting-state functional magnetic resonance imaging signature of resilience to recurrent depression Workman Cl, Lythe KE, McKie S, Moll J, Gethin JA, Deakin JFW, Elliott R and Zahn R	597	An adaptive randomized trial of dialectical behaviour therapy and cognitive behavior therapy for binge-eating Chen EY, Cacioppo J, Fettich K, Gallop R, McCloskey MS, Olino T and Zeffiro TA	703
Performance characteristics and clinical utility of diagnostic criteria proposals in bereaved treatment-seeking patients Mauro C, Shear MK, Reynolds III CF, Simon NM, Zisook S, Skritskaya N, Wang Y, Lebowitz B, Duan N, First MB, Ghesquiere A, Gribbin C and Glickman K	608	Insight as a social identity process in the evolution of psychosocial functioning in the early phase of psychosis Klaas HS, Clémence A, Marion-Veyron R, Antonietti J-P, Alameda L, Golay P and Conus P Transdiagnostic brain responses to disorder-related threat	718
Cannabis-induced attenuated psychotic symptoms: implications for prognosis in young people at ultra-high risk for psychosis McHugh MJ, McGorry PD, Yung AR, Lin A, Wood SJ,	across four psychiatric disorders Feldker K, Heitmann CY, Neumeister P, Tupak SV, Schramr Moeck R, Zwitserlood P, Bruchmann M and Straube T	across four psychiatric disorders Feldker K, Heitmann CY, Neumeister P, Tupak SV, Schrammen E, Moeck R, Zwitserlood P, Bruchmann M and Straube T	730
Hartmann JA and Nelson B Neighborhood deprivation and depression in adult twins: genetics and genexenvironment interaction Strachan E, Duncan G, Horn E and Turkheimer E	616	Exploring DSM-5 ADHD criteria beyond young adulthood: phenomenology, psychometric properties and prevalence in a large three-decade birth cohort Vitola ES, Bau CHD, Salum GA, Horta BL, Quevedo L, Barros FC, Pinheiro RT, Kieling C, Rohde LA and Grevet EH	744
Shared atypical brain anatomy and intrinsic functional architecture in male youth with autism spectrum disorder and their unaffected brothers Lin H-Y, Tseng W-Y I, Lai M-C, Chang Y-T and Gau SS-F	639	Relationship of amotivation to neurocognition, self-efficacy and functioning in first-episode psychosis: a structural equation modeling approach Chang WC, Kwong WWY, Hui CLM, Chan SKW, Lee EHM and Chen EYH	755
Alcohol use is associated with thinner cerebral cortex and larger ventricles in schizophrenia, bipolar disorder and healthy controls Lange EH, Nerland S, Jørgensen KN, Mørch-Johnsen L, Nesvåg R, Hartberg CB, Haukvik UK, Osnes K, Melle I, Andreassen OA and Agartz I	655	Antenatal depressive symptoms as a predictor of deterioration in perceived social support across the perin period: a four-wave cohort study in Turkey Senturk V, Abas M, Dewey M, Berksun O and Stewart R 655 Educational achievement in psychiatric patients and their	766
Childhood trauma is associated with increased brain responses to emotionally negative as compared with positive faces in patients with psychotic disorders Aas M, Kauppi K, Brandt CL, Tesli M, Kaufmann T, Steen NE, Agartz I, Westlye LT, Andreassen OA and Melle I Maternal depressive symptoms and early childhood cognitive	s N To	siblings: a register-based study in 30 000 individuals in The Netherlands Tempelaar WM, Termorshuizen F, MacCabe JH, Boks MPM and Kahn RS	776
development: a meta-analysis			



680



cambridge.org/psm

Liu Y, Kaaya S, Chai J, McCoy DC, Surkan PJ, Black MM, Sutter-Dallay A-L, Verdoux H and Smith-Fawzi MC

