

AN ESCAPE FROM VARDANYAN'S THEOREM

ANA DE ALMEIDA BORGES  AND JOOST J. JOOSTEN 

Abstract. Vardanyan's Theorems [36, 37] state that $\text{QPL}(\text{PA})$ —the quantified provability logic of Peano Arithmetic—is Π_2^0 complete, and in particular that this already holds when the language is restricted to a single unary predicate. Moreover, Visser and de Jonge [38] generalized this result to conclude that it is impossible to computably axiomatize the quantified provability logic of a wide class of theories. However, the proof of this fact cannot be performed in a strictly positive signature. The system QRC_1 was previously introduced by the authors [1] as a candidate first-order provability logic. Here we generalize the previously available Kripke soundness and completeness proofs, obtaining constant domain completeness. Then we show that QRC_1 is indeed complete with respect to arithmetical semantics. This is achieved via a Solovay-type construction applied to constant domain Kripke models. As corollaries, we see that QRC_1 is the strictly positive fragment of QGL and a fragment of $\text{QPL}(\text{PA})$.

§1. Introduction. Provability is a fundamental concept in mathematics, logic, and philosophy alike. Gödel proved his famous incompleteness results in [22] by formalizing provability. Thus, for formal theories like Peano Arithmetic (PA), we have a natural arithmetical predicate $\Box_{\text{PA}}(\cdot)$ that is true exactly for the Gödel numbers of PA-provable sentences:

$$\text{PA} \vdash A \iff \mathbb{N} \models \Box_{\text{PA}}(\ulcorner A \urcorner). \quad (1)$$

For readability we shall not distinguish formulas from their Gödel numbers or from syntactic terms (numerals) denoting these Gödel numbers in the future.

Gödel observed various provable structural properties of the provability predicate. For example, for any formula A , if $\text{PA} \vdash A$ then $\text{PA} \vdash \Box_{\text{PA}}A$. Moreover, this can be formalized itself: for any formula A , $\text{PA} \vdash \Box_{\text{PA}}A \rightarrow \Box_{\text{PA}}\Box_{\text{PA}}A$. Using such properties, and after showing how self-reference can be obtained in PA, Gödel derived his first incompleteness theorem (here presented in a slightly weakened form for PA) by observing that the sentence B such that $\text{PA} \vdash B \leftrightarrow \neg\Box_{\text{PA}}B$ can neither be proved nor refuted in PA, provided that PA only proves true theorems.

In light of this, it makes sense to design a system that collects all provable structural properties of the provability predicate. The language \mathcal{L}_{\Box} of propositional modal logic is optimally suited for this purpose. The formulas of this language are as the ones of propositional logic together with a unary modality \Box that is syntactically treated as negation. Thus we can write $\text{Form}_{\Box} ::= \perp \mid \text{Prop} \mid \text{Form}_{\Box} \rightarrow \text{Form}_{\Box} \mid \Box\text{Form}_{\Box}$, where Prop represents propositional symbols.

Received August 31, 2021.

2020 *Mathematics Subject Classification.* 03B45, 03B60, 03F03, 03F30, 03F45, 03F55.

Key words and phrases. modal logic, provability logic, strictly positive logics, quantified modal logic, arithmetic interpretations, feasible fragments.

© The Author(s), 2022. Published by Cambridge University Press on behalf of The Association for Symbolic Logic. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted re-use, distribution, and reproduction in any medium, provided the original work is properly cited.

0022-4812/23/8804-0014

DOI:10.1017/jsl.2022.38



The modal logical formulas are linked to arithmetic using so-called *realizations*. A particular realization \cdot^* maps propositional variables to sentences in the language of PA and this map is extended to any formula by stipulating that it commutes with implication, \perp is mapped to $(0 = 1)$, and the modal operator is mapped to formalized provability, i.e., $(\Box A)^{*\text{PA}} = \Box_{\text{PA}} A^{*\text{PA}}$.

We can now express what it means to be a provable structural property. For example, for any realization \cdot^* we have $\text{PA} \vdash (\Box(p \rightarrow q) \rightarrow (\Box p \rightarrow \Box q))^{*\text{PA}}$, since $\text{PA} \vdash \Box_{\text{PA}}(p^* \rightarrow q^*) \rightarrow (\Box_{\text{PA}} p^* \rightarrow \Box_{\text{PA}} q^*)$ for any formulas p^* and q^* . The structural properties are thus captured by the modal formulas that are provable under any realization. We call this $\text{PL}(\text{PA})$, the *provability logic* of PA, and write

$$\text{PL}(\text{PA}) := \{A \in \mathcal{L}_{\Box} \mid \text{for any } \cdot^*, \text{ we have } \text{PA} \vdash A^{*\text{PA}}\}.$$

Via (1) we know that $(\text{PA} \vdash A^{*\text{PA}}$ for any \cdot^*) if and only if $(\mathbb{N} \models \Box_{\text{PA}} A^{*\text{PA}}$ for any \cdot^*). Clearly $A^{*\text{PA}}$ only depends on the value of \cdot^* for the finitely many propositional variables that occur in A , and so the universal quantifier “for any \cdot^* ” can be coded and made internal, making it possible to characterize $\text{PL}(\text{PA})$ by $\{A \mid \mathbb{N} \models \forall \cdot^* \Box_{\text{PA}} A^{*\text{PA}}\}$. We now observe that $\Box_{\text{PA}} B$ is a Σ_1^0 formula. That is, $\Box_{\text{PA}} B$ is of the form $\exists p \text{Prf}_{\text{PA}}(p, B)$, where $\text{Prf}_{\text{PA}}(\cdot, \cdot)$ is a decidable predicate. Thus, $\text{PL}(\text{PA}) = \{A \mid \mathbb{N} \models \forall \cdot^* \Box_{\text{PA}} A^{*\text{PA}}\}$ has a Π_2^0 definition. If we moreover realize that $\Box_{\text{PA}}(\cdot)$ is Σ_1^0 complete in that any computably enumerable¹ set U can be defined using a formula A_U as $\{n \mid \mathbb{N} \models \Box_{\text{PA}}(A_U(n))\}$, then it seems that there is little hope that the Π_2^0 set $\text{PL}(\text{PA})$ allows for a simple characterization.

However, a little miracle happens and by Solovay’s Completeness Theorem [34] we know that $\text{PL}(\text{PA})$ is decidable, corresponding to what we nowadays call the *Gödel–Löb provability logic* GL. Likewise, Solovay proved that the set of *true* structural provability principles

$$\text{TPL}(\text{PA}) := \{\varphi \in \mathcal{L}_{\Box} \mid \text{for any } \cdot^*, \text{ we have } \mathbb{N} \models \varphi^{*\text{PA}}\}$$

is also decidable and described by the well-behaved modal logic GLS. We observe that, *a priori*, $\text{TPL}(\text{PA})$ falls outside the arithmetical hierarchy due to Tarski’s [35] result on the undefinability of arithmetical truth.

After these positive results, there was hope that the nice characterizations could be extended to the realm of quantified modal logic. Thus, the focus switched to the language $\mathcal{L}_{\Box, \forall}$ of quantified modal logic without identity, which contains \perp , relation symbols, Boolean connectives, $\forall x$, and \Box , as well as the usual abbreviations such as \Diamond and $\exists x$. We define arithmetical realizations \cdot^* for $\mathcal{L}_{\Box, \forall}$ formulas as before with the only difference that we now map n -ary relation symbols to arithmetical formulas with n free variables and set $(\forall x \varphi)^{*\text{PA}} := \forall y \varphi^{*\text{PA}}$, where y is the arithmetical variable corresponding to x . The *quantified provability logic* of PA can now be defined by analogy (cf. [11]):²

$$\text{QPL}(\text{PA}) := \{A \in \mathcal{L}_{\Box, \forall} \mid \text{for any } \cdot^*, \text{ we have } \text{PA} \vdash A^{*\text{PA}}\}$$

¹Also known as c.e., recursively enumerable, or computably axiomatizable.

²In general, $\text{QPL}(T)$ may change depending on the chosen axiomatization τ for T (see [3, 28, 29]), but here we omit the axiomatization for simplicity.

as well as

$$\text{TQPL}(\text{PA}) := \{A \in \mathcal{L}_{\Box, \forall} \mid \text{for any } \cdot^*, \text{ we have } \mathbb{N} \models A^{*\text{PA}}\}.$$

It is not hard to see that $\Box \forall x A \rightarrow \forall x \Box A$ is in $\text{QPL}(\text{PA})$, and for some time it was believed that one could obtain $\text{QPL}(\text{PA})$ by adding such principles together with predicate logical reasoning to GL. A first wrinkle in this hope appeared when Montagna [32] published a new always provable principle falling outside this easily generated set. Soon after, Artemov [2] proved that $\text{TQPL}(\text{PA})$ is not even arithmetically definable. Last hopes were scattered when Vardanyan [36] and McGee [31] independently proved that $\text{QPL}(\text{PA})$ is as complex as it can possibly be: Π_2^0 complete. Moreover, Boolos and McGee [12] showed that $\text{TQPL}(\text{PA})$ is also as complex as possible: Π_1^0 complete in the set of true arithmetic.

These negative results are quite robust in various ways. For example, Vardanyan [37] showed that restricting the language to the fragment with a single unary predicate and without any modality nesting doesn't break the Π_2^0 completeness.

Restricting the complexity of the realizations to, for example, Σ_1^0 formulas doesn't help either: the corresponding set is still Π_2^0 complete, as shown by Berarducci [10].

One can consider $\text{QPL}(T)$ for an arbitrary c.e. theory T . It is easy to see that the degenerate case of $\text{QPL}(\text{PA} + \Box_{\text{PA}} \perp)$ is axiomatized by predicate logic together with $\Box \perp$. However, Visser and de Jonge [38] proved that for most theories T , $\text{QPL}(T)$ is indeed Π_2^0 complete. The title of their paper is *No Escape from Vardanyan's Theorem*.

Be that as it may, there have already been some "escapes" from Vardanyan's Theorem. For example, the one-variable fragment of $\text{QPL}(\text{PA})$ is decidable, as shown by Artemov and Japaridze [4]. Furthermore, an arithmetically complete quantified modal logic was proposed by Yavorsky [40] (see also [24]). This logic, called QGL^b , assumes that in $\Box A$ every free variable of A is bound under the box. In other words, QGL^b is roughly QGL extended with the axiom schema $\Box A \rightarrow \Box \forall x A$.

Our proposed fragment includes countably many variables and allows for open variables under the box. In order to "escape," we restrict the language to its strictly positive fragment instead. This work already began in [1], where we described the system QRC_1 and proved its decidability. Here we improve on the modal results presented there and prove the arithmetical completeness theorem for QRC_1 , previously left as a conjecture.

There is an ongoing formalization³ of this paper in the Coq Proof Assistant [14], covering Sections 3 and 4 (and part of Section 5) as of May 2022.

1.1. Overview of the paper. We start with a brief overview of strictly positive logics in Section 2, followed by the definition of QRC_1 , our main object of study, in Section 3.

After that the paper is divided into two parts, the first dealing with purely modal results (Sections 4–6), and the second delving into arithmetic (Sections 7–10). The two parts are not completely independent, but a reader who wishes to skip to Section 7 need only be familiar with the definition of Kripke model with constant domain (Definition 4.1) and the constant domain completeness theorem for QRC_1 (Theorem 5.11).

³<https://gitlab.com/ana-borges/QRC1-Coq>.

In the first part, Section 4 starts by presenting an extended definition of Kripke model that does not depend on inclusive models, with a respective soundness proof for QRC_1 . Then in Section 5 we show that QRC_1 is complete with respect to constant domain Kripke models. Section 6 remarks that QRC_1 is the strictly positive fragment of every logic between QK4 and QGL , and also of other quantified modal logics such as $\text{QK4} + \text{BF}$, where $\text{BF} = \forall x \Box A \rightarrow \Box \forall x A$ is the well-known and arithmetically unsound Barcan Formula.

In the second part, Section 7 explains our arithmetic reading of strictly positive formulas, followed by Section 8, where we prove the arithmetical completeness of QRC_1 . Section 9 then makes use of the arithmetical completeness theorem to show that QRC_1 is a fragment of $\text{QPL}(\text{PA})$. Section 10, which can be read right after Section 7, shows the arithmetical soundness of QRC_1 with respect to Heyting Arithmetic.

We end with some proposed avenues for future work in Section 11.

§2. Strictly positive logics. A key feature of our escape is given by restricting the language. Given variables x, x_i, \dots and a signature Σ fixing the constants c, c_i, \dots and relation symbols S, S_i, \dots , the formulas of QRC_1 are built up from \top , n -ary relation symbols applied to n terms (which are either variables or constants), the binary \wedge , the unary \diamond , and the quantifier $\forall x$. The provable judgments in QRC_1 are all of the form $\varphi \vdash \psi$ with φ and ψ in the above language.

Our formulas are related to arithmetic through realizations \cdot° that map n -ary predicate symbols to c.e. axiomatizations of theories indexed by n parameters (details can be found in Section 7). Given a Σ_1^0 axiomatization τ of some theory T , we extend \cdot° to non-predicate formulas such that conjunctions are interpreted as the unions of the corresponding axiom sets ($(\varphi \wedge \psi)^{\circ\tau} := \varphi^{\circ\tau} \vee \psi^{\circ\tau}$), and universal quantification corresponds to an infinite union ($(\forall x \varphi)^{\circ\tau} := \exists y \varphi^{\circ\tau}$, where y is the arithmetical variable corresponding to the modal variable x).

This interpretation is common in the study of reflection calculi [7] and is more general than the usual, finitary one, mentioned in the previous section as \cdot^* . In the latter case, conjunctions of modal formulas are simply interpreted as conjunctions of their arithmetical counterparts. We further define and make use of this finitary notion in Section 8. While it is possible to represent finite extensions of a base theory through the finitary \cdot^* -style realizations, the \cdot° approach allows for the possibility of infinitary extensions.

Using this restricted fragment and infinitary arithmetical interpretation, we define the *Strictly Positive Quantified Provability Logic* of a theory T as follows.

$$\text{QPL}^{\text{SP}}(T) := \{ \langle \varphi, \psi \rangle \mid \forall \cdot^\circ T \vdash \forall \theta (\Box_{\psi^{\circ\tau}} \theta \rightarrow \Box_{\varphi^{\circ\tau}} \theta) \}.$$

The main result of this paper is that, for a large class of theories T , the logic $\text{QPL}^{\text{SP}}(T)$ is decidable and given by the system QRC_1 as introduced in [1]. As such our paper was inspired by and contributes to three recent developments in the literature: strictly positive logics, reflection calculi, and polymodal logics.

The quintessential polymodal provability logic is GLP , introduced by Japaridze in 1986 [26]. The Reflection Calculus, RC , was first introduced by Dashkov [16] as the set of GLP -equivalences between strictly positive formulas. It was then axiomatized by Beklemishev [6], and it is the latter formulation that appears in most of the ensuing literature.

Even though RC has a strictly positive language, it is remarkably expressive, giving rise to an ordinal notation system [19] and being an appropriate tool for Π_1^0 ordinal analysis [9]. This is remarkable because, while GLP is PSPACE complete [33], RC has a polynomial-time decision procedure. Thus, at least in the case of GLP, restricting the language to the strictly positive fragment is very worthwhile.

QRC_1 was inspired by RC_1 , which is the unimodal fragment of RC. We hoped to emulate RC's success in these two dimensions, obtaining a useful calculus with a simpler complexity than the original (QPL(PA) in this case). We already see in this paper that the latter goal was achieved, since QRC_1 is decidable while QPL(PA) is Π_2^0 complete. However, the main expressibility tool available in RC, the iterated consistency statements (also known as worms [5]), are not very interesting when there is only one modality. Thus we plan to extend QRC_1 to QRC_Λ in the future.

When considering a strictly positive logic P , one may ask whether there is some modal logic L whose strictly positive fragment L^{SP} coincides with P (cf. [8]). In that case we would have

$$\varphi \vdash_P \psi \iff \varphi \vdash_{L^{\text{SP}}} \psi \iff L \vdash \varphi \rightarrow \psi, \tag{2}$$

where φ and ψ are strictly positive formulas.

We know that RC is the strictly positive fragment of GLP (in the sense of (2)), and in Section 6 we show that QRC_1 is the strictly positive fragment of QGL. However, there are strictly positive logics with no such counterpart, such as RC together with the persistence axiom $\langle \omega \rangle \varphi \vdash \varphi$. This system was described in [7] as $\text{RC}\omega$.

The study of strictly positive logics has also been developed in other communities, most notably in the field of description logics, as in [27, 30]. Furthermore, there is a significant body of work done about (non-strictly) positive logics (see [13, 17]).

§3. Quantified Reflection Calculus with one modality. The *Quantified Reflection Calculus with one modality*, or QRC_1 , is a sequent logic in a strictly positive predicate modal language introduced in [1].

The free variables of a formula φ are defined as usual, and denoted by $\text{fv}(\varphi)$. The expression $\varphi[x \leftarrow t]$ denotes the formula φ with all free occurrences of the variable x simultaneously replaced by the term t . We say that t is free for x in φ if no occurrence of a free variable in t becomes bound in $\varphi[x \leftarrow t]$.

The axioms and rules of QRC_1 are listed in the following definition from [1]. Here we removed the axiom $\diamond \forall x \varphi \vdash \forall x \diamond \varphi$ because it is an easy consequence of the calculus without it.

DEFINITION 3.1 (QRC_1 [1]). Let Σ be a signature and φ, ψ , and χ be any formulas in that language. The axioms and rules of QRC_1 are the following:

- (i) $\varphi \vdash \top$ and $\varphi \vdash \varphi$;
- (ii) $\varphi \wedge \psi \vdash \varphi$ and $\varphi \wedge \psi \vdash \psi$;
- (iii) if $\varphi \vdash \psi$ and $\varphi \vdash \chi$, then $\varphi \vdash \psi \wedge \chi$;
- (iv) if $\varphi \vdash \psi$ and $\psi \vdash \chi$, then $\varphi \vdash \chi$;
- (v) if $\varphi \vdash \psi$, then $\diamond \varphi \vdash \diamond \psi$;
- (vi) $\diamond \diamond \varphi \vdash \diamond \varphi$;
- (vii) if $\varphi \vdash \psi$, then $\varphi \vdash \forall x \psi$ ($x \notin \text{fv}(\varphi)$);
- (viii) if $\varphi[x \leftarrow t] \vdash \psi$, then $\forall x \varphi \vdash \psi$ (t free for x in φ);
- (ix) if $\varphi \vdash \psi$, then $\varphi[x \leftarrow t] \vdash \psi[x \leftarrow t]$ (t free for x in φ and ψ);
- (x) if $\varphi[x \leftarrow c] \vdash \psi[x \leftarrow c]$, then $\varphi \vdash \psi$ (c not in φ nor ψ).

If $\varphi \vdash \psi$, we say that ψ follows from φ in QRC_1 . When the signature is not clear from the context, we write $\varphi \vdash_{\Sigma} \psi$ instead.

We observe that our axioms do not include universal quantifier elimination. However, this and various other rules are readily available via the following easy lemma.

LEMMA 3.2. *The following are theorems (or derivable rules) of QRC_1 :*

1. $\forall x \forall y \varphi \vdash \forall y \forall x \varphi$;
2. $\forall x \varphi \vdash \varphi[x \leftarrow t]$ (t free for x in φ);
3. $\diamond \forall x \varphi \vdash \forall x \diamond \varphi$;⁴
4. $\forall x \varphi \vdash \forall y \varphi[x \leftarrow y]$ (y free for x in φ and $y \notin \text{fv}(\varphi)$);
5. if $\varphi \vdash \psi$, then $\varphi \vdash \psi[x \leftarrow t]$ (x not free in φ and t free for x in ψ);
6. if $\varphi \vdash \psi[x \leftarrow c]$, then $\varphi \vdash \forall x \psi$ (x not free in φ and c not in φ nor ψ).

The following are two useful complexity measures on the formulas of QRC_1 .

DEFINITION 3.3 ($d_{\diamond}, d_{\forall}$ [1]). Given a formula φ , its modal depth $d_{\diamond}(\varphi)$ is defined inductively as follows:

- $d_{\diamond}(\top) := d_{\diamond}(S(x_0, \dots, x_{n-1})) := 0$;
- $d_{\diamond}(\psi \wedge \chi) := \max\{d_{\diamond}(\psi), d_{\diamond}(\chi)\}$;
- $d_{\diamond}(\forall x \psi) := d_{\diamond}(\psi)$;
- $d_{\diamond}(\diamond \psi) := d_{\diamond}(\psi) + 1$.

Given a finite set of formulas Γ , its modal depth is $d_{\diamond}(\Gamma) := \max_{\varphi \in \Gamma} \{d_{\diamond}(\varphi)\}$.

The definition of quantifier depth d_{\forall} is analogous except for:

- $d_{\forall}(\forall x \psi) := d_{\forall}(\psi) + 1$ and
- $d_{\forall}(\diamond \psi) := d_{\forall}(\psi)$.

The modal depth provides a necessary condition for derivability, which in particular implies irreflexivity.

LEMMA 3.4 [1]. *Let φ and ψ be formulas in the language of QRC_1 .*

- If $\varphi \vdash \psi$, then $d_{\diamond}(\varphi) \geq d_{\diamond}(\psi)$.
- $\varphi \not\vdash \diamond \varphi$.

Finally, the signature of QRC_1 can be extended without strengthening the calculus.

LEMMA 3.5 [1]. *Let Σ be a signature and let C be a collection of constants not yet occurring in Σ . By Σ_C we denote the signature obtained by including these new constants C in Σ . Let φ, ψ be formulas in the language of Σ . Then, if $\varphi \vdash_{\Sigma_C} \psi$, so does $\varphi \vdash_{\Sigma} \psi$.*

§4. Relational semantics. QRC_1 was proven sound and complete with respect to relational semantics in [1]. Here we extend both of those results in the following ways: we relax the requirement for the adequateness of a frame, and we prove constant domain completeness: that if $\varphi \not\vdash \psi$ then there exists a counter model that, in addition to being finite and irreflexive, also has a constant domain.

⁴In [1] this was presented as an axiom of QRC_1 , but it is easily provable through the two \forall introduction rules and necessitation.

We begin by slightly changing the definition of frame and relational model presented in [1]. There, models for QRC_1 were described as a number of first-order models (the worlds) connected through a transitive relation R . We additionally required *inclusiveness*: that whenever w and u are worlds connected through R , the domain of w be included in the domain of u . We then used the inclusion (identity) function $\iota_{w,u}$ to refer to the element of the domain of u corresponding to an element in the domain of w .

Here, we no longer have the inclusiveness restriction. In fact, as we will see below, any configuration of domains is sound as long as the functions $\eta_{w,u}$ relating the domain of w with the domain of u respect the transitivity of R . This is clearly the case if the frame is inclusive and $\eta_{w,u} = \iota_{w,u}$, so the definitions presented in [1] are a particular case of the ones presented here.

DEFINITION 4.1. A *relational model* \mathcal{M} in a signature Σ is a tuple $\langle W, R, \{M_w\}_{w \in W}, \{\eta_{w,v}\}_{wRv}, \{I_w\}_{w \in W}, \{J_w\}_{w \in W} \rangle$ where:

- W is a non-empty set (the set of worlds, where individual worlds are referred to as w, u, v , etc.);
- R is a binary relation on W (the accessibility relation);
- each M_w is a finite set (the domain of the world w , whose elements are referred to as d, d_0, d_1 , etc.);
- if wRv , then $\eta_{w,v}$ is a function from M_w to M_v ;
- for each $w \in W$, the interpretation I_w assigns an element of the domain M_w to each constant $c \in \Sigma$, written c^{I_w} ; and
- for each $w \in W$, the interpretation J_w assigns a set of tuples $S^{J_w} \subseteq \wp((M_w)^n)$ to each n -ary relation symbol $S \in \Sigma$.

The $\langle W, R, \{M_w\}_{w \in W}, \{\eta_{w,v}\}_{wRv} \rangle$ part of the model is called its *frame*. We say that the frame (or model) is *finite* if W is finite, and that it is *constant domain* if all the M_w coincide and all the $\eta_{w,u}$ are the identity function.

The relevant frames and models will need to satisfy a number of requisites.

DEFINITION 4.2. A frame \mathcal{F} is *adequate* if:

- R is transitive: if wRu and uRv , then wRv ; and
- η functions respect transitivity: if wRu and uRv , then $\eta_{w,v} = \eta_{u,v} \circ \eta_{w,u}$.⁵

A model is *adequate* if it is based on an adequate frame and it is:

- concordant: if wRu , then $c^{I_u} = \eta_{w,u}(c^{I_w})$ for every constant c .

Note that in an adequate and rooted model the interpretation of the constants is fully determined by their interpretation at the root.

As in [1], we use assignments to define truth at a world in a first-order model. A w -assignment g is a function assigning a member of the domain M_w to each variable in the language. Any w -assignment can be seen as a v -assignment as long as wRv , by composing it with $\eta_{w,v}$ on the left. We write g^v to shorten $\eta_{w,v} \circ g$ when w is clear from the context.

Two w -assignments g and h are x -*alternative*, written $g \sim_x h$, if they coincide on all variables other than x . A w -assignment g is extended to terms by defining

⁵We only require extensional equality.

$g(c) := c^{J_w}$ for any constant c . Note that this meshes nicely with the concordant restriction of an adequate model: for any term t , if wRv then $g^v(t) = \eta_{w,v}(g(t))$.

We are finally ready to define satisfaction at a world. This definition is a straightforward adaptation of the one presented in [1] to our current definition of model. The only difference is in the case of $\diamond\varphi$, where we use g^v instead of g^t .

DEFINITION 4.3. Let $\mathcal{M} = \langle W, R, \{M_w\}_{w \in W}, \{\eta_{w,u}\}_{wRu}, \{I_w\}_{w \in W}, \{J_w\}_{w \in W} \rangle$ be an adequate model in some signature Σ , and let $w \in W$ be a world, g be a w -assignment, S be an n -ary relation symbol, and φ, ψ be formulas in the language of Σ .

We define $\mathcal{M}, w \Vdash^g \varphi$ (φ is true at w under g) by induction on φ as follows.

- $\mathcal{M}, w \Vdash^g \top$;
- $\mathcal{M}, w \Vdash^g S(t_0, \dots, t_{n-1})$ iff $\langle g(t_0), \dots, g(t_{n-1}) \rangle \in S^{J_w}$;
- $\mathcal{M}, w \Vdash^g \varphi \wedge \psi$ iff both $\mathcal{M}, w \Vdash^g \varphi$ and $\mathcal{M}, w \Vdash^g \psi$;
- $\mathcal{M}, w \Vdash^g \diamond\varphi$ iff there is a $v \in W$ such that wRv and $\mathcal{M}, v \Vdash^{g^v} \varphi$;
- $\mathcal{M}, w \Vdash^g \forall x \varphi$ iff for all w -assignments h such that $h \sim_x g$, we have $\mathcal{M}, w \Vdash^h \varphi$.

THEOREM 4.4 (Relational soundness). *If $\varphi \vdash \psi$, then for any adequate model \mathcal{M} , for any world $w \in W$, and for any w -assignment g ,*

$$\mathcal{M}, w \Vdash^g \varphi \implies \mathcal{M}, w \Vdash^g \psi.$$

PROOF. By induction on the proof of $\varphi \vdash \psi$, making the same arguments as in [1]. Here we highlight only the transitivity axiom, where the transitivity of the η functions comes into play, and also remark on the generalization on constants rule.

The transitivity axiom is $\diamond\varphi \vdash \diamond\varphi$, so assume that $\mathcal{M}, w \Vdash^g \diamond\varphi$. Then there is a world v such that wRv and $\mathcal{M}, v \Vdash^{\eta_{w,v} \circ g} \varphi$, and also a subsequent world u such that vRu and $\mathcal{M}, u \Vdash^{\eta_{v,u} \circ (\eta_{w,v} \circ g)} \varphi$. Since R is transitive, we know that wRu and thus that $\eta_{v,u} \circ (\eta_{w,v} \circ g)$ coincides with $\eta_{w,u} \circ g$. Then $\mathcal{M}, u \Vdash^{\eta_{w,u} \circ g} \varphi$, and consequently $\mathcal{M}, w \Vdash^g \diamond\varphi$, as desired.

The soundness of the generalization on constants rule, Rule 3.1(x), is the most involved part of the proof presented in [1] and depends on the construction of a model where the interpretation of a constant is changed. Building that model in this context is a simple matter of taking care to propagate that change to all future worlds using the η functions. ⊣

We end this section by noting that, even though the language of QRC_1 is quite restricted, even its \forall fragment requires counter-models with arbitrarily large domains. For example, the sequent

$$\forall x, y S(x, x, y) \wedge \forall x, y S(x, y, x) \wedge \forall x, y S(y, x, x) \vdash \forall x, y, z S(x, y, z)$$

is unprovable in QRC_1 , but satisfied by every world with at most two domain elements. This reasoning can be extended to any n : if S is an n -ary predicate symbol, let φ_n be the conjunction of the $n(n-1)/2$ formulas of the form $\forall x_0, \dots, x_{n-2} S(\dots, x_0, \dots, x_0, \dots)$, with x_0 appearing in every possible pair of positions and every other position filled by a unique variable. Then φ_n does not entail $\psi_n := \forall x_0, \dots, x_{n-1} S(x_0, \dots, x_{n-1})$ but any world with at most $n-1$ domain elements that satisfies φ_n must also satisfy ψ_n .

§5. Constant domain completeness. In [1] we proved relational completeness by building a term model that satisfies φ and doesn't satisfy ψ when $\varphi \not\vdash \psi$. That construction provides a finite, irreflexive, and rooted model with increasing domains. Here we show that it is possible to build a constant domain model instead, that is, a model where the domain of every world is exactly the same. This is extremely useful to prove the arithmetical completeness theorem in Section 8.

Before starting the formal proof, we briefly describe the main idea. The term models we build are such that each world is a pair of sets of closed formulas $p = \langle p^+, p^- \rangle$. The first set, p^+ , is the set of formulas that will be satisfied at that world, or the *positive part*. The second set, p^- , is the set of formulas that will not be satisfied at that world, or the *negative part*. All worlds must be well-formed with respect to some finite set of closed formulas Φ , which means that:

- p is closed: every formula in p^+ and every formula in p^- is closed;
- p is Φ -maximal: every formula of Φ is in either p^+ or p^- (and there are no formulas in p but not in Φ);
- p is consistent: if $\delta \in p^-$ then $\bigwedge p^+ \not\vdash \delta$; and
- p is fully-witnessed: if $\forall x \varphi \in p^-$ then there is a constant c such that $\varphi[x \leftarrow c] \in p^-$.

In that case we say that p is Φ -maximal consistent and fully witnessed, or Φ -MCW for short (the closeness condition is included in the concept, although in practice almost every formula in this section will be closed). If p and q are pairs, we write $p \subseteq q$ when $p^+ \subseteq q^+$ and $p^- \subseteq q^-$. Furthermore, if Γ is a set of formulas we write $p \subseteq \Gamma$ instead of $p^+ \cup p^- \subseteq \Gamma$.

We want to have Φ -MCW pairs where Φ is closed under subformulas. However, the naive subformulas of $\forall x \varphi$ might be open. In order to avoid that, we use the notion of closure with respect to a set of constants C defined in [1], where $\varphi[x \leftarrow c]$ is a valid subformula of $\forall x \varphi$ as long as $c \in C$.

DEFINITION 5.1 (\mathcal{Cl}_C [1]). Given a set of constants C , the closure of a formula φ under C , written $\mathcal{Cl}_C(\varphi)$, is defined by induction on the formula as such: $\mathcal{Cl}_C(\top) := \{\top\}$; $\mathcal{Cl}_C(\mathcal{S}(t_0, \dots, t_{n-1})) := \{\mathcal{S}(t_0, \dots, t_{n-1}), \top\}$; $\mathcal{Cl}_C(\varphi \wedge \psi) := \{\varphi \wedge \psi\} \cup \mathcal{Cl}_C(\varphi) \cup \mathcal{Cl}_C(\psi)$; $\mathcal{Cl}_C(\diamond \varphi) := \{\diamond \varphi\} \cup \mathcal{Cl}_C(\varphi)$; and

$$\mathcal{Cl}_C(\forall x \varphi) := \{\forall x \varphi\} \cup \bigcup_{c \in C} \mathcal{Cl}_C(\varphi[x \leftarrow c]).$$

The closure under C of a set of formulas Γ is the union of the closures under C of each of the formulas in Γ :

$$\mathcal{Cl}_C(\Gamma) := \bigcup_{\gamma \in \Gamma} \mathcal{Cl}_C(\gamma).$$

The closure of a pair p is defined as the closure of $p^+ \cup p^-$.

Going back to the overview of the completeness proof, suppose that $\varphi \not\vdash \psi$, (assuming for now that φ and ψ are closed). Defining $p := \langle \{\varphi\}, \{\psi\} \rangle$, the counter-model will be rooted on a $\mathcal{Cl}_C(p)$ -MCW extension of p , where C is a set of constants to determine. The set of constants C will be used as the domain of the root. Note that p is already closed and consistent, so taking the step to maximality is as simple as

deciding whether to add each $\chi \in \mathcal{C}l_C(p)$ to the positive or negative part of the root without ruining its consistency. The hard part is doing so in a way that guarantees that the resulting pair is fully witnessed.

The completeness proof shown in [1] uses the observation that if $\bigwedge p^+ \not\vdash \forall x \chi$, then also $\bigwedge p^+ \not\vdash \chi[x \leftarrow c]$, as long as c does not appear in either p^+ or χ (Lemma 3.2.6). This suggests a way of sorting the formulas of $\mathcal{C}l_C(p)$ into positive and negative: mark a formula as positive if and only if it is a consequence of p^+ . This guarantees that there are witnesses for the negative universal formulas as long as there are enough constants to go around. The way to make sure that there are enough constants is precisely the difference between the proof presented in [1] and the proof presented here. To that end, we introduce the following definition.

DEFINITION 5.2 (d_{const}). The number of different constants in a formula φ is represented by $d_{\text{const}}(\varphi)$. The maximum number of different constants per formula in a set of formulas Γ is defined as $d_{\text{const}}(\Gamma) := \max_{\varphi \in \Gamma} \{d_{\text{const}}(\varphi)\}$.

Note that $d_{\text{const}}(\Gamma)$ is *not* the number of different constants appearing in Γ , but the maximum number of different constants in any single formula of Γ . For example, $d_{\text{const}}(\{S_0(c_0, c_1), S_1(c_2)\}) = 2$.

We observe that the maximum number of distinct constants per formula in the closure under C of a set of formulas can be bounded by a number that *does not depend on C*.

FACT 5.3. For any formula φ , set of formulas Φ , and set of constants C :

- $d_{\text{const}}(\varphi) \leq d_{\text{const}}(\varphi[x \leftarrow c]) \leq d_{\text{const}}(\varphi) + 1$;
- $d_{\text{const}}(\varphi) \leq d_{\text{const}}(\mathcal{C}l_C(\varphi)) \leq d_{\text{const}}(\varphi) + d_{\forall}(\varphi)$;
- $d_{\text{const}}(\Phi) \leq d_{\text{const}}(\mathcal{C}l_C(\Phi)) \leq d_{\text{const}}(\Phi) + d_{\forall}(\Phi)$.

We are now ready to prove a Lindenbaum-like lemma.

LEMMA 5.4. Given a finite signature Σ with constants C and a finite set of closed formulas Φ in the language of Σ such that $|C| > 2d_{\text{const}}(\Phi) + 2d_{\forall}(\Phi)$, if $p \subseteq \mathcal{C}l_C(\Phi)$ is a closed consistent pair and p^+ is a singleton, then there is a pair $q \supseteq p$ in the language of Σ such that q is $\mathcal{C}l_C(\Phi)$ -MCW, and $d_{\diamond}(q^+) = d_{\diamond}(p^+)$.

PROOF. Like in [1], we start by defining a pair $q \supseteq p$ such that for each $\chi \in \mathcal{C}l_C(\Phi)$, $\chi \in q^+$ if and only if $p^+ \vdash \chi$ (otherwise $\chi \in q^-$). It is easy to see that this pair q is $\mathcal{C}l_C(\Phi)$ -maximal consistent, and we have $d_{\diamond}(q^+) = d_{\diamond}(p^+)$ by Lemma 3.4.

It remains to show that q is fully witnessed. Let $\forall x \psi$ be a formula in q^- . We claim that there is $d \in C$ such that d does not appear either in p^+ or in $\forall x \psi$. For this it is enough to see that $|C| > d_{\text{const}}(\bigwedge p^+) + d_{\text{const}}(\forall x \psi)$, which is the same as $|C| > d_{\text{const}}(p^+) + d_{\text{const}}(\forall x \psi)$ because p^+ is a singleton by assumption. Since $p^+ \subseteq \mathcal{C}l_C(\Phi)$, we know that $d_{\text{const}}(p^+) \leq d_{\text{const}}(\mathcal{C}l_C(\Phi))$, and similarly for $\forall x \psi$. Then by Fact 5.3 we may conclude that $d_{\text{const}}(p^+) + d_{\text{const}}(\forall x \psi) \leq 2d_{\text{const}}(\Phi) + 2d_{\forall}(\Phi)$, which suffices by our assumption on the size of C .

Since $d \in C$ does not appear in either p^+ or $\forall x \psi$, we conclude that $p^+ \not\vdash \psi[x \leftarrow d]$ by Lemma 3.2.6, and consequently $\psi[x \leftarrow d] \in q^-$ as desired. \dashv

We now recall the definition of \hat{R} from [1], which is the relation we use to connect the worlds of the term model.

DEFINITION 5.5 ($p\hat{R}q$ [1]). The relation \hat{R} between pairs is such that $p\hat{R}q$ if and only if both of following hold:

- for any formula $\diamond\varphi \in p^-$ we have $\varphi, \diamond\varphi \in q^-$; and
- there is some formula $\diamond\psi \in p^+ \cap q^-$.

LEMMA 5.6 [1]. *The relation \hat{R} restricted to consistent pairs is transitive and irreflexive.*

We now see that if w is a $\mathcal{C}l_C(\Phi)$ -MCW pair with $\diamond\varphi \in w^+$, we can find a $\mathcal{C}l_C(\Phi)$ -MCW pair v with $\varphi \in v^+$ and $w\hat{R}v$. The proof is the same as in [1], except that we now use Lemma 5.4 to obtain constant domains throughout the model.

LEMMA 5.7 (Pair existence). *Let Σ be a signature with a finite set of constants C , and Φ be a finite set of closed formulas in the language of Σ such that $|C| > 2d_{\text{const}}(\Phi) + 2d_{\forall}(\Phi)$. If p is a $\mathcal{C}l_C(\Phi)$ -MCW pair and $\diamond\varphi \in p^+$, then there is a $\mathcal{C}l_C(\Phi)$ -MCW pair q such that $p\hat{R}q$, $\varphi \in q^+$, and $d_{\diamond}(q^+) < d_{\diamond}(p^+)$.*

PROOF. Consider the pair r defined as $r^+ := \{\varphi\}$ and $r^- := \{\delta, \diamond\delta \mid \diamond\delta \in p^-\} \cup \{\diamond\varphi\}$. It is easy to check that r is consistent and that $p\hat{R}r$ (details can be found in [1]), and clearly $r \subseteq \mathcal{C}l_C(\Phi)$. We then use Lemma 5.4 to obtain a $\mathcal{C}l_C(\Phi)$ -MCW pair $q \supseteq r$ such that $d_{\diamond}(q^+) = d_{\diamond}(r^+) = d_{\diamond}(\varphi) < d_{\diamond}(p^+)$. We obtain $p\hat{R}q$ as a straightforward consequence of $p\hat{R}r$. -1

We can now define an adequate and constant domain model $\mathcal{M}[p]$ from any given finite and consistent pair p such that $\mathcal{M}[p]$ satisfies the formulas in p^+ and doesn't satisfy the formulas in p^- . The idea is exactly the same as in [1]: build a term model where each world w is a $\mathcal{C}l_M(p)$ -MCW pair, and the worlds are related by (a sub-relation of) \hat{R} .

DEFINITION 5.8. Let Σ be a signature. Given a finite consistent pair p of closed formulas in Σ such that p^+ is a singleton,⁶ we define an adequate model $\mathcal{M}[p]$. Here we will use $\Phi := p^+ \cup p^-$.

Let C be a set of at least $2d_{\text{const}}(\Phi) + 2d_{\forall}(\Phi) + 1$ different constants, including all of the ones appearing in Σ and adding more if necessary. The pairs of formulas we work with are in the signature Σ extended by C .

We start by defining the underlying frame in an iterative manner. The root is given by Lemma 5.4 applied to C and p , obtaining the $\mathcal{C}l_C(\Phi)$ -MCW pair q . Frame \mathcal{F}^0 is then defined such that its set of worlds is $W^0 := \{q\}$, its relation R^0 is empty, and the domain of q is $M_q^0 := C$.

Assume now that we already have a frame \mathcal{F}^i , and we set out to define \mathcal{F}^{i+1} as an extension of \mathcal{F}^i . For each leaf w of \mathcal{F}^i , i.e., each world such that there is no world $v \in \mathcal{F}^i$ with $wR^i v$, and for each formula $\diamond\varphi \in w^+$, use Lemma 5.7 to obtain a $\mathcal{C}l_C(\Phi)$ -MCW pair v such that $w\hat{R}v$, $\varphi \in v^+$, and $d_{\diamond}(v^+) < d_{\diamond}(w^+)$. Now add v to W^{i+1} , add $\langle w, v \rangle$ to R^{i+1} , define M_v^{i+1} as C , and define $\eta_{w,v}$ as the identity function.

The process described above terminates because each pair is finite and the modal depth of $\mathcal{C}l_C(\Phi)$ (and consequently of w^+ , for any $w \subseteq \mathcal{C}l_C(\Phi)$) is also finite. Thus

⁶This is without loss of generality, as otherwise we could take the conjunction of every formula in p^+ as the new p^+ .

there is a final frame \mathcal{F}^m , for some natural number m . This frame is constant domain by construction, but not transitive. We obtain $\mathcal{F}[p]$ as the transitive closure of \mathcal{F}^m , which is clearly still constant domain. The η functions are all the identity in C , thus satisfying the transitivity condition. We conclude that the frame $\mathcal{F}[p]$ is adequate.

In order to obtain the model $\mathcal{M}[p]$ based on the frame $\mathcal{F}[p]$, let I_q take constants in Σ to their corresponding version as domain elements. If w is any other world, let I_w coincide with I_q . This is necessary to make sure that the model is concordant, because q sees every other world, and is sufficient to see that $\mathcal{M}[p]$ is adequate. Finally, given an n -ary predicate letter S and a world w , define S^{J_w} as the set of n -tuples $\langle d_0, \dots, d_{n-1} \rangle \subseteq (M_w)^n$ such that $S(d_0, \dots, d_{n-1}) \in w^+$.

Since everything up until now was meant for closed formulas, and furthermore we are potentially adding new constants to the signature of the formulas we care about, we provide a way of replacing the free variables of a formula with constants.

DEFINITION 5.9 (φ^g [1]). Given a formula φ in a signature Σ and a function g from the set of variables to a set of constants in some signature $\Sigma' \supseteq \Sigma$, we define the formula φ^g in the signature Σ' as φ with each free variable x simultaneously replaced by $g(x)$.

The constant domain model defined above coincides with the non-constant domain definition provided in [1] in everything other than that it refers to the stronger Lemmas 5.4 and 5.7 that keep the domain constant. Thus the Truth Lemma holds with exactly the same proof.

LEMMA 5.10 (Truth Lemma [1]). *Let Σ be a signature. For any finite non-empty consistent pair p of closed formulas in the language of Σ , world $w \in \mathcal{M}[p]$, w -assignment g , and formula φ in the language of Σ such that $\varphi^g \in \mathcal{C}\ell_{M_w}(p)$, we have that*

$$\mathcal{M}[p], w \Vdash^g \varphi \iff \varphi^g \in w^+.$$

We are now ready to prove the constant domain completeness theorem.

THEOREM 5.11 (Constant domain completeness). *Let Σ be a signature and φ, ψ be formulas in Σ . If $\varphi \not\vdash \psi$, then there is an adequate, finite, irreflexive, and constant domain model \mathcal{M} , a world $w \in W$, and a w -assignment g such that*

$$\mathcal{M}, w \Vdash^g \varphi \quad \text{and} \quad \mathcal{M}, w \not\vdash^g \psi.$$

PROOF. As in the original proof of relational completeness presented in [1], define a new constant c_x for each free variable x of φ and ψ and let Σ' be the signature Σ augmented with these new constants and a dummy constant c_0 . Let g be the assignment taking each free variable x of φ and ψ to c_x and every other variable to c_0 . Note that $\varphi^g \not\vdash_{\Sigma'} \psi^g$ by Rule 3.1(x) and Lemma 3.5. Build $\mathcal{M} := \mathcal{M}[\{\{\varphi^g\}, \{\psi^g\}\}]$ as described in Definition 5.8, with root w . Then by Lemma 5.10 we have both $\mathcal{M}, w \Vdash^g \varphi$ and $\mathcal{M}, w \not\vdash^g \psi$, as desired. ⊣

§6. The strictly positive fragment of QK4 and QGL. Consider the language of full quantified modal logic with no equality nor constant nor function symbols, $\mathcal{L}_{\square, \forall}$. We use upper case letters to refer to formulas in this language. A propositional

modal logic S can be extended to a quantified modal logic QS as described in [25] by extending the language and adding the following axiom schema and rule:

- [∀E] $\forall x A \rightarrow A[x \leftarrow y]$ (y free for x in A);
- [∀I] if $QS \vdash A \rightarrow B$, then $QS \vdash A \rightarrow \forall x B$ ($x \notin \text{fv}(A)$).

The logic K4 is obtained from the smallest normal modal logic K by adding the 4 Axiom $\Box A \rightarrow \Box \Box A$. The logic GL is K4 extended with Löb's Axiom $\Box(\Box A \rightarrow A) \rightarrow \Box A$. Dashkov [16] showed that RC_1 is the strictly positive fragment of both K4 and GL (and of any logic between them), in the sense that, if φ and ψ are built up from \top , propositional symbols, conjunctions, and diamonds, then $\varphi \vdash_{RC_1} \psi$ if and only if $K4 \vdash \varphi \rightarrow \psi$, and similarly for GL. In this section we show that this equivalence is maintained in the predicate case.

Since QRC_1 includes constants in its language and predicate modal logics are often presented without them, we define a map from the language of QRC_1 to its constant-free subset, replacing constants with fresh variables. Thus, given a finite set of QRC_1 formulas Φ where the constants appearing in Φ are $c = c_0, \dots, c_{n-1}$, let $\mathbf{x} = x_0, \dots, x_{n-1}$ be fresh variables with respect to Φ . Then we define $\varphi^\Phi := \varphi[c_0 \leftarrow x_0] \cdots [c_{n-1} \leftarrow x_{n-1}]$ for each $\varphi \in \Phi$, and abbreviate this long substitution with the notation $\varphi[c \leftarrow \mathbf{x}]$. Thus we also have $\varphi = \varphi^\Phi[\mathbf{x} \leftarrow c]$.

It is straightforward to iterate Rules 3.1(x) and 3.1(ix) in order to confirm that the above translation is harmless when it comes to provability.

FACT 6.1. *Let Φ be a finite set of QRC_1 formulas such that $\varphi, \psi \in \Phi$. Then*

$$\varphi \vdash \psi \iff \varphi^\Phi \vdash \psi^\Phi.$$

The semantics presented here for QRC_1 is a generalization of more traditional semantics for quantified modal logics, since it uses maps between domains. However, for this section, we will only need constant domain models. These can be easily interpreted in the usual framework as described in [25]. The following is a well-known (and easily provable) result.

THEOREM 6.2 (Soundness for QS [25]). *If $\mathcal{F} = \langle W, R \rangle$ satisfies every theorem of S, then any constant domain model based on \mathcal{F} satisfies every theorem of QS.*

Thus any transitive constant domain model is a QK4 model, and any transitive and conversely well-founded constant domain model is a QGL model.⁷

We are now ready to show the main theorem of this section.

THEOREM 6.3. *Let φ and ψ be QRC_1 formulas and let QS be any logic between QK4 and QGL. Then $\varphi \vdash \psi$ if and only if there is a finite set Φ such that $\varphi, \psi \in \Phi$ and $QS \vdash \varphi^\Phi \rightarrow \psi^\Phi$.*

PROOF. For the left-to-right implication, let Φ be the set of formulas appearing in the proof of $\varphi \vdash \psi$. It is easy to see by induction on the length of the QRC_1 proof that $QK4 \vdash \varphi^\Phi \rightarrow \psi^\Phi$, which then implies that $QS \vdash \varphi^\Phi \rightarrow \psi^\Phi$.

For the right-to-left implication, we prove the contrapositive. If $\varphi \not\vdash \psi$, then by Fact 6.1 we know that $\varphi^\Phi \not\vdash \psi^\Phi$ for any Φ including both φ and ψ . Let \mathcal{M} be a QRC_1 model satisfying φ^Φ and not satisfying ψ^Φ , as given by Theorem 5.11. This

⁷We assume constant domain models for simplicity, but these results also hold in the inclusive case.

model is transitive, irreflexive, and has finitely-many worlds, which means it is a QGL model by Theorem 6.2. We conclude that $QGL \not\vdash \varphi^\Phi \rightarrow \psi^\Phi$, and consequently that $QS \not\vdash \varphi^\Phi \rightarrow \psi^\Phi$. ⊣

We end with the following observation. The proof of Theorem 6.3 holds for any quantified modal logic that extends QK4 and is sound for any combination of constant-domain, finite, transitive, and irreflexive models. The Barcan Formula (BF), $\forall x \Box A \rightarrow \Box \forall x A$, is popular in the world of quantified modal logic and is always sound in constant-domain models. Thus the proof of Theorem 6.3 also serves to see that, for example, $\varphi \vdash \psi$ if and only if $QK4 + BF \vdash \varphi^\Phi \rightarrow \psi^\Phi$. This is curious because BF does not hold in QGL and is in fact unsound with the usual provability interpretation of \Box . From this we conclude that it would be worthwhile to study QRC_1 from points of view unrelated to provability.

§7. Arithmetical semantics. Recall that the language of arithmetic is that of first-order logic together with the symbols $\{0, 1, +, \times, \leq, =\}$ with their usual arities (see [23] for details). We will rely on Σ_1^0 collection throughout the sections on arithmetic, so we take $\mathbf{I}\Sigma_1$ as our base theory.

Let T be an elementary presented extension of $\mathbf{I}\Sigma_1$, meaning that there is a bounded formula $Ax_T(u)$ that is true in the standard model if and only if u is (the Gödel number of) an axiom of T . In this setting it is traditional to define Gödel’s provability predicate $\Box_T \varphi$ as $\exists p \text{Prf}_T(p, \ulcorner \varphi \urcorner)$, where

$$\begin{aligned} \text{Prf}_T(p, n) &:= \text{sequence}(p) \wedge p_{|p|-1} = n \wedge \\ &\quad \forall k < |p| (Ax_T(p_k) \vee \exists i, j < k \text{MP}(p_i, p_j, p_k) \vee \exists i < k \text{Gen}(p_i, p_k)). \end{aligned}$$

Roughly, $\Box_T \varphi$ formalizes that there is a Hilbert-style proof of φ , that is, a finite sequence p_0, \dots, p_m such that p_m is φ and that each p_k is either (the Gödel number of) an axiom of T or follows from previous elements of the sequence through either *modus ponens* or generalization.

Note that all the functions used in the definition of Prf_T can be naturally defined as bounded formulas in $\mathbf{I}\Sigma_1$, and thus Prf_T is itself a bounded formula. This means that $\Box_T \varphi$ is Σ_1^0 . We write $\Diamond_T \varphi$ as shorthand for $\neg \Box_T \neg \varphi$.

In what follows we will be interested in c.e. theories, which are theories whose axioms can be defined by a Σ_1^0 formula. It is known by Craig’s Trick [15, 18] that any such theory has an equivalent elementary presentation, allowing us to use the regular definition of \Box_T . However, we will work with specific axiomatizations and thus it is sometimes more convenient to allow Prf to be a Σ_1^0 formula. For a given Σ_1^0 axiomatization τ of T , we define $\Box_\tau \varphi := \exists p \text{Prf}_\tau(p, \ulcorner \varphi \urcorner)$, where

$$\begin{aligned} \text{Prf}_\tau(p, n) &:= \text{sequence}(p) \wedge p_{|p|-1} = n \wedge \\ &\quad \forall k < |p| (\tau(p_k) \vee \exists i, j < k \text{MP}(p_i, p_j, p_k) \vee \exists i < k \text{Gen}(p_i, p_k)), \end{aligned}$$

the only difference from Prf_T being that we use the Σ_1^0 formula τ instead of the bounded Ax_T . Thus Prf_τ is equivalent to a Σ_1^0 formula (provably in $\mathbf{I}\Sigma_1$), and so $\Box_\tau \varphi$ is still Σ_1^0 . We define $\Diamond_\tau \varphi$ as $\neg \Box_\tau \neg \varphi$, as usual.

We wish to interpret the strictly positive formulas in the language of QRC_1 as parameterized axiomatizations of arithmetical theories extending IS_1 . Let τ be a bounded axiomatization of a sound c.e. base theory T extending IS_1 .

A realization \cdot° interprets each n -ary relation symbol $S(\mathbf{x}, \mathbf{c})$ as an $(n + 1)$ -ary Σ_1^0 formula $\sigma(u, \mathbf{y}, \mathbf{z})$ in the language of arithmetic, where \mathbf{y} matches with \mathbf{x} and \mathbf{z} matches with \mathbf{c} .⁸ We then extend this notion to any formula as follows (we add $\tau(u)$ to the axioms of the interpretation of any relation symbol to guarantee that every theory is an extension of T).

DEFINITION 7.1 ($\cdot^{\circ\tau}$ [1]). Let \cdot° be a realization such that, for a given n -ary predicate S and terms \mathbf{t} , $S(\mathbf{t})^\circ$ is an $(n + 1)$ -ary Σ_1^0 arithmetical formula where modal variables x_k are interpreted as y_k and modal constants c_k are interpreted as z_k . We extend this realization to QRC_1 formulas as follows:

- $\top^{\circ\tau} := \tau(u)$;
- $S(\mathbf{x}, \mathbf{c})^{\circ\tau} := S(\mathbf{x}, \mathbf{c})^\circ \vee \tau(u)$;
- $(\psi \wedge \delta)^{\circ\tau} := \psi^{\circ\tau} \wedge \delta^{\circ\tau}$;
- $(\diamond\psi)^{\circ\tau} := \tau(u) \vee (u = \ulcorner \diamond_{\psi^{\circ\tau}} \top \urcorner)$;
- $(\forall x_i \psi)^{\circ\tau} := \exists y_i \psi^{\circ\tau}$.

We remark that, if the free variables of φ are x_0, \dots, x_{n-1} and the constants appearing in φ are c_0, \dots, c_{m-1} , then the free variables of $\varphi^{\circ\tau}$ are $u, y_0, \dots, y_{n-1}, z_0, \dots, z_{m-1}$. For example, if $S(x_0, c_1)^{\circ\tau} = \sigma(u, y_0, z_1) \vee \tau(u)$, then

$$(\diamond S(x_0, c_1))^{\circ\tau} = \tau(u) \vee (u = \ulcorner \diamond_{\sigma(u, y_0, z_1) \vee \tau(u)} \top \urcorner).$$

The dotted variables y_0 and z_1 in the expression $u = \ulcorner \diamond_{\sigma(u, y_0, z_1) \vee \tau(u)} \top \urcorner$ indicate that y_0 and z_1 are free variables of this expression that, upon being instantiated by natural numbers n and m , shall be replaced by the numerals \bar{n} and \bar{m} instead.

Since we have Σ_1^0 collection, $\varphi^{\circ\tau}$ is always provably equivalent to a Σ_1^0 formula, and it represents the axiomatization of an extension of T .

We briefly inspect our intuitions regarding this arithmetic interpretation and the interaction between \forall and \diamond . As mentioned in Lemma 3.2, the sequent $\diamond\forall x \varphi \vdash \forall x \diamond\varphi$ is provable in QRC_1 . In the arithmetical reading it says that if the union of many theories (the $\forall x \varphi$ part) is consistent, then so is each individual theory. The converse, on the other hand, must not be a consequence of QRC_1 , for it is not in general the case that the union of many consistent theories is itself consistent.

We can now define $\text{QRC}_1(T)$:

$$\text{QRC}_1(T) := \{\varphi(\mathbf{x}, \mathbf{c}) \vdash \psi(\mathbf{x}, \mathbf{c}) \mid \forall \cdot^\circ T \vdash \forall \theta \forall \mathbf{y}, \mathbf{z} (\Box_{\psi^{\circ\tau}} \theta \rightarrow \Box_{\varphi^{\circ\tau}} \theta)\},$$

where θ is a closed formula and $\varphi^{\circ\tau}, \psi^{\circ\tau}$ in general depend on \mathbf{y} and \mathbf{z} .

We will show that

$$\text{QRC}_1 = \text{QRC}_1(T)$$

for any sound c.e. theory T extending IS_1 . The left-to-right inclusion was already proved in [1] for IS_1 , and the same proof goes through for any such T . The most

⁸In the sections on arithmetic, we always use x for variables of QRC_1 and y, z, u for variables of arithmetic. Furthermore, u is reserved for the Gödel numbers of axioms of theories.

notable part of the proof is the case of Rule 3.1(vii) (\forall introduction on the right), which uses Σ_1^0 collection. We go into detail on the very similar proof of Theorem 10.3.

THEOREM 7.2 (Arithmetical soundness [1]). $\text{QRC}_1 \subseteq \text{QRC}_1(T)$.

We dedicate the next section to the proof of the other inclusion.

§8. Arithmetical completeness. The proof of arithmetical completeness closely follows the proof of Solovay’s Theorem found in [11]. The arithmetical semantics used there is different from the one presented in the previous section, allowing only for finite extensions of the base theory instead of arbitrary c.e. ones. This is not a problem because finite axiomatizations are enough to show completeness. We use \cdot° in the general setting and \cdot^* in the finite one. The base theory T considered here is any sound c.e. extension of IS_1 . Note that we use the bounded Prf_T formula in the definition of \cdot^{*T} (depending on an elementary axiomatization τ of T), as otherwise Solovay’s proof wouldn’t go through.

DEFINITION 8.1 (\cdot^{*T}). Let \cdot^* be a realization such that, for a given predicate S and terms \mathbf{t} , $S(\mathbf{t})^*$ is an arithmetical formula with the same arity as S where modal variables x_k are interpreted as y_k and modal constants c_k are interpreted as z_k . We extend this realization to QRC_1 formulas as follows:

- $\top^{*T} := \top$;
- $S(\mathbf{t})^{*T} := S(\mathbf{t})^*$;
- $(\varphi \wedge \psi)^{*T} := \varphi^{*T} \wedge \psi^{*T}$;
- $(\diamond\varphi)^{*T} := \diamond_T \varphi^{*T}$;
- $(\forall x_k \varphi)^{*T} := \forall y_k \varphi^{*T}$.

The idea of Solovay’s proof is to embed a Kripke model not satisfying the desired unprovable formula in the language of arithmetic. If the embedding is done correctly, it is possible to prove that a formula is satisfied at a world of the Kripke model exactly when it is a consequence of the representation of that world in the desired theory T .

Given two strictly positive formulas φ and ψ such that $\varphi \not\vdash \psi$ in QRC_1 , let $\mathcal{M}_{\varphi,\psi}$ be a finite, irreflexive, and constant domain adequate model satisfying φ and not satisfying ψ at the root 1 under a 1-assignment $g_{\varphi,\psi}$. This model and assignment exist by Theorem 5.11. Since $\mathcal{M}_{\varphi,\psi}$ has constant domain, we refer to $g_{\varphi,\psi}$ and any other i -assignments as just assignments, omitting the relevant world.

We assume that the worlds of $\mathcal{M}_{\varphi,\psi}$ are $W = \{1, 2, \dots, N\}$, where 1 is the root. We define a new (adequate) model \mathcal{M} that is a copy of $\mathcal{M}_{\varphi,\psi}$, except that it has an extra world 0 as the new root. This world 0 is connected to all the other worlds through R and has the same domain, constant interpretation, and relational symbol interpretation as 1. The functions $\eta_{0,i}$ with $0 < i \leq N$ are all defined as the identity function.

Let λ_i be the Solovay sentences for T as defined in [11], satisfying the following Embedding Lemma.

LEMMA 8.2 (Embedding [11]).

1. $\mathbb{N} \models \lambda_0$;
2. $T \vdash \bigvee_{i \leq N} \lambda_i$;
3. $T \vdash \lambda_i \rightarrow \bigwedge_{j \leq N, j \neq i} \neg \lambda_j$, for $i \leq N$;

4. $T \vdash \lambda_i \rightarrow \bigwedge_{j \leq N, iRj} \diamond_T \lambda_j$, for $i \leq N$;
5. $T \vdash \lambda_i \rightarrow \square_T \bigvee_{j \leq N, iRj} \lambda_j$, for $0 < i \leq N$.

The domain of every world is M . Let m be the size of M , and $[\cdot]$ be a bijection between M and the set of numerals $\{\overline{0}, \dots, \overline{m-1}\}$.

We now define for a given n -ary predicate symbol S and terms $\mathbf{t} = t_0, \dots, t_{n-1}$ the \cdot^{\otimes} interpretation as follows:

$$S(\mathbf{t})^{\otimes} := \bigvee_{i \leq N} \left(\lambda_i \wedge \Phi_i^{S(\mathbf{t})} \right), \text{ where}$$

$$\Phi_i^{S(\mathbf{t})} := \bigvee_{\langle a_0, \dots, a_{n-1} \rangle \in S^{J_i}} \bigwedge_{l < n} \left([a_l] = \begin{cases} y_k \bmod m & \text{if } t_l = x_k \\ z_k \bmod m & \text{if } t_l = c_k \end{cases} \right).$$

Here x_k is any QRC_1 variable, y_k is the corresponding T variable, c_k is any QRC_1 constant, z_k is the corresponding T variable, and J_i is the denotation of a predicate symbol at world i . Note that $x_k \in \text{fv}(S(\mathbf{t}))$ if and only if $y_k \in \text{fv}(S(\mathbf{t})^{\otimes})$ and c_k appears in $S(\mathbf{t})$ if and only if $z_k \in \text{fv}(S(\mathbf{t})^{\otimes})$. The \cdot^{\otimes} interpretation is extended to non-predicate formulas as described in Definition 8.1.

FACT 8.3. *For any QRC_1 formula φ , variable x_k , and constant c_k , we have that $x_k \in \text{fv}(\varphi)$ if and only if $y_k \in \text{fv}(\varphi^{\otimes T})$ and that c_k appears in φ if and only if $z_k \in \text{fv}(\varphi^{\otimes T})$.*

We further note that $\varphi^{\otimes T}$ is invariant under replacing variables by themselves modulo m , provably in T .

LEMMA 8.4. *For any QRC_1 formula φ and any arithmetical variable y :*

1. $T \vdash \varphi^{\otimes T} \leftrightarrow \varphi^{\otimes T}[y \leftarrow y \bmod m]$;
2. $T \vdash \forall y \varphi^{\otimes T} \leftrightarrow \forall y < m \varphi^{\otimes T}$.

PROOF. The second item is a straightforward consequence of the first, which is proved by an external induction on the complexity of φ , noting that T proves $(y \bmod m) \bmod m = y \bmod m$ for any y . ⊢

We now prove two versions of a Truth Lemma, one for when φ is forced at a world i of \mathcal{M} , and one for when it isn't. We wish to show that T proves $\varphi^{\otimes T}$ (respectively $\neg \varphi^{\otimes T}$) as a consequence of λ_i , as long as the free variables of φ are interpreted in the same way in both settings.

In order to use concise notation, we shall say \mathbf{y} instead of y_0, \dots, y_{n-1} , and similarly for other terms. We also abbreviate iterated substitutions, writing $\varphi^{\otimes T}[\mathbf{y} \leftarrow [g(\mathbf{x})]]$ instead of $\varphi^{\otimes T}[y_0 \leftarrow [g(x_0)]] \cdots [y_{n-1} \leftarrow [g(x_{n-1})]]$ and writing $\varphi^{\otimes T}[\mathbf{z} \leftarrow [c^{I_0}]]$ instead of $\varphi^{\otimes T}[z_0 \leftarrow [c_0^{I_0}]] \cdots [z_{n'-1} \leftarrow [c_{n'-1}^{I_0}]]$.

LEMMA 8.5 (Truth Lemma: positive). *Let φ be a QRC_1 formula with free variables $\mathbf{x} = x_0, \dots, x_{n-1}$ and constants $\mathbf{c} = c_0, \dots, c_{n'-1}$. Then for any world $i \leq N$ and any assignment g ,*

$$\mathcal{M}, i \Vdash^g \varphi \implies T \vdash \lambda_i \rightarrow \varphi^{\otimes T}[\mathbf{y} \leftarrow [g(\mathbf{x})]][\mathbf{z} \leftarrow [c^{I_0}]].$$

PROOF. By external induction on the complexity of φ . The cases of \top and \wedge are straightforward.

In the case of relational symbols, we assume without loss of generality that the relevant formula is $S(x_0, c_0)$. If $\mathcal{M}, i \Vdash^g S(x_0, c_0)$, then $\langle g(x_0), (c_0)^{I_i} \rangle \in S^{J_i}$. Reason in T and assume λ_i . It suffices to prove

$$\Phi_i^{S(x_0, c_0)} [y_0 \leftarrow \ulcorner g(x_0) \urcorner] [z_0 \leftarrow \ulcorner (c_0)^{I_0} \urcorner],$$

which implies $S(x_0, c_0)^{\otimes} [y_0 \leftarrow \ulcorner g(x_0) \urcorner] [z_0 \leftarrow \ulcorner (c_0)^{I_0} \urcorner]$ under the assumption of λ_i . We need to find $\langle a_0, a_1 \rangle \in S^{J_i}$ such that $\ulcorner a_0 \urcorner = \ulcorner g(x_0) \urcorner \pmod m$ and $\ulcorner a_1 \urcorner = \ulcorner (c_0)^{I_0} \urcorner \pmod m$. Noting that $\ulcorner b \urcorner \pmod m$ is provably equal to $\ulcorner b \urcorner$ for any b , we pick $a_0 := g(x_0)$ and $a_1 := (c_0)^{I_0}$. This concludes this step of the proof because $(c_0)^{I_0}$ is equal to $(c_0)^{I_i}$, for any world i .

For $\forall x_0 \varphi$, assume without loss of generality that the free variables of φ are x_0 and x_1 . If $\mathcal{M}, i \Vdash^g \forall x_0 \varphi$ then for every assignment $h \sim_{x_0} g$ we have $\mathcal{M}, i \Vdash^h \varphi$. We wish to show

$$T \vdash \lambda_i \rightarrow (\forall y_0 \varphi^{\otimes T}) [y_1 \leftarrow \ulcorner g(x_1) \urcorner] [z \leftarrow \ulcorner \mathbf{c}^{I_0} \urcorner].$$

Reason in T and assume λ_i . By Lemma 8.4, it is enough to show

$$(\forall y_0 < m \varphi^{\otimes T}) [y_1 \leftarrow \ulcorner g(x_1) \urcorner] [z \leftarrow \ulcorner \mathbf{c}^{I_0} \urcorner].$$

Since y_0, y_1 , and z are all different, we can push the substitutions inside and prove

$$\forall y_0 < m \varphi^{\otimes T} [y_1 \leftarrow \ulcorner g(x_1) \urcorner] [z \leftarrow \ulcorner \mathbf{c}^{I_0} \urcorner]$$

instead. Let $y_0 < m$ be arbitrary. Since $\ulcorner \cdot \urcorner$ is a bijection, there is $a \in M$ such that $\ulcorner a \urcorner = y_0$. We define an assignment h such that $h \sim_{x_0} g$ and $h(x_0) := a$. By assumption, $\mathcal{M}, i \Vdash^h \varphi$, so by the induction hypothesis we obtain

$$\varphi^{\otimes T} [y_0 \leftarrow \ulcorner h(x_0) \urcorner] [y_1 \leftarrow \ulcorner h(x_1) \urcorner] [z \leftarrow \ulcorner \mathbf{c}^{I_0} \urcorner].$$

This concludes the argument because $\ulcorner h(x_0) \urcorner = y_0$ and $h(x_1) = g(x_1)$.

Finally, consider the case of $\diamond \varphi$. If $\mathcal{M}, i \Vdash^g \diamond \varphi$, then there is $j \leq N$ such that iRj and $\mathcal{M}, j \Vdash^g \varphi$. Reason in T and assume λ_i . By Lemma 8.2.4, we obtain $\diamond_T \lambda_j$. Then the induction hypothesis under the box gives us the desired $\diamond_T (\varphi^{\otimes T} [y \leftarrow \ulcorner \mathbf{g}(\mathbf{x}) \urcorner] [z \leftarrow \ulcorner \mathbf{c}^{I_0} \urcorner])$. \dashv

LEMMA 8.6 (Truth Lemma: negative). *Let φ be a QRC_1 formula with free variables $\mathbf{x} = x_0, \dots, x_{n-1}$ and constants $\mathbf{c} = c_0, \dots, c_{n'-1}$. Then for any world $0 < i \leq N$ and any assignment g ,*

$$\mathcal{M}, i \nVdash^g \varphi \implies T \vdash \lambda_i \rightarrow \neg \varphi^{\otimes T} [y \leftarrow \ulcorner \mathbf{g}(\mathbf{x}) \urcorner] [z \leftarrow \ulcorner \mathbf{c}^{I_0} \urcorner].$$

PROOF. By external induction on the complexity of φ . The cases of \top and \wedge are straightforward.

For the relational symbols, consider $S(x_0, c_0)$ without loss of generality. If $\mathcal{M}, i \nVdash^g S(x_0, c_0)$, then $\langle g(x_0), (c_0)^{I_i} \rangle \notin S^{J_i}$. Reason in T and assume λ_i . We obtain $\neg \lambda_j$ for every $j \neq i$ by Lemma 8.2.3, and hence need only show $\neg \Phi_i^{S(x_0, c_0)} [y_0 \leftarrow \ulcorner g(x_0) \urcorner] [z_0 \leftarrow \ulcorner (c_0)^{I_0} \urcorner]$. In other words, we need to check that if $\langle a_0, a_1 \rangle \in S^{J_i}$, either $\ulcorner a_0 \urcorner \neq \ulcorner g(x_0) \urcorner \pmod m$, or $\ulcorner a_1 \urcorner \neq \ulcorner (c_0)^{I_0} \urcorner \pmod m$. This follows from our observation that $\langle g(x_0), (c_0)^{I_i} \rangle \notin S^{J_i}$, taking into account that $\ulcorner b \urcorner \pmod m$ is equal to $\ulcorner b \urcorner$ for any b , that $\ulcorner \cdot \urcorner$ is injective, and that $(c_0)^{I_0} = (c_0)^{I_i}$.

Consider now the case of $\forall x_0 \varphi$. We assume without loss of generality that $\text{fv}(\varphi) = \{x_0, x_1\}$. If $\mathcal{M}, i \Vdash^g \forall x_0 \varphi$, then there is an assignment $h \sim_{x_0} g$ such that $\mathcal{M}, i \Vdash^h \varphi$. Reason in T and assume λ_i . From the induction hypothesis we obtain $\neg \varphi^{\otimes T} [y_0 \leftarrow \ulcorner h(x_0) \urcorner] [y_1 \leftarrow \ulcorner h(x_1) \urcorner] [z \leftarrow \ulcorner c^{I_0} \urcorner]$, which then implies $\neg \forall y_0 \varphi^{\otimes T} [y_1 \leftarrow \ulcorner h(x_1) \urcorner] [z \leftarrow \ulcorner c^{I_0} \urcorner]$. This is what we wanted, taking into account that $h(x_1) = g(x_1)$.

Finally, in the case of $\diamond \varphi$, assume that $\mathcal{M}, i \not\Vdash^g \diamond \varphi$. Then for every j such that iRj , we have $\mathcal{M}, j \not\Vdash^g \varphi$ and thus for each such j the induction hypothesis gives us $T \vdash \lambda_j \rightarrow \neg \varphi^{\otimes T} [y \leftarrow \ulcorner g(x) \urcorner] [z \leftarrow \ulcorner c^{I_0} \urcorner]$, which put together implies $T \vdash \bigvee_{iRj} \lambda_j \rightarrow \neg \varphi^{\otimes T} [y \leftarrow \ulcorner g(x) \urcorner] [z \leftarrow \ulcorner c^{I_0} \urcorner]$. Reason in T and assume λ_i . By Lemma 8.2.5 and our assumption, we obtain $\Box_T \bigvee_{iRj} \lambda_j$. Taking the previous observation under the box, we conclude $\Box_T \neg \varphi^{\otimes T} [y \leftarrow \ulcorner g(x) \urcorner] [z \leftarrow \ulcorner c^{I_0} \urcorner]$, which is precisely $\neg (\diamond \varphi)^{\otimes T} [y \leftarrow \ulcorner g(x) \urcorner] [z \leftarrow \ulcorner c^{I_0} \urcorner]$. \dashv

We are ready to prove something analogous to Solovay's Theorem, which is the precursor to our desired completeness theorem.

THEOREM 8.7. *If φ, ψ are QRC₁ formulas with free variables x and constants c such that $\varphi \not\vdash \psi$, we have $T \not\vdash (\varphi^{\otimes T} \rightarrow \psi^{\otimes T}) [y \leftarrow \ulcorner g_{\varphi, \psi}(x) \urcorner] [z \leftarrow \ulcorner c^{I_0} \urcorner]$.*

PROOF. Recall that \mathcal{M} satisfies φ and not ψ at world 1 under the assignment $g := g_{\varphi, \psi}$.

Since $\mathcal{M}, 1 \Vdash^g \varphi$, we obtain $T \vdash \lambda_1 \rightarrow \varphi^{\otimes T} [y \leftarrow \ulcorner g(x) \urcorner] [z \leftarrow \ulcorner c^{I_0} \urcorner]$ from Lemma 8.5. Since $\mathcal{M}, 1 \not\Vdash^g \psi$, we obtain $T \vdash \lambda_1 \rightarrow \neg \psi^{\otimes T} [y \leftarrow \ulcorner g(x) \urcorner] [z \leftarrow \ulcorner c^{I_0} \urcorner]$ from Lemma 8.6. Thus $T \vdash \lambda_1 \rightarrow \neg (\varphi^{\otimes T} \rightarrow \psi^{\otimes T}) [y \leftarrow \ulcorner g(x) \urcorner] [z \leftarrow \ulcorner c^{I_0} \urcorner]$.

By Lemma 8.2.4 and the fact that $0R1$, we obtain $T \vdash \lambda_0 \rightarrow \diamond_T \lambda_1$, so putting this together with the previous observation under the box, $T \vdash \lambda_0 \rightarrow \diamond_T \neg (\varphi^{\otimes T} \rightarrow \psi^{\otimes T}) [y \leftarrow \ulcorner g(x) \urcorner] [z \leftarrow \ulcorner c^{I_0} \urcorner]$.

By Lemma 8.2.1, we know that $\mathbb{N} \models \lambda_0$, and thus by the soundness of T , we know that $\mathbb{N} \models \diamond_T \neg (\varphi^{\otimes T} \rightarrow \psi^{\otimes T}) [y \leftarrow \ulcorner g(x) \urcorner] [z \leftarrow \ulcorner c^{I_0} \urcorner]$. Then it must be the case that $T \not\vdash (\varphi^{\otimes T} \rightarrow \psi^{\otimes T}) [y \leftarrow \ulcorner g(x) \urcorner] [z \leftarrow \ulcorner c^{I_0} \urcorner]$. \dashv

We now define an arithmetical realization \cdot^{\odot} in the style of Section 7 that behaves like \cdot^{\otimes} . Recall that if τ is a Σ_1^0 formula axiomatizing T , the realization $\cdot^{\circ\tau}$ sends QRC₁ formulas to Σ_1^0 axiomatizations of theories extending T . So in particular we always have $(\varphi \wedge \psi)^{\circ\tau} = \varphi^{\circ\tau} \vee \psi^{\circ\tau}$, because χ is an axiom of the union of the theories axiomatized by $\varphi^{\circ\tau}$ and $\psi^{\circ\tau}$ precisely when χ is an axiom of one of them. This is different from \cdot^{*T} realizations such as $\cdot^{\otimes T}$, which send QRC₁ formulas to generic arithmetical formulas. We then interpret these formulas as finite extensions of T , so in particular $(\varphi \wedge \psi)^{*T} = \varphi^{*T} \wedge \psi^{*T}$, because the union of $T + \varphi^{*T}$ and $T + \psi^{*T}$ is the same as $T + \varphi^{*T} \wedge \psi^{*T}$.

We define \cdot^{\odot} such that $S(\mathbf{t})^{\odot} := \tau(u) \vee (u = \ulcorner S(\mathbf{t})^{\otimes T} \urcorner)$, and extend it to non-atomic formulas as described in Definition 7.1. Note that $x_k \in \text{fv}(\varphi)$ if and only if $y_k \in \text{fv}(\varphi^{\circ\tau})$ and c_k appears in φ if and only if $z_k \in \text{fv}(\varphi^{\circ\tau})$.

LEMMA 8.8. *For any QRC₁ formula φ with free variables x and constants c ,*

$$T \vdash \forall \theta \forall \mathbf{y}, \mathbf{z} (\Box_{\varphi^{\circ\tau}} \theta \leftrightarrow \Box_T (\varphi^{\otimes T} \rightarrow \theta)),$$

where θ is (the Gödel number of) a closed formula.

PROOF. By external induction on φ . There is nothing to show for \top , because since τ is a bounded formula, \Box_τ is the same as \Box_T .

The case of relational symbols is a straightforward consequence of the formalized Deduction Theorem [18].

In the case of \wedge , we take $\varphi = \psi \wedge \delta$, and we omit the variables \mathbf{y} and \mathbf{z} , as they introduce visual clutter but don't make the proof any more complex.

(\rightarrow) Reason in T and fix an arbitrary θ , assuming $\Box_{\psi^{\otimes\tau} \vee \delta^{\otimes\tau}} \theta$. Then there is a finite sequence $\boldsymbol{\pi} = \pi_0, \dots, \pi_n$ with $\pi_n = \theta$ that is a proof of θ in the theory axiomatized by $\psi^{\otimes\tau} \vee \delta^{\otimes\tau}$. Each formula χ_i occurring in $\boldsymbol{\pi}$ that is not a consequence of previous formulas in the sequence through a rule satisfies either $\psi^{\otimes\tau}$ or $\delta^{\otimes\tau}$. Then we have in particular that either $\Box_{\psi^{\otimes\tau}} \chi_i$ or $\Box_{\delta^{\otimes\tau}} \chi_i$ for each such χ_i , whence by the induction hypothesis either $\Box_T(\psi^{\otimes T} \rightarrow \chi_i)$ or $\Box_T(\delta^{\otimes T} \rightarrow \chi_i)$ holds. In both cases we have $\Box_T(\psi^{\otimes T} \wedge \delta^{\otimes T} \rightarrow \chi_i)$, and thus the proof of θ can be repeated in T under the assumption of $(\psi \wedge \delta)^{\otimes T}$, as desired.

(\leftarrow) Fix θ , \mathbf{y} , and \mathbf{z} . By the induction hypothesis (taking θ to be $\psi^{\otimes T}$) we see that $\Box_{\psi^{\otimes\tau}} \psi^{\otimes T}$ and likewise $\Box_{\delta^{\otimes\tau}} \delta^{\otimes T}$. Thus $\Box_{\psi^{\otimes\tau} \vee \delta^{\otimes\tau}} (\psi^{\otimes T} \wedge \delta^{\otimes T})$. By assumption we have $\Box_T(\psi^{\otimes T} \wedge \delta^{\otimes T} \rightarrow \theta)$, and since $\varphi^{\otimes\tau}$ extends T , we may conclude $\Box_{\varphi^{\otimes\tau}} \theta$ as desired.

The \forall case follows the same idea as the \wedge case. Consider $\varphi = \forall x_0 \psi$, with $\text{fv}(\psi) = \{x_0, x_1\}$ without loss of generality. Note that x_0 is represented by y_0 in T , and this is always a different variable from any z used to represent QRC_1 constants. As there is no further complication with constants, we omit them. Let θ and l be arbitrary and reason in T .

(\rightarrow) If $\Box_{\exists y_0 \psi^{\otimes\tau} [y_1 \leftarrow l]} \theta$, then there is a proof $\boldsymbol{\pi} = \pi_0, \dots, \pi_n$ where $\pi_n = \theta$ and each axiom χ_i in $\boldsymbol{\pi}$ satisfies $\psi^{\otimes\tau} [y_1 \leftarrow l] [y_0 \leftarrow k_i]$ for some number k_i , and consequently $\Box_T(\psi^{\otimes T} [y_1 \leftarrow l] [y_0 \leftarrow k_i] \rightarrow \chi_i)$ by the induction hypothesis for each i . Then by weakening we conclude $\Box_T(\forall y_0 \psi^{\otimes T} [y_1 \leftarrow l] \rightarrow \chi_i)$ for each i , and we are done.

(\leftarrow) Assume $\Box_T(\forall y_0 \psi^{\otimes T} [y_1 \leftarrow l] \rightarrow \theta)$. By Lemma 8.4 under the box, we obtain $\Box_T(\forall y_0 < m \psi^{\otimes T} [y_1 \leftarrow l] \rightarrow \theta)$, where m is the size of the domain of \mathcal{M} . Using the induction hypothesis for each $k < m$ with $\theta := \psi^{\otimes T} [y_1 \leftarrow l] [y_0 \leftarrow k]$, $y_0 := k$, and $y_1 := l$, we get $\forall k < m \Box_{\psi^{\otimes\tau} [y_1 \leftarrow l] [y_0 \leftarrow k]} \psi^{\otimes T} [y_1 \leftarrow l] [y_0 \leftarrow k]$, and in particular $\forall k < m \Box_{\exists y_0 \psi^{\otimes\tau} [y_1 \leftarrow l]} \psi^{\otimes T} [y_1 \leftarrow l] [y_0 \leftarrow k]$. Then by Σ_1^0 collection we can change the order of the quantifier and the box, concluding $\Box_{\exists y_0 \psi^{\otimes\tau} [y_1 \leftarrow l]} \forall y_0 < m \psi^{\otimes T} [y_1 \leftarrow l]$, which is enough to conclude this part of the proof.

Finally, for the case of $\varphi = \diamond\psi$, we start by observing that applying the induction hypothesis to \perp yields $T \vdash \forall \mathbf{y}, \mathbf{z} (\diamond_{\varphi^{\otimes\tau}} \top \leftrightarrow \diamond_T \varphi^{\otimes T})$. Note that $(\diamond\psi)^{\otimes\tau} = \tau(u) \vee (u = \ulcorner \diamond_{\psi^{\otimes\tau}} \top \urcorner)$, and thus $\Box_{(\diamond\psi)^{\otimes\tau}} \theta$ is equivalent to $\Box_T(\diamond_{\psi^{\otimes\tau}} \top \rightarrow \theta)$ by the formalized Deduction Theorem. The previous observation under the box then suffices to finish the proof. \dashv

We are finally ready to prove arithmetical completeness for any sound c.e. theory extending IS_1 .

THEOREM 8.9 (Arithmetical completeness). $\text{QRC}_1 \supseteq \text{QRC}_1(T)$.

PROOF. Recall the definition of $\text{QRC}_1(T)$:

$$\text{QRC}_1(T) = \{\varphi(\mathbf{x}, \mathbf{c}) \vdash \psi(\mathbf{x}, \mathbf{c}) \mid \forall \cdot^\circ T \vdash \forall \theta \forall \mathbf{y}, \mathbf{z} (\Box_{\psi^{\otimes\tau}} \theta \rightarrow \Box_{\varphi^{\otimes\tau}} \theta)\},$$

where θ is closed.

We show that if $\varphi \not\vdash \psi$, then the realization \cdot^\circledast defined above is such that

$$T \not\vdash \forall \theta \forall y, z (\Box_{\psi^\circledast \tau} \theta \rightarrow \Box_{\varphi^\circledast \tau} \theta).$$

Suppose towards a contradiction that T does prove this formula. Then by Lemma 8.8,

$$T \vdash \forall \theta \forall y, z (\Box_T(\psi^{\circledast T} \rightarrow \theta) \rightarrow \Box_T(\varphi^{\circledast T} \rightarrow \theta)).$$

Taking $\theta := \psi^{\circledast T}[y \leftarrow \ulcorner g_{\varphi, \psi}(x) \urcorner][z \leftarrow \ulcorner c^{I_0} \urcorner]$, $y := \ulcorner g_{\varphi, \psi}(x) \urcorner$, and $z := \ulcorner c^{I_0} \urcorner$, we conclude

$$T \vdash \Box_T(\varphi^{\circledast T} \rightarrow \psi^{\circledast T})[y \leftarrow \ulcorner g_{\varphi, \psi}(x) \urcorner][z \leftarrow \ulcorner c^{I_0} \urcorner].$$

This together with the soundness of T contradicts Theorem 8.7. ⊥

We have seen that $\text{QRC}_1 = \text{QRC}_1(T)$ for any sound c.e. theory T extending IS_1 . Thus $\text{QRC}_1(T)$ is constant over a large class of theories, and for these theories it does not depend on the specific axiomatization τ chosen. This is similar to the propositional case, but simpler than the full predicate case, where $\text{QPL}(T)$ is known to depend on both T (as shown by Montagna [32]) and τ (as shown by Artemov [3]; see also [28, 29]).

§9. A decidable fragment of QPL(PA). As we mentioned before, Vardanyan's results are very robust and Π_2^0 completeness can already be obtained for the language with just one unary predicate symbol and no nested occurrences of \Box , as shown in [37]. However, Artemov and Japaridze [4] managed to carve out a non-trivial decidable fragment of QPL(PA): all formulas that are decidable on finite Kripke frames correspond directly to a fragment of QPL(PA). They further observed that as a corollary one can conclude the decidability of the one variable fragment of QPL(PA).

The above may seem contradictory with Vardanyan's result on Π_2^0 completeness of the fragment of QPL(PA) with just one unary predicate symbol. However, although it is easy to see that in predicate logic any sentence with just one unary predicate is equivalent to one in the one-variable fragment, this does not hold when modalities are involved, as exhibited by the formula $\forall x \forall y \Box(P(x) \vee \neg P(y))$.

In this section we shall see that QRC_1 gives rise to a new decidable fragment of QPL(PA). Let $\mathcal{L}_{\Box, \forall}$ be the language of full quantified modal logic, based on the same signatures as the language of QRC_1 and extending the latter with \rightarrow, \perp , and the usual abbreviations. We extend the notion of finite arithmetical realization described in Definition 8.1 to $\mathcal{L}_{\Box, \forall}$ as follows:

- $\perp^{*PA} := \perp$;
- $(A \rightarrow B)^{*PA} := A^{*PA} \rightarrow B^{*PA}$.

We now define the quantified provability logic of PA as the set of always provable $\mathcal{L}_{\Box, \forall}$ formulas:

$$\text{QPL(PA)} := \{A(x, c) \in \mathcal{L}_{\Box, \forall} \mid \text{for any } \cdot^*, \text{ we have } \text{PA} \vdash \forall y, z A^{*PA}\},$$

where y are the arithmetical counterpart of x and z the arithmetical counterpart of c , as before.

THEOREM 9.1. *Let φ and ψ be QRC_1 formulas. Then*

$$\varphi \vdash \psi \iff \varphi \rightarrow \psi \in \text{QPL}(\text{PA}).$$

PROOF. The left-to-right implication is proved by induction on $\varphi \vdash \psi$. We note that the formalized converse Barcan formula is used in the case of the necessitation rule and provable Σ_1^0 completeness is used in the case of the transitivity axiom (see [11] for details on these results).

For the right-to-left implication, we take the contrapositive. Thus, let φ and ψ be strictly positive formulas with free variables x and constants c such that $\varphi \not\vdash \psi$. By Theorem 8.7, we have

$$\text{PA} \not\vdash (\varphi^{\otimes \text{PA}} \rightarrow \psi^{\otimes \text{PA}})[y \leftarrow \ulcorner \mathbf{g}_{\varphi, \psi}(\mathbf{x}) \urcorner][z \leftarrow \ulcorner \mathbf{c}^{I_0} \urcorner],$$

and thus $\varphi \rightarrow \psi \notin \text{QPL}(\text{PA})$. ⊣

§10. Heyting Arithmetic. We end this paper with a foray into intuitionistic arithmetic. We show that QRC_1 is sound with respect to Heyting Arithmetic (HA) and conjecture that it is complete as well.

Let $\eta(u)$ be a Σ_1^0 formula naturally axiomatizing HA.⁹ The HA-provability of φ can thus be expressed by $\Box_\eta \varphi$, as defined in Section 7.

We observe that a number of standard results in the realm of classical provability logic still hold in the intuitionistic case.

LEMMA 10.1 (Derivability conditions [39]). *Let τ be a Σ_1^0 axiomatization of an arithmetical theory T extending HA, and φ, ψ be formulas in the language of arithmetic. Then:*

1. if $T \vdash \varphi$ then $\text{HA} \vdash \Box_\tau \varphi$;
2. $\text{HA} \vdash \Box_\tau(\varphi \rightarrow \psi) \rightarrow (\Box_\tau \varphi \rightarrow \Box_\tau \psi)$;
3. $\text{HA} \vdash \Box_\tau \varphi \rightarrow \Box_\eta \Box_\tau \varphi$.

LEMMA 10.2 (Collection [21, Proposition 5.13]). *Let φ be an arithmetical formula without x as a free variable. Then*

$$\text{HA} \vdash \forall y < x \exists z \varphi \rightarrow \exists w \forall y < x \exists z < w \varphi.$$

Note that, since HA proves full collection, Prf_η is equivalent to a Σ_1^0 formula, provably in HA.

As in the classical case, we define $\Diamond_\tau \varphi$ as $\neg \Box_\tau \neg \varphi$ when τ is an axiomatization of an extension of HA.

We extend a generic realization \cdot° to non-predicate formulas as in the classical case (cf. Definition 7.1). Note that $\varphi^{\circ \eta}$ is equivalent to a Σ_1^0 formula and extends HA, both of these provably in HA.

THEOREM 10.3 (Arithmetical soundness w.r.t. HA).

$$\text{QRC}_1 \subseteq \{ \varphi \vdash \psi \mid \forall \cdot^\circ \text{HA} \vdash \forall \theta, \mathbf{y}, \mathbf{z} (\Box_{\varphi^{\circ \eta}} \theta \rightarrow \Box_{\psi^{\circ \eta}} \theta) \}.$$

⁹HA has less complex axiomatizations, but Σ_1^0 suffices for this section.

PROOF. Let φ and ψ be formulas such that $\varphi \vdash \psi$. The proof is similar to the classical case, and proceeds by external induction on $\varphi \vdash \psi$.

The soundness of $\varphi \vdash \top$ states that $\varphi^{\circ\eta}$ extends HA, which can readily be checked by induction on φ . The soundness of $\varphi \vdash \varphi$ and of the cut rule are straightforward, and the soundness of $\varphi \wedge \psi \vdash \varphi$ is a simple weakening.

We proceed with the soundness of the conjunction introduction rule, that if $\varphi \vdash \psi$ and $\varphi \vdash \chi$ then $\varphi \vdash \psi \wedge \chi$. Reason in HA and let θ be a closed formula, and \mathbf{y} and \mathbf{z} be arbitrary. Assume $\Box_{\psi^{\circ\eta} \vee \chi^{\circ\eta}} \theta$, taking $\boldsymbol{\pi} = \pi_0, \dots, \pi_{n-1}$ as a proof of this fact. Some of the π_i are axioms of $\psi^{\circ\eta}$, some are axioms of $\chi^{\circ\eta}$, and some follow from previous steps in the proof through a rule. Let $\{\delta_i\}_{i \in I}$ be the finite set of $\psi^{\circ\eta}$ axioms appearing in $\boldsymbol{\pi}$. Thus $\Box_{\psi^{\circ\eta}} \bigwedge_{i \in I} \delta_i$ and by the induction hypothesis for $\varphi \vdash \psi$ we obtain $\Box_{\varphi^{\circ\eta}} \bigwedge_{i \in I} \delta_i$. On the other hand, we know $\Box_{\chi^{\circ\eta}} (\bigwedge_{i \in I} \delta_i \rightarrow \theta)$ by the Deduction Theorem. Through the induction hypothesis for $\varphi \vdash \chi$ we conclude $\Box_{\varphi^{\circ\eta}} (\bigwedge_{i \in I} \delta_i \rightarrow \theta)$. Putting these two observations together, we obtain the desired $\Box_{\varphi^{\circ\eta}} \theta$.

We turn to the necessitation rule, that if $\varphi \vdash \psi$ then $\diamond\varphi \vdash \diamond\psi$. Reason in HA and let θ , \mathbf{y} , and \mathbf{z} be arbitrary. Consider the induction hypothesis with $\theta := \neg\top$ (and \mathbf{y} , \mathbf{z} as given by our assumption): $\Box_{\psi^{\circ\eta}} \neg\top \rightarrow \Box_{\varphi^{\circ\eta}} \neg\top$. Taking the contrapositive, we conclude $\diamond_{\varphi^{\circ\eta}} \top \rightarrow \diamond_{\psi^{\circ\eta}} \top$, and consequently $\Box_{\eta} (\diamond_{\varphi^{\circ\eta}} \top \rightarrow \diamond_{\psi^{\circ\eta}} \top)$ by Lemma 10.1. Assume now that $\Box_{(\diamond\psi)^{\circ\eta}} \theta$. By the Deduction Theorem, we obtain $\Box_{\eta} (\diamond_{\psi^{\circ\eta}} \top \rightarrow \theta)$. Thus our previous observation gives us $\Box_{\eta} (\diamond_{\varphi^{\circ\eta}} \top \rightarrow \theta)$, which is $\Box_{(\diamond\varphi)^{\circ\eta}} \theta$ by the Deduction Theorem.

Consider the transitivity axiom: $\diamond\diamond\varphi \rightarrow \diamond\varphi$. We start by observing that $(\diamond\diamond\varphi)^{\circ\eta}$ is equivalent to $\eta(u) \vee u = \ulcorner \diamond_{\eta} \diamond_{\varphi^{\circ\eta}} \top \urcorner$. Note that we can derive $\Box_{\eta} (\diamond_{\eta} \diamond_{\varphi^{\circ\eta}} \top \rightarrow \diamond_{\varphi^{\circ\eta}} \top)$ from Lemma 10.1. Assume now $\Box_{(\diamond\varphi)^{\circ\eta}} \theta$. By the Deduction Theorem we have $\Box_{\eta} (\diamond_{\varphi^{\circ\eta}} \top \rightarrow \theta)$. Then we obtain $\Box_{\eta} (\diamond_{\eta} \diamond_{\varphi^{\circ\eta}} \top \rightarrow \theta)$ by our previous observation, and we finish with one more application of the Deduction Theorem.

The \forall introduction rule on the right states that if $x_0 \notin \text{fv}(\varphi)$ and $\varphi \vdash \psi$, then $\varphi \vdash \forall x_0 \psi$. Reason in HA and let θ , \mathbf{y} , and \mathbf{z} be arbitrary, where y_0 does not appear in \mathbf{y} . Assume $\Box_{\exists y_0 \psi^{\circ\eta}} \theta$ and let $\boldsymbol{\pi} = \pi_0, \dots, \pi_{n-1}$ be a proof of this fact. Let $\{\delta_i\}_{i \in I}$ be the finite set of axioms of $\exists y_0 \psi^{\circ\eta}$ appearing in $\boldsymbol{\pi}$. Note that $\Box_{\eta} (\forall i \in I \delta_i \rightarrow \theta)$ holds by the Deduction Theorem. Then for each $i \in I$ there is k_i such that δ_i is an axiom of $\psi^{\circ\eta}[y_0 \leftarrow k_i]$, so in particular $\Box_{\psi^{\circ\eta}[y_0 \leftarrow k_i]} \delta_i$. We can use the induction hypothesis for each $i \in I$ to conclude $\Box_{\varphi^{\circ\eta}[y_0 \leftarrow k_i]} \delta_i$, and since $x_0 \notin \text{fv}(\varphi)$, we also know that $y_0 \notin \text{fv}(\varphi^{\circ\eta})$, and thus we obtain $\forall i \in I \Box_{\varphi^{\circ\eta}} \delta_i$. We now use collection to obtain $\Box_{\varphi^{\circ\eta}} \forall i \in I \delta_i$, and the result follows from our observation that $\Box_{\eta} (\forall i \in I \delta_i \rightarrow \theta)$, noting that $\varphi^{\circ\eta}$ extends HA.

The \forall introduction rule on the left states that if $\varphi[x_0 \leftarrow t] \vdash \psi$, then $\forall x_0 \varphi \vdash \psi$. Let w be the arithmetical counterpart of t (so if t is x_k then $w := y_k$ and if t is c_k then $w := z_k$). We have $\varphi[x_0 \leftarrow t]^{\circ\eta} = \varphi^{\circ\eta}[y_0 \leftarrow w]$. Let θ , \mathbf{y} , and \mathbf{z} be arbitrary, where y_0 appears in \mathbf{y} if and only if x_0 is a free variable of ψ . We assume $\Box_{\psi^{\circ\eta}} \theta$. If t appears in φ or in ψ , then the value of w was already fixed when we picked arbitrary \mathbf{y} and \mathbf{z} . Otherwise, fix $w := 0$ and use the induction hypothesis to obtain $\Box_{\varphi^{\circ\eta}[y_0 \leftarrow w]} \theta$. It is then clear that $\Box_{\exists y_0 \varphi^{\circ\eta}} \theta$ holds as well.

Finally, the term instantiation and constant elimination rules boil down to variable renaming. ⊣

The arithmetical completeness of QRC_1 with respect to HA remains an open question. We conjecture that the fact that the substitutions in the completeness proofs are of restricted complexity, and the fact that PA is Π_2^0 conservative over HA (see [20]) can be essential ingredients in a possible completeness proof.

§11. Future work. There are many unexplored paths surrounding QRC_1 . It would be worthwhile to extend it to a polymodal language (in analogy with RC, as in [6]), and to the positive setting (as in [13, 17]). Whether this is possible without loosing decidability remains to be seen. A hypothetical QRC_Λ would presumably lead to some interesting applications to Π_1^0 ordinal analysis and ordinal notation systems.

There are several proof theoretical properties of interest, such as interpolation, cut-free proofs, fixpoints, etc. One could also strive for uniform completeness.

Moreover, the set of always true QRC_1 sequents should be a productive avenue of study.

The completeness of QRC_1 with respect to HA remains an open question.

Finally, the current approximation for the computational complexity of QRC_1 is super-exponential space, since the canonical model grows quite large. A more dedicated study might lead to a significant reduction in complexity.

REFERENCES

- [1] A. DE ALMEIDA BORGES and J. J. JOOSTEN, *Quantified reflection calculus with one modality*, *Advances in Modal Logic*, vol. 13 (N. Olivetti, R. Verbrugge, S. Negri, and G. Sandu, editors), College Publications, Cambridge, 2020, pp. 13–32.
- [2] S. N. ARTEMOV, *Nonarithmeticity of truth predicate logics of provability*. *Doklady Akademii Nauk SSSR*, vol. 284 (1985), no. 2, pp. 270–271 (in Russian). English translation in *Soviet Mathematics Doklady*, vol. 33 (1985), pp. 403–405.
- [3] ———, *Numerically correct logics of provability*. *Doklady Akademii Nauk SSSR*, vol. 290 (1986), no. 6, pp. 1289–1292 (in Russian).
- [4] S. N. ARTEMOV and G. K. JAPARIDZE, *Finite Kripke models and predicate logics of provability*, *Journal of Symbolic Logic*, vol. 55 (1990), no. 3, pp. 1090–1098.
- [5] L. D. BEKLEMISHEV, *The worm principle*, *Logic Colloquium '02* (Z. Chatzidakis, P. Koepke, and W. Pohlers, editors), Lecture Notes in Logic, vol. 27, ASL Publications, Cambridge, 2006, pp. 75–95.
- [6] ———, *Calibrating provability logic: From modal logic to reflection calculus*, *Advances in Modal Logic*, vol. 9 (T. Bolander, T. Braüner, T. S. Ghilardi, and L. Moss, editors), College Publications, London, 2012, pp. 89–94.
- [7] ———, *Positive provability logic for uniform reflection principles*. *Annals of Pure and Applied Logic*, vol. 165 (2014), no. 1, pp. 82–105.
- [8] ———, *A note on strictly positive logics and word rewriting systems*, *Larisa Maximova on Implication, Interpolation, and Definability*, (S. Odintsov, editor), Outstanding Contributions to Logic, vol. 15, Springer, Berlin–Heidelberg, 2018, pp. 61–70.
- [9] LEV D. BEKLEMISHEV and FEDOR N. PAKHOMOV, *Reflection algebras and conservation results for theories of iterated truth*. *Annals of Pure and Applied Logic*, vol. 173 (2022), no. 5, pp. 103093.
- [10] A. BERARDUCCI, Σ_n^0 -interpretations of modal logic. *Bollettino dell'Unione Matematica Italiana*, vol. 7 (1989), no. 3-A, pp. 177–184.
- [11] G. S. BOLOS, *The Logic of Provability*, Cambridge University Press, Cambridge, 1993.
- [12] G. S. BOLOS and V. R. MCGEE, *The degree of the set of sentences of predicate provability logic that are true under every interpretation*, *Journal of Symbolic Logic*, vol. 52 (1987), pp. 165–171.

- [13] S. CELANI and R. JANSANA, *A note on the model theory for positive modal logic*. *Fundamenta Informaticae*, vol. 114 (2012), no. 1, pp. 31–54.
- [14] Coq Development Team, *The Coq Proof Assistant* (1989–2022), 2022.
- [15] W. CRAIG, *On axiomatizability within a system*, *Journal of Symbolic Logic*, vol. 18 (1953), pp. 30–32.
- [16] E. V. DASHKOV, *On the positive fragment of the polymodal provability logic GLP*. *Mathematical Notes*, vol. 91 (2012), nos. 3–4, pp. 318–333.
- [17] J. M. DUNN, *Positive modal logic*. *Studia Logica*, vol. 55 (1995), pp. 301–317.
- [18] S. FEFERMAN, *Arithmetization of metamathematics in a general setting*. *Fundamenta Mathematicae*, vol. 49 (1960), pp. 35–92.
- [19] D. FERNÁNDEZ-DUQUE, *Worms and spiders: Reflection calculi and ordinal notation systems*. *IfCoLoG Journal of Logics and their Applications*, vol. 4 (2017), no. 10, pp. 3277–3356.
- [20] H. FRIEDMAN, *Classically and intuitionistically provably recursive functions*. *Higher Set Theory* (D. S. Scott and G. H. Muller, editors), Lecture Notes in Mathematics, vol. 699, Springer, Berlin, 1978, pp. 21–28.
- [21] M. FUJIWARA and T. KURAHASHI, *Refining the arithmetical hierarchy of classical principles*, preprint, 2022, [arXiv:2010.11527](https://arxiv.org/abs/2010.11527) [math.LO].
- [22] K. GÖDEL, *Über formal unentscheidbare Sätze der Principia Mathematica und verwandter Systeme I*. *Monatshefte für Mathematik und Physik*, vol. 38 (1931), pp. 173–198.
- [23] P. HÁJEK and P. PUDLÁK, *Metamathematics of First Order Arithmetic*. Springer, Berlin–Heidelberg–New York, 1993.
- [24] Y. HAO and G. TOURLAKIS, *An arithmetically complete predicate modal logic*. *Bulletin of the Section of Logic*, vol. 50 (2021), no. 4, pp. 513–541.
- [25] G. E. HUGHES and M. J. CRESSWELL, *A New Introduction to Modal Logic*. Routledge, London, 1968.
- [26] G. K. JAPARIDZE, *The modal logical means of investigation of provability*. Ph.D. thesis, Moscow State University, Moscow, 1986 (in Russian).
- [27] S. KIKOT, A. KURUCZ, Y. TANAKA, F. WOLTER, and M. ZAKHARYASCHEV, *Kripke completeness of strictly positive modal logics over meet-semilattices with operators*. *Journal of Symbolic Logic*, vol. 84 (2019), no. 2, pp. 533–588.
- [28] T. KURAHASHI, *On predicate provability logics and binumerations of fragments of Peano Arithmetic*. *Archive for Mathematical Logic*, vol. 52 (2013), pp. 871–880.
- [29] ———, *On inclusions between quantified provability logics*. *Studia Logica*, vol. 110 (2022), pp. 165–188.
- [30] A. KURUCZ, F. WOLTER, and M. ZAKHARYASCHEV, *Islands of tractability for relational constraints: towards dichotomy results for the description logic EL*. *Advances in Modal Logic*, vol. 8 (L. D. Beklemishev, V. Goranko, and V. Shehtman, editors), College Publications, Moscow, 2010, pp. 271–291.
- [31] V. R. MCGEE, *Truth and necessity in partially interpreted languages*. Ph.D. thesis, University of California, Berkeley, 1985.
- [32] F. MONTAGNA, *The predicate modal logic of provability*. *Notre Dame Journal of Formal Logic*, vol. 25 (1984), no. 2, pp. 179–189.
- [33] I. SHAPIROVSKY, *PSPACE-decidability of Japaridze's polymodal logic*. *Advances in Modal Logic*, vol. 7 (C. Areces and R. Goldblatt, editors), College Publications, Nancy, 2008, pp. 289–304.
- [34] R. M. SOLOVAY, *Provability interpretations of modal logic*. *Israel Journal of Mathematics*, vol. 28 (1976), pp. 33–71.
- [35] A. TARSKI, *The concept of truth in formalized languages*. *Logic, Semantics and Metamathematics*, Hackett, 1983, pp. 152–278, English translation of Tarski's 1936 *Der Wahrheitsbegriff in den Formalisierten Sprachen*.
- [36] V. A. VARDANYAN, *Arithmetic complexity of predicate logics of provability and their fragments*. *Doklady Akad. Nauk SSSR*, vol. 288 (1986), no. 1, pp. 11–14 (in Russian). English translation in *Soviet Mathematics Doklady*, vol. 33 (1986), pp. 569–572.
- [37] ———, *Bounds on the arithmetical complexity of predicate logics of provability*. *Questions of Cybernetics: Complexity of Computation and Applied Mathematical Logic* (S. N. Adyan, editor), vol. 134, Scientific Council of the USSR Academy of Sciences on the complex problem “Cybernetics”, Moscow, 1988, pp. 46–72 (in Russian).

- [38] A. VISSER and M. DE JONGE, *No escape from Vardanyan's theorem*. *Archive for Mathematical Logic*, vol. 45 (2006), no. 5, pp. 539–554.
- [39] A. VISSER and J. ZOETHOUT, *Provability logic and the completeness principle*. *Annals of Pure and Applied Logic*, vol. 170 (2019), no. 6, pp. 718–753.
- [40] R. E. YAVORSKY, *On arithmetical completeness of first-order logics of provability*, *Advances in Modal Logic*, vol. 3 (F. Wolter, H. Wansing, M. de Rijke, and M. Zakharyashev, editors), World Scientific, Leipzig, 2002, pp. 1–16.

PHILOSOPHY DEPARTMENT
UNIVERSITY OF BARCELONA
08001 BARCELONA, SPAIN

E-mail: ana.agvb@gmail.com

URL: aborges.eu

E-mail: jjoosten@ub.edu

URL: joostjjoosten.nl