

46. Centers for Disease Control and Prevention. Outbreaks of pneumococcal pneumonia among unvaccinated residents in chronic-care facilities—Massachusetts, October 1995, Oklahoma, February 1996, and Maryland, May-June 1996. *MMWR* 1997;46:60-62.
47. McNeeley DF, Lyons J, Conte S, Labowitz A, Layton M. A cluster of drug-resistant *Streptococcus pneumoniae* among nursing home patients. *Infect Control Hosp Epidemiol* 1998;19:476-477.
48. Centers for Disease Control and Prevention. Surveillance for penicillin-nonsusceptible *Streptococcus pneumoniae*—New York City, 1995. *MMWR* 1997;46:297-299.
49. Centers for Disease Control and Prevention. Prevalence of penicillin-resistant *Streptococcus pneumoniae*—Connecticut, 1992-1993. *MMWR* 1994;43:216-217,223.
50. Butler JC, Hofmann J, Cetron MS, Elliott JA, Facklam RR, Breiman RF. The continued emergence of drug-resistant *Streptococcus pneumoniae* in the United States: an update from the Centers for Disease Control and Prevention's Pneumococcal Sentinel Surveillance System. *J Infect Dis* 1996;174:986-993.
51. Centers for Disease Control and Prevention. Recommendations of the Immunization Practices Advisory Committee. Pneumococcal polysaccharide vaccine. *JAMA* 1989;261:1265-1267.
52. US Public Health Service. *Healthy People 2000 Review, 1995-96*. Washington, DC: US Department of Health and Human Services, Public Health Service; 1996.
53. Beck-Sague C, Villarino E, Giuliano D, Welbel S, Latts L, Manangan LM, et al. Infectious diseases and death among nursing home residents: results of surveillance in 13 nursing homes. *Infect Control Hosp Epidemiol* 1994;15:494-496.
54. Beck-Sague C, Banerjee S, Jarvis WR. Infectious diseases and mortality among US nursing home residents. *Am J Public Health* 1993;83:1739-1742.
55. Kunin CM. Care of the urinary catheter. In: Kunin CM, ed. *Detection, Prevention and Management of Urinary Tract Infections*. Philadelphia, PA: Lea and Febiger; 1972.
56. Wong ES. Guideline for prevention of catheter-associated urinary tract infections. *Am J Infect Control* 1983;11:28-31.
57. Nicolle LE, Muir P, Harding GK, Norris M. Localization of urinary tract infection in elderly, institutionalized women with asymptomatic bacteriuria. *J Infect Dis* 1988;157:65-70.
58. Nicolle LE, Brunka J, McIntyre M, Murray D, Harding GK. Asymptomatic bacteriuria, urinary antibody, and survival in the institutionalized elderly. *J Am Geriatr Soc* 1992;40:607-613.
59. Nordenstam GR, Brandberg CA, Oden AS, Svanborg Eden CM, Svanborg A. Bacteriuria and mortality in an elderly population. *N Engl J Med* 1986;314:1152-1156.
60. Kunin CM, Douthitt S, Dancing J, Anderson J, Moeschberger M. The association between the use of urinary catheters and morbidity and mortality among elderly patients in nursing homes. *Am J Epidemiol* 1992;135:291-301.
61. Garner JS, Favero MS. CDC Guideline for handwashing and hospital environmental control, 1985. *Infect Control* 1986;7:231-243.
62. Larson EL. APIC guideline for handwashing and hand antisepsis in health care settings. *Am J Infect Control* 1995;23:251-269.
63. Thompson BL, Dwyer DM, Ussery XT, Denman S, Vacek P, Schwartz B. Handwashing and glove use in a long-term-care facility. *Infect Control Hosp Epidemiol* 1997;18:97-103.
64. Crossley K, SHEA Long-Term Care Committee. Vancomycin-resistant enterococci in long-term-care facilities. *Infect Control Hosp Epidemiol* 1998;19:521-525.
65. Nicolle LE, Bentley D, Garibaldi R, Neuhaus E, Smith P. Antimicrobial use in long-term-care facilities. *Infect Control Hosp Epidemiol* 1996;17:119-128.
66. Smith PW, Seip CW, Schaefer SC, Bell-Dixon C. Microbiologic survey of long-term care. 36th Annual Meeting of the Infectious Diseases Society of America, November 1998; Denver, CO. Abstract 6435a.
67. Mylotte JM. Antimicrobial prescribing in long-term care facilities: prospective evaluation of potential antimicrobial use and cost indicators. *Am J Infect Control* 1999;27:10-19.
68. Zimmer JG, Eggert GM, Treat A, Brodows B. Nursing homes as acute care providers. A pilot study of incentives to reduce hospitalizations. *J Am Geriatr Soc* 1988;36:124-129.

Effectiveness of a Manual Disinfection Procedure in Eliminating HCV From Experimentally Contaminated Endoscopes

Gina Pugliese, RN, MS
Martin S. Favero, PhD

Transmission of hepatitis C virus (HCV) through endoscopy has been reported, but the implications as a public health concern remain controversial. Some hospital hygienists believe that extraordinary means of cleaning and disinfection must be used on instruments of endoscopy after they are exposed to patients with HCV infection. Others believe that current cleaning and disinfection procedures are conservative enough and do not have to be modified to eliminate the risk of HCV transmission. Chanzy and co-investigators from France conducted a study that investigated the degree to which a thorough manual cleaning-washing-disinfection procedure can

decontaminate all channels of a flexible submersible endoscope experimentally contaminated with HCV. To assess the accuracy of the method currently in use, the initial investigation focused on sampling effectiveness. Nine endoscopes were contaminated with high-titer HCV-positive plasma and flushed with 150 mL of sampling solution (distilled water) before disinfection. To assess the effectiveness of the disinfection procedure, the following sequence was performed on another 10 endoscopes: inoculation, disinfection, and sampling. After concentration, residual viruses were detected by means of RNA amplification with commercial assays.

The study showed that sampling alone can reduce viral titer to one-fourth its original value. Within the limits of this method, HCV RNA was

never detected by means of polymerase chain reaction after disinfection, whereas all internal amplification controls were positive. This reduction to less than 1/100,000 of original titer exceeds the criterion expected for the virucidal activity of disinfectants.

The authors conclude that the results of this *in vitro* experiment provided evidence that patient-to-patient endoscopic transmission of HCV can be reduced, if not eliminated, with the current mechanical cleaning-washing-disinfection procedure.

FROM: Chanzy B, Duc-Bin DL, Rousset B, Morand P, Morel-Baccard C, Marchetti B, et al. Effectiveness of a manual disinfection procedure in eliminating hepatitis C virus from experimentally contaminated endoscopes. *Gastrointest Endosc* 1999; 50:147-151.