

Effects of prepartum positive and negative dietary cation/anion differences on postpartum calcium concentration and risk factors for subclinical hypocalcemia in Holstein cows

Research Article

Cite this article: Sami M, Badieli A, Beiranvand H, Moosakhani F, Shaghayegh A and Ahmadi F (2024). Effects of prepartum positive and negative dietary cation/anion differences on postpartum calcium concentration and risk factors for subclinical hypocalcemia in Holstein cows. *Journal of Dairy Research* **91**, 190–194. <https://doi.org/10.1017/S0022029924000451>

Received: 3 February 2024

Revised: 17 April 2024

Accepted: 6 May 2024

Keywords:

Dairy farm systems; dairy production; hypocalcemia; mammary gland

Corresponding author:

Mehrdad Sami;

Email: Samimehrdad@gmail.com

Mehrdad Sami¹, Arya Badieli², Hamed Beiranvand^{3,4}, Farhad Moosakhani⁵,

Alireza Shaghayegh² and Farhad Ahmadi⁶

¹Large Animal Internal Medicine Specialist, Veterinarian in the Private Sector, Tehran, Iran; ²Department of Clinical Sciences, Faculty of Veterinary Medicine, Islamic Azad University, Karaj, Iran; ³Department of Animal and Poultry Sciences, College of Aburaihan, University of Tehran, Tehran, Iran; ⁴R&D Department of Chaltasian & Talise Asil Jahan Agro-Animal Husbandry Co., Varamin, Tehran, Iran; ⁵Department of Pathobiology, Faculty of Veterinary Medicine, Karaj Branch, Islamic Azad University, Karaj, Iran and ⁶Department of Eco-friendly Livestock Science, Institute of Green Bio Science and Technology, Seoul National University, Pyeongchang 25354, South Korea

Abstract

Insufficient data are available about the association between prepartum risk factors including dietary cation–anion difference (DCAD) status and postpartum outcomes of Holstein dairy cows within commercial management systems. The first objective of this experiment was to assess the association between postpartum serum calcium (Ca) dynamics and the risk of metritis development. The second objective was to identify the association of risk factors, including DCAD status of prepartum ration, prepartum serum macrominerals and parity, with the development of subclinical hypocalcemia (SCH) in two commercial herds. Herd A ($n = 32$) fed a negative DCAD close-up ration and herd B ($n = 30$) fed a positive DCAD close-up diet. A receiver operating characteristic (ROC) curve was run to evaluate the association of serum Ca concentration at 1, 2, and 4 DIM with the risk of developing metritis. A second ROC curve was also created to assess the association of prepartum serum Mg, P, and Ca concentration with the postpartum serum Ca concentration dichotomized into normocalcemic (>8.82 mg/dl) and subclinical hypocalcemic (≤ 8.82 mg/dl). A logistic regression model was created to assess prepartum DCAD status (negative vs. positive) and parity (classified into a 3-level variable as first, second and third or greater lactations) as potential predictors of SCH classification. Serum Ca concentration at DIM 4 was a significant predictor of metritis (area under the curve = 0.87; $P < 0.01$). Cows fed positive vs. negative prepartum DCAD diet were more likely to be classified as SCH at 4 DIM. Parity did not show a significant association with the classification of SCH at 4 DIM. Our results demonstrate the importance of the DIM of blood Ca concentration assessment in connection with the diagnosis of metritis, which is crucial for the most accurate categorization of SCH and the related risk factors.

Hypocalcemia occurs in cows when their blood calcium (Ca) concentration decreases below the normal physiological range due to their inability to adjust to the sudden increase in Ca demand following calving (Seely *et al.*, 2021). Although data from modern dairy farms indicate a low incidence of clinical hypocalcemia (Goff, 2008), subclinical hypocalcemia (SCH) is still prevalent in commercial dairy farms, affecting a significant proportion of early-lactation cows (McArt and Neves, 2020; Serrenho *et al.*, 2021).

Owing to the considerable variability in blood Ca dynamics within a few days after calving, a single blood sample obtained shortly after calving for SCH classification may be a poor predictor of postpartum illness (Neves *et al.*, 2018; Seely *et al.*, 2021; McArt and Neves, 2020). A temporary drop in blood Ca may occur immediately after parturition as a normal physiological response to the sudden surge in Ca demand (McArt and Neves, 2020). In contrast to transient blood Ca reductions, delayed or persistent blood Ca reductions may indicate failure to maintain the Ca homeostatic system and, thus, be a better predictor of postpartum diseases, notably metritis in dairy cows (Seely *et al.*, 2021). For example, Neves *et al.* (2018) reported that in primiparous Holstein cows, there was no association between blood Ca concentration at 1 DIM and the risk of metritis development. However, they identified a significant association between blood Ca concentration at 2, 3, and 4 DIM and the risk of metritis. More information is required, particularly under commercial management systems, to identify the most reliable time point for SCH diagnosis (Rodríguez *et al.*, 2017; Venjakob *et al.*, 2018; McArt and Neves, 2020).

Creation of a negative dietary cation–anion difference (DCAD) by adding anionic salts to a close-up diet has been adopted as an effective nutritional strategy to help alleviate Ca deficit, and thus prevent clinical and subclinical hypocalcemia in dairy cows (Megahed *et al.*, 2018; Lean *et al.*, 2019; Santos *et al.*, 2019). Feeding an acidogenic diet causes the blood pH to decrease, allowing the bones to release Ca into the extracellular fluid to counteract the high anion concentration in the blood (Goff and Horst, 2003; Caixeta and Omontese, 2021). Martinez *et al.* (2018) reported a 30% reduction in the incidence of SCH in cows fed negative vs. positive DCAD diets during the prepartum period. However, less information is available to assess the association of DCAD status of prepartum diet and SCH classification according to DIM of blood Ca assessment. The first objective of this investigation was to identify the association between serum Ca concentration at 1, 2, and 4 DIM and the likelihood of cows being diagnosed with metritis. The second objective was to assess the association of risk factors, including prepartum serum magnesium (Mg), phosphorus (P) and Ca concentration as well as the DCAD status of the close-up diet and parity with SCH dichotomized into normocalcemic (>8.82 mg/dl) and subclinical hypocalcemic (≤ 8.82 mg/dl; Seely *et al.*, 2021; McArt and Neves, 2020; McArt and Oetzel, 2023).

Materials and methods

Herd description, analysis, and definitions

Two commercial Holstein herds (A and B) participated in data collection from July to September 2018. The herds were located in the central north of Iran (Tehran province). The second herd had originated from the first herd primarily because of stocking density challenges. Thus, both herds shared a similar genetic background among their lactating cows. Both herds had consistent management practices of having free-stall housing with a total mixed ration feeding program as well as a robust herd health program, including regular veterinary care, vaccination protocols, and disease prevention measures. Cows chosen for the trial met the criteria of being Holstein breed, aged between 2 and 8 years old, without major chronic health issues, not being introduced to the herd within the last 6 months and having consistent diet and management system for the last 6 months. In total, 62 cows (19 primiparous and 43 multiparous Holstein cows) were recruited for this study (herd A $n=32$, herd B $n=30$). During the prepartum period, herd A received a total mixed ration with a negative DCAD (-34 mEq/kg of DM), which falls within the typical negative DCAD diet for Iranian commercial dairy farms (usually ranging from 0 to -50 mEq/kg of DM). Ammonium sulfate and calcium chloride were used as anionic salts in the close-up diet. Herd B received a total mixed ration with a positive DCAD ($+201$ mEq/kg of DM) and did not include any anionic salts. DCAD (mEq/kg) was calculated using the formula $(\text{Na} + \text{K}) - (\text{Cl} + \text{S})$ (Sanchez *et al.*, 1994). Both herds were fed their respective diet once daily in the morning. Ingredients and chemical composition of prepartum diets formulated using NRC (2001) are listed in online Supplementary Table S1. Both farms A (2300 cows) and B (1800 cows) had a milking frequency of 3 times/d.

Blood was drawn from the coccygeal vein using a 20-gauge Vacutainer needle into an evacuated tube containing no anticoagulant (FL Medical, Torreglia, Italy) from approximately 1 d before the expected parturition date and then at 1, 2, 4, 6, 8, or 10 d after calving. The collected pre-calving blood samples were used for measurement of Mg, P, and Ca. The collected postpartum

blood samples were analyzed for Ca only. After centrifugation at $2000 \times g$ for 15 min, serum was recovered and stored at -20°C until mineral analysis. The total Ca, P, and Mg concentrations were determined using commercial kits (Pars Azmoon Co. Tehran, Iran) and an automated analyzer (RA-1000, Pharmacia Co., LKB, Novaspec, USA). The intra- and inter-assay coefficient of variations were less than 5%. Body condition score (BCS) was determined using a 5-point scale (Wildman *et al.*, 1982) and categorized into a 3-level variable as: thin: $\text{BCS} \leq 2.75$; normal: $3.0 \leq \text{BCS} \leq 3.5$; over-conditioned: $\text{BCS} \geq 3.75$ (Neves *et al.*, 2018). Parity was classified into a 3-level variable as first, second, and third or greater lactations. The case definition of the disease in both farms was established through the active participation and expertise of the first author, a qualified veterinarian, ensuring a standardized and consistent approach to identifying the diseases. Calving ease was also classified into a 2-level variable as cows with no or negligible assistance or suffering from dystocia. Failure to discharge fetal membranes within 24 h of calving was characterized as retained placenta (Qu *et al.*, 2014). Farm veterinarians checked uterine health, and metritis was diagnosed when reddish to brownish discharges were evident, often accompanied by symptoms of systemic diseases, including elevated rectal temperature ($\geq 39.5^\circ\text{C}$) or fever (Sheldon *et al.*, 2006).

Data analysis

Data of serum Ca collected over time from the two herds were analyzed using the MIXED Proc of SAS (SAS 9.4, SAS Institute Inc., Cary, NC) to detect the main effects of prepartum DCAD status, time (blood sampling relative to calving), and the DCAD-by-time interaction, with time variable specified in the REPEATED statement. Three covariance structures, namely Autoregressive type 1, Toeplitz, and Compound symmetry were tested to account for within-cow variations. The lowest Akaike's information criterion was achieved using Autoregressive type 1 and was selected. A significant difference ($P \leq 0.05$) was identified using the PDIFF option with a Tukey's test. The influence of prepartum DCAD status on the incidence of dystocia, retained placenta, or metritis was assessed using the GENMOD Proc in SAS, employing a Binomial distribution and logit link function. A ROC curve was constructed using SPSS (IBM SPSS Statistics, Version 24.0 Armonk, NY, USA, INM Corp., USA) to identify whether serum Ca concentration at 1, 2, and 4 DIM (continuous variables) could be a potential predictor of occurring metritis dichotomized into 0 = no disease and 1 = diseased. A second ROC analysis was also built to identify whether prepartum serum Mg, P, and Ca concentrations (continuous variables) were associated with SCH at 4 DIM (the classification variable). Continuous variables underwent a normality check, but no transformation was required. A multivariate binary logistic regression model was constructed in SAS to assess the association of prepartum DCAD status (negative vs. positive DCAD) and parity (a 3-level variable) with odds of developing SCH classification at 4 DIM based on the cut-off point of 8.82 mg/dl. The models initially included the BCS category at calving as a covariate, however, it was non-significant and removed from the model. Odds ratios with confidence intervals (95%) are reported for logistic regression models.

Results and discussion

Table 1 describes group proportions of BCS and parity in 62 recruited Holstein cows. Table 2 reports the effect of prepartum

Table 1. Description of group proportions of body condition score (BCS) and parity in 62 Holstein cows from two commercial herds

Items	Prepartum DCAD status ^a	
	Negative	Positive
BCS classification at calving, % (n)		
BCS ≤ 2.75 (thin)	3.13 (1)	6.67 (2)
3.0 ≤ BCS ≤ 3.5 (normal)	50.0 (16)	40.0 (12)
BCS ≥ 3.75 (over-conditioned)	46.9 (15)	53.3 (16)
Parity classification, % (n)		
1	28.1 (9)	26.7 (8)
2	28.1 (9)	23.3 (7)
≥3	43.8 (14)	50.0 (15)

^aThe negative DCAD ration was fed in herd A (n = 32), and the positive DCAD ration was fed in herd B (n = 30).

Table 2. Effect of prepartum DCAD status on prepartum serum Mg and P concentrations (mean ± standard deviation) and disease incidence in 62 Holstein cows from two commercial herds


Items	Prepartum DCAD status*		P value
	Negative	Positive	
Prepartum serum macrominerals ^a , mg/dl			
Mg	2.57 ± 0.37	2.42 ± 0.31	0.11
P	6.18 ± 1.19	5.89 ± 1.08	0.33
Disease incidence, % (n)			
Dystocia ^b	3.13 (1)	10.0 (3)	0.26
Retained placenta ^c	9.38 (3)	6.67 (2)	0.69
Metritis	12.5 (4)	10.0 (3)	0.76

^aPrepartum blood was collected approximately 24 h before the expected parturition.

^bCalving ease was classified as a 2-level variable, with cows requiring no or negligible assistance or suffering from dystocia.

^cFailure to expel fetal membranes within 24 h after calving was defined as retained placenta.

* The negative DCAD ration was fed in Herd A (n = 32), and the positive DCAD ration was fed in Herd B (n = 30).

Figure 1. Serum Ca concentration at different time points from 1 d before the expected calving date to 10 DIM in Holstein cows from two commercial herds. The negative DCAD ration (solid line) was fed in Herd A (n = 32). The positive DCAD ration (dashed line) was fed in herd B (n = 30). Time indicates serum Ca changes over time. An asterisk at each time point designates a statistical difference (P < 0.05). Error bars indicate standard error. There were significant effects of DCAD status, time and their interaction (P ≤ 0.01).

DCAD status on prepartum serum Mg and P, and the incidence of dystocia, retained placenta, and metritis. Prepartum serum Mg and P concentrations were not different between cows fed prepartum negative vs. positive DCAD close-up diet. Prepartum DCAD status also had no effect on the incidence of dystocia, retained placenta, and metritis.

Serum Ca dynamics and association with metritis

The effect of prepartum DCAD status on serum Ca concentration at different time points is shown in Fig. 1. The DCAD-by-time interaction was significant ($P = 0.01$) as negative DCAD-fed cows had greater serum Ca concentration before calving (-1) and d 1 and 8 post-calving. Prepartum DCAD status had no effect on serum Ca concentration at other time points (2, 4, 6, and 10 DIM).

Association of serum Ca concentration at 1, 2, and 4 DIM with the risk of metritis diagnosis, and association of prepartum blood macrominerals (Mg, P, and Ca) with postpartum blood Ca concentration dichotomized into 2 categories of normocalcemic (>8.82 mg/dl) and subclinical hypocalcemic (≤ 8.82 mg/dl), is presented in Table 3. According to ROC curve analysis, serum Ca concentration at 4 DIM but not 1 and 2 was a significant predictor of metritis (AUC = 0.87; $P < 0.01$), implying the adequate accuracy of the diagnostic test (AUC = 0.7 to 0.9; Swets, 1988). A growing body of evidence indicates the important involvement of Ca metabolism in the development of metritis and the heightened vulnerability of hypocalcemic cows to this condition (Martinez *et al.*, 2012; Rodríguez *et al.*, 2017). Impairment of immune function in SCH cows diminishes muscle contraction, which is associated with the elevated risk of developing uterine diseases (Martinez *et al.*, 2012). Our findings support those of Neves *et al.* (2018), who reported that blood Ca concentration at 1 DIM did not show an association with the risk of both primiparous and multiparous cows being diagnosed with metritis, but blood Ca concentration at 2, 3, and 4 DIM was a significant predictor of metritis development only in primiparous cows.

Association of prepartum serum macrominerals with SCH classification

According to the ROC curve analysis, a non-significant numerical difference was identified between prepartum serum Mg and SCH at 4 DIM (AUC = 0.65; $P = 0.06$). Association between prepartum

Table 3. Association of total serum Ca concentration (continuous variable) at DIM 1, 2, and 4 with the risk of being diagnosed with metritis (categorized variable) and association of prepartum serum macrominerals (continuous variable) with subclinical hypocalcemia classification (DIM 4)^a in two commercial herds

Items	P value	AUC (95% CI)	Sensitivity	Specificity	% of cows below the threshold
DIM of serum Ca concentration					
1	0.10	–	–	–	–
2	0.15	–	–	–	–
4	<0.01	0.87 (0.75–0.98)	70.9	85.7	34.4
Prepartum serum macrominerals ^b					
Mg	0.06	0.65 (0.51–0.79)	48.7	69.6	56.2
P	0.69	–	–	–	–
Ca	0.21	–	–	–	–

AUC, area under the curve.

^aDichotomization of serum Ca concentration (4 DIM) was based on the optimized Ca cut-off point from ROC curve analysis. Cows were classified into normocalcemic if total serum Ca concentration was >8.82 mg/dl at 4 DIM or subclinical hypocalcemia if total blood Ca concentration ≤8.82 mg/dl at 4 DIM.

^bPrepartum blood was collected approximately 24 h before the expected parturition.

Mg and postpartum SCH was anticipated as Mg status in the periparturient cow plays an important role in Ca metabolism, and hypomagnesemia has been recognized as a contributing factor for the development of clinical hypocalcemia (Lean *et al.*, 2006). Neves *et al.* (2017) also identified no link between prepartum blood Mg concentration and the risk of cows being categorized as SCH at calving. They suggested that their inability to identify any association may have originated from the low prevalence of prepartum subclinical hypomagnesemia as feeding prepartum rations containing Mg sources with proper bioavailability and adequate amounts may minimize the risk of developing hypomagnesemia and the subsequent effect on the SCH occurrence due to Mg deficient states. Although the cut-off point defining subclinical hypomagnesemia has not been characterized formally, a prepartum Mg concentration <1.95 mg/dl has been used for the classification of prepartum subclinical hypomagnesemia (Goff, 2008; Neves *et al.*, 2017). In this experiment, prepartum serum Mg concentration in cows fed negative and positive DCAD diets averaged 2.57 and 2.42 mg/dl, respectively, with only 3.03 and 3.23% of cows fed negative and positive DCAD diets having serum Mg concentrations below the cut-off point of <1.95 mg/dl. This low prevalence might not have had enough statistical power to establish a strong and reliable

association between prepartum serum Mg concentration and postpartum SCH classification. To precisely optimize the prepartum Mg cut-off point, additional information is required to identify whether the greater variability in prepartum Mg concentration may affect the risk of developing SCH.

The association of prepartum DCAD and parity with SCH dichotomized into 2 categories of normocalcemic (>8.82 mg/dl) and subclinical hypocalcemic (≤8.82 mg/dl) is listed in Table 4. Cows fed positive vs. negative DCAD prepartum ration had a numerically greater incidence of SCH classification at 4 DIM [odds ratio = 2.71; 95% confidence interval = 0.87–8.41; *P* = 0.09]. Our findings support the benefits of feeding a negative-DCAD prepartum diet that reduces the risk of low serum Ca concentration postpartum. In agreement, dairy cow studies have reported that a reduction in DCAD of prepartum diet from 300 to 0 mEq/kg resulted in reduced risk of being diagnosed with clinical hypocalcemia (16.4 to 3.2%) and increased blood ionized Ca pool before and after parturition (Charbonneau *et al.*, 2006; Lean *et al.*, 2006). This thereby contributed to the diminished risk of postpartum diseases such as retained placenta and metritis (Charbonneau *et al.*, 2006; Lean *et al.*, 2019).

Parity was not found to be a significant predictor of SCH classification at 4 DIM, which is consistent with the findings of Neves

Table 4. Final logistic regression models evaluating the association of DCAD^a status of prepartum diet (negative DCAD is reference point) and parity (primiparity is reference point) with subclinical hypocalcemia classification (normocalcemia is reference point)^b in Holstein cows from two commercial herds

Items	Incidence, %	Estimate	SE ^c	Odds ratio	95% confidence interval	P value
Prepartum DCAD status						
Negative	21.9			Referent		
Positive	43.3	0.99	0.57	2.71	0.87 to 8.41	0.09
Parity						
1	35.3			Referent		0.81
2	25.0	−0.48	0.79	0.62	0.13 to 2.97	
≥3	34.5	−0.09	0.66	0.92	0.25 to 3.43	

^aDCAD, dietary cation–anion difference.

^bCows were classified into normocalcemic if the total serum Ca concentration at 4 DIM was >8.82 mg/dl or subclinical hypocalcemia if the total serum Ca concentration at 4 DIM was ≤8.82 mg/dl.

^cSE, standard error of the estimate.

et al. (2017), who reported that parity was a significant predictor of SCH classification at the immediate postpartum period (within 4 h of parturition) but not 2 DIM. The physiological processes related to Ca metabolism and vitamin D receptor function tend to decline with age in cows, particularly in older parity individuals. Consequently, older parity cows exhibit reduced capacity to adequately maintain Ca turnover rates during the immediate postpartum period (Horst et al., 1990; Neves et al., 2017). However, as the DIM of blood Ca analysis advances, this effect becomes less pronounced.

It is important to highlight that this experiment suffers from a limitation in that the categorization of positive and negative DCAD-fed cows from two different herds might have introduced confounding variables into the results. Despite our initial attempts to minimize the herd-associated confounding by selecting the second herd that had originally emerged from the first herd (cows with comparatively similar genetic backgrounds), this approach may not have fully eliminated the residual confounding related to herd-to-herd variations. Future research should prioritize controlled trials, enabling greater control over variables for more accurate conclusions.

In conclusion, our findings validate distinct differences in the relationship between the timing of blood sampling relative to parturition and the risk of Holstein cows being diagnosed with metritis. We identified that serum Ca concentration at 4 DIM demonstrated a significant and consistent association with the development of metritis. Serum prepartum Mg but not P and Ca tended to be a significant predictor of SCH classification based on a serum Ca threshold at 4 DIM (8.82 mg/dl). Feeding a prepartum diet with a positive DCAD compared to a negative DCAD was associated with higher odds of being classified as SCH at 4 DIM. This study provided herd-level information about the DIM of blood Ca assessment for more accurate categorization of SCH and the associated risk factors that could help set targets for preventative strategies under commercial farm management conditions.

Supplementary material. The supplementary material for this article can be found at <https://doi.org/10.1017/S0022029924000451>

References

- Caixeta LS and Omontese BO (2021) Monitoring and improving the metabolic health of dairy cows during the transition period. *Animals* **11**, 352.
- Charbonneau E, Pellerin D and Oetzel GR (2006) Impact of lowering dietary cation-anion difference in nonlactating dairy cows: a meta-analysis. *Journal of Dairy Science* **89**, 537–548.
- Goff JP (2008) The monitoring, prevention, and treatment of milk fever and subclinical hypocalcemia in dairy cows. *The Veterinary Journal* **176**, 50–57.
- Goff JP and Horst RL (2003) Role of acid-base physiology on the pathogenesis of parturient hypocalcaemia (milk fever) – the DCAD theory in principal and practice. *Acta Veterinaria Scandinavica* **97**, 51–56.
- Horst RL, Goff JP and Reinhardt TA (1990) Advancing age results in reduction of intestinal and bone 1, 25-dihydroxyvitamin D receptor. *Endocrinology* **126**, 1053–1057.
- Lean IJ, DeGaris PJ, McNeil DM and Block E (2006) Hypocalcemia in dairy cows: meta-analysis and dietary cation anion difference theory revisited. *Journal of Dairy Science* **89**, 669–684.
- Lean IJ, Santos JEP, Block E and Golder HM (2019) Effects of prepartum dietary cation-anion difference intake on production and health of dairy cows: a meta-analysis. *Journal of Dairy Science* **102**, 2103–2133.
- Martinez N, Risco CA, Lima FS, Bisinotto RS, Greco LF, Ribeiro ES, Maunsell F, Galvão K and Santos JE (2012) Evaluation of periparturient calcium status, energetic profile, and neutrophil function in dairy cows at low or high risk of developing uterine disease. *Journal of Dairy Science* **95**, 7158–7172.
- Martinez N, Rodney RM, Block E, Hernandez LL, Nelson CD, Lean IJ and Santos JE (2018) Effects of prepartum dietary cation-anion difference and source of vitamin D in dairy cows: health and reproductive responses. *Journal of Dairy Science* **101**, 2563–2578.
- McArt JAA and Neves RC (2020) Association of transient, persistent, or delayed subclinical hypocalcemia with early lactation disease, removal, and milk yield in Holstein cows. *Journal of Dairy Science* **103**, 690–701.
- McArt JAA and Oetzel GR (2023) Considerations in the diagnosis and treatment of early lactation calcium disturbances. *Veterinary Clinics: Food Animal Practice* **39**, 241–259.
- Megahed AA, Hiew MW, El Badawy SA and Constable PD (2018) Plasma calcium concentrations are decreased at least 9 hours before parturition in multiparous Holstein-Friesian cattle in a herd fed an acidogenic diet during late gestation. *Journal of Dairy Science* **101**, 1365–1378.
- National Research Council (2001) *Nutrient Requirements of Dairy Cattle*, 7th rev. ed., Washington, DC: National Academy Press.
- Neves RC, Leno BM, Stokol T, Overton TR and McArt JAA (2017) Risk factors associated with postpartum subclinical hypocalcemia in dairy cows. *Journal of Dairy Science* **100**, 3796–3804.
- Neves RC, Leno BM, Bach KD and McArt JAA (2018) Epidemiology of subclinical hypocalcemia in early-lactation Holstein dairy cows: the temporal associations of plasma calcium concentration in the first 4 days in milk with disease and milk production. *Journal of Dairy Science* **101**, 9321–9331.
- Qu Y, Fadden AN, Traber MG and Bobe G (2014) Potential risk indicators of retained placenta and other diseases in multiparous cows. *Journal of Dairy Science* **97**, 4151–4165.
- Rodríguez EM, Aris A and Bach A (2017) Associations between subclinical hypocalcemia and postparturient diseases in dairy cows. *Journal of Dairy Science* **100**, 7427–7434.
- Sanchez WK, Beede DK and Delorenzo MA (1994) Macromineral element interrelationships and lactational performance: empirical models from a large data set. *Journal of Dairy Science* **77**, 3096–3110.
- Santos JEP, Lean IJ, Golder H and Block E (2019) Meta-analysis of the effects of prepartum dietary cation-anion difference on performance and health of dairy cows. *Journal of Dairy Science* **102**, 2134–2154.
- Seely CR, Leno BM, Kerwin AL, Overton TR and McArt JA (2021) Association of subclinical hypocalcemia dynamics with dry matter intake, milk yield, and blood minerals during the periparturient period. *Journal of Dairy Science* **104**, 4692–4702.
- Serrenho RC, DeVries TJ, Duffield TF and LeBlanc SJ (2021) Graduate student literature review: what do we know about the effects of clinical and subclinical hypocalcemia on health and performance of dairy cows? *Journal of Dairy Science* **104**, 6304–6326.
- Sheldon IM, Lewis GS, LeBlanc S and Gilbert RO (2006) Defining postpartum uterine disease in cattle. *Theriogenology* **65**, 1516–1530.
- Swets JA (1988) Measuring the accuracy of diagnostic systems. *Science (New York, N.Y.)* **240**, 1285–1293.
- Venjakob PL, Pieper L, Heuwieser W and Borchardt S (2018) Association of postpartum hypocalcemia with early-lactation milk yield, reproductive performance, and culling in dairy cows. *Journal of Dairy Science* **101**, 9396–9405.
- Wildman EE, Jones GM, Wagner PE, Boman RL, Troutt Jr HF and Lesch TN (1982) A dairy cow body condition scoring system and its relationship to selected production variables in high producing Holstein dairy cattle. *Journal of Dairy Science* **65**, 495–501.