

quality of life (QoL). However, how they impact QoL in long-term LT survivors is unclear.

Objectives: The study aimed to assess the associations between personality traits and cognition and their impact on the QoL in long-term LT survivors.

Methods: First time LT recipients due to end-stage alcohol liver disease without long-term complications were consecutively included during standard outpatient care. Sociodemographic and clinical data was collected. Personality traits were assessed using 50-item International Personality Item Pool of the Five-factor model (IPIP), cognition using Mini Mental State Examination (MMSE), and QoL using EuroQoL-5D (EQ-5D) questionnaire.

Results: Eighty-three participants were included (mean age 62.9 ±7.03y, 90.6% male). Median MMSE score was 27±2.00, and median years since LT 5±2.91. Significant positive associations were found between IPIP dimensions Extraversion (B=0.297, p<0.01), Agreeableness (B=0.384, p<0.01), Conscientiousness (B=0.511, p<0.01), and Emotional stability (B=0.432, p<0.01) with EQ-5D visual analogue scale (EQ-VAS). IPIP dimension Conscientiousness (B=0.338, p<0.01) and Emotional Stability (B=0.379, p<0.01) were significantly associated with descriptive dimension of EQ-5D (EQ-5D-3L). MMSE score was significantly associated with QoL (EQ-VAS B=0.291, p<0.01; EQ-5D-3L B=0.283, p<0.05, respectively). However, MMSE score was not shown to be a statistically significant predictor of QoL, whereas Conscientiousness was a significant predictor of EQ-VAS (β 1.404, t 3.125), and Emotional stability of EQ-5D-3L (β 0.011, t 2.132).

Conclusions: Some personality traits predicted QoL in long-term LT survivors. Therefore, assessment of personality traits should be considered as a part of pre-LT evaluation within a regular psychiatric clearance evaluation.

Disclosure of Interest: S. Medved: None Declared, B. Aukst Margetić: None Declared, A. Ražić Pavičić: None Declared, T. Filipec Kanižaj : None Declared, V. Medved Grant / Research support from: This work was done as a part of the “Genetic Background of End Stage Alcoholic Liver Disease and Liver Transplantation” study supported by the University of Zagreb, Croatia grant 2017 and 2018.

Post-Traumatic Stress Disorder

EPP0509

Prevalence and predictors of PTSD and resilience among Adolescents and Young Adults: Findings from the MoreGoodDays Support Program in Alberta, Canada

A. Belinda^{1*}, R. Shalaby¹, K. Hay², R. Pattison², E. Eboime³, M. Korhuis⁴, Y. Wei¹ and V. I. O. Agyapong^{1,3}

¹University of Alberta; ²Kickstand, Edmonton; ³Dalhousie University, Halifax and ⁴Glenrose Rehabilitation Hospital Foundation, Edmonton, Canada

*Corresponding author.

doi: 10.1192/j.eurpsy.2024.637

Introduction: Adolescents and young adults have particularly been impacted by the COVID-19 pandemic, leading to a rise in the incidence of mental health issues. Increased exposure to traumatic

events may lead to decreased resilience and subsequently increased likely PTSD.

Objectives: This study sets out to examine the predictors and prevalence of likely PTSD and determine the level of resilience among adolescents and young adults.

Methods: A cross-sectional study using an online survey questionnaire was adopted to collect sociodemographic and clinical information from the subscribers of MoreGoodDays. The PTSD Checklist Civilian (PCL-C) and the Brief Resilience Scale (BRS) was respectively used to assess likely PTSD and resilience Data was analyzed with SPSS version 25 using chi-squared tests and multivariate logistic regression analysis.

Results: 343 of MoreGoodDays subscribers who participated in the survey were about 343. Most were female (79.0%), and 13.7% were male. Overall, 95 (45.7%) of respondents had likely PTSD and 109 (51.7%) had likely low resilience. Approximately 176 (51.3%) respondents had received mental health counselling, and 64 (35.4%) expressed the desire to receive mental health counselling. When all other variables are controlled in the regression model, respondents who have received mental health counselling in the past year were 13.7 times more likely to experience likely PTSD (OR = 13.70; 95% CI: 1.23- 142.86) and 15.15 times more likely to experience low resilience than those who did not (OR = 15.15; 95% CI: 1.46- 166.67). Again, those who would like to receive mental health counselling were 20.8 times more likely to experience PTSD than those who did not (OR = 20.76; 95% CI: 2.61- 165.401) and 29.4 times more likely to experience low resilience than those who did not (OR = 29.42; 95% CI: 3.31- 261.445). Finally, those with four or more ACE scores were 6.2 times more likely to experience likely PTSD than those who had zero scores (OR = 6.24; 95% CI: 1.46- 26.67).

Conclusions: MoreGoodDays subscribers were disproportionately affected by likely PTSD and low resilience, reflecting the devastating effect of the COVID-19 pandemic. Increased ACE has been linked to low resilience, which may also lead to a rise in mental health issues. Strategies to promote resilience may reduce the incidence of likely PTSD. Educational institutions may adopt innovative mental health interventions, including psychological interventions such as mobile text technology, to support the mental health of this cohort. Policymakers and government agencies are encouraged to give the mental health of young adults and youth more prominence on their agenda.

Disclosure of Interest: None Declared

EPP0510

PTSD, dissociative experiences, and depressive symptoms in a clinical sample of women who featured in pornographic productions

F. K. El Khoury¹, V. Hérouin^{2*} and M. Fareng²

¹Department of social epidemiology, Sorbonne Université, INSERM, IPLESP and ²Clinical Psychologist, Paris, France

*Corresponding author.

doi: 10.1192/j.eurpsy.2024.638

Introduction: The mental health of the people who featured in Pornographic Productions (PP) is underexamined. However, PP frequently involve unsimulated violent acts mostly experienced by