

Concepts in Disaster Medicine

Cite this article: Levy E, Alkan M and Gidron Y (2024). Painful Call About the Under-Reporting of Children's Pain During Humanitarian Crises. *Disaster Medicine and Public Health Preparedness*, 18, e213, 1–3
<https://doi.org/10.1017/dmp.2024.252>

Received: 26 July 2023
Revised: 19 October 2024
Accepted: 19 September 2024

Keywords:
pain; palliative care; children; humanitarian crises; disasters; stress

Corresponding author:
Einav Levy;
Email: levygaea@gmail.com

Painful Call About the Under-Reporting of Children's Pain During Humanitarian Crises

Einav Levy^{1,2} , Michael Alkan³ and Yori Gidron⁴ 

¹Dept. of Social Work, The Research Center for Innovation in Social Work, Tel-Hai College, Qiryat Shmona, Israel; ²The Israeli School of Humanitarian Action and Lucien Research Center, Tel Aviv, Israel; ³School of International Health, Ben-Gurion University, Beer-Sheva, Israel and ⁴Faculty of Welfare and Health Sciences, University of Haifa, Haifa, Israel

Abstract

Humanitarian crises often require urgent medical care to people of concern. Such medical aid includes assessing and treating acute medical needs and ongoing chronic health conditions. Among the people of concern there are children, who are often the most vulnerable population in humanitarian contexts because they often lack the experience, independence, and cognitive and verbal skills to deal with the ordeals they are facing. These limitations might prevent identification and diagnosis of pain. The under-diagnosis and under-treated pain by health care providers might be also due to the perceived urgency of more acute or life-threatening medical needs with limited medical equipment and personnel, lack of awareness, or assessment tools in such contexts. Additionally, due to issues of anonymity and lack of formal guidelines, there is a severe lack of standardized registration of children's pain conditions in humanitarian crises. Finally, acute pain is also a predictor of post-traumatic stress disorder, a common outcome in such disasters. We call on health care providers to use standardized scales to assess children's pain intensity, frequency, and duration, and to treat it appropriately. These will not only reduce children's physical suffering but may also prevent subsequent risk of PTSD.

In humanitarian settings, the priorities of assistance provision are often focused on urgent medical support. Within this realm, data collection and usage of validated measurements of medical conditions are generally low. Several organizational and personal reasons are associated with the failure to collect data in these settings. First, in many cases there is no data collection procedure as part of the organizational protocol. Yet, if such a procedure does exist, the mere sense of emergency does not always enable the medical staff to record its actions. Moreover, humanitarian agencies are often avoiding sharing or publishing data due to considerations of confidentiality, organizational concerns, awareness, or low quality of the data collected. Even if published, it is difficult to draw a policy from such publications due to the lack of standardization among the measures described in the publications. The above challenges were addressed by the World Health Organization (WHO) standards, enabling databases to be analyzed, compared, and validated. Yet few agencies will follow the standard reporting system developed by the WHO.¹

The phenomena of under-reporting and the lack of data may have more severe implications when it comes to the condition of pain.² Pain in these humanitarian contexts could be chronic or acute, due to disaster-related injuries or its consequences (migration, malnutrition, infections, etc). Generally, pain is defined by the International Association of the Study of Pain (IASP) as: "An unpleasant sensory and emotional experience associated with, or resembling that associated with, actual or potential tissue damage."³ In the general population, based on reviews, the prevalence of different types of pain is 10.6% for chronic widespread pain⁴, 52.2% for headaches, 14% for migraines⁵, and 3.9%–19.6% for chronic low back pain subject to age.⁶

Data on various types of pain and palliative care are even more difficult to obtain from published field reports compiled during or after humanitarian assistance. Several publications have shown under-reporting and under-treatment of pain in such contexts.^{7–12} This under-reporting is in line with the general perception around addressing pain as presented in a systematic review of a handful of programs that addressed palliative care or pain in humanitarian contexts.¹³

Within the described programs in that review, we failed to find a report addressing the subject of pain related to age, and specifically to pediatric pain. This leads to the conclusion that pain in children is a topic that has been overlooked and neglected as an important issue in humanitarian actions.

Neglecting pain assessment and treatment in children is a known phenomenon in general hospitals out of the humanitarian context as well.^{14–23} This phenomenon may be a result of children's ways of expressing pain, which are different from those of adults,^{17–24} or due to barriers held by the medical staff, such as unfamiliarity and discomfort with pediatrics, unfamiliarity with the protocols, insufficient clinical education, or concerns from adverse effects of analgesics.²⁵ As it comes to humanitarian settings, research shows that children are more vulnerable to

© The Author(s), 2024. Published by Cambridge University Press on behalf of Society for Disaster Medicine and Public Health, Inc. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted re-use, distribution and reproduction, provided the original article is properly cited.

consequences of disasters and conflicts because they often lack the experience, independence, and cognitive and verbal skills to deal with the ordeals they are facing. Therefore, these all impair children's pre-existing ability to cope with such stressful events.^{26–31} Such limited skills and the inability to verbalize their pain, especially among infants, might also affect the level and nature of reported pain. In such cases, children learn fast that expressing pain is futile and it does not lead to pain relief. This might lead to developing helplessness, particularly in prolonged and uncontrollable disasters. Such a sense of helplessness is part of the broader concept of catastrophizing, a set of negative pain-related cognitions, which is correlated with pain in general and more specifically with pediatric pain.³²

Another reason for under-reporting pain could be related to culture. In some cultures, or situations, children may have been told not to complain or cry because they should be brave or silent in cases of a consistent violent threat to their life and the lives of their close ones. Finally, observed and reported pain are not always a result of a measurable physical malfunction or injury, but rather a stress related symptom requiring psychosocial care rather than typical medical care.^{33–35} This requires adequate communication with children who speak other languages, which is not always possible in urgent humanitarian contexts.

Assessment of Pain

The science of assessing pain, algometry,³⁶ faces methodological challenges, partly because of imperfect inter-rater variability of pain. Among children, assessment of pain includes visual analogue scales (VAS) of pain with numbers or faces. The face scale is especially suitable for younger children and in cases of existence of a linguistic gap between the child and the health care providers. Some biases and inaccuracies using face scales emerge when using drawn faces because children may have difficulties in distinguishing the sensory dimension of pain from its affective dimension. In addition, using smiling faces in the scale can result in overestimation of pain levels.³⁷ Lastly, according to a meta-analysis of 40 studies (4 628 children), there is a moderate association between pain ratings of children and their caregiver and a weaker association between child and health care provider,³⁸ but in some studies the pattern was opposite.³⁹

Implications

Considering these difficulties, we make the following suggestions. First, health care teams involved in humanitarian actions should be aware of pain in children as a hidden burden. Second, this should be incorporated into staff training programs and policies prior to deployments. Third, humanitarian organizations should be familiar with the existing standards of palliative care and pain treatment. Moreover, these standards should be implemented into the protocols of the medical staff or clinics providing medical care. Fourth, tools to assess pain, such as VAS-pain and face scales, should be available to the teams. Fifth, following inconsistent findings concerning estimating children's pain, it is recommended that both caregivers and health care providers assess the child's pain levels. In addition, clinics should include a checklist for caregivers concerning the observed behavioral expression of their child's pain (e.g., grimaces, unusual body movements), especially for children with cognitive or linguistic difficulties. These would raise the general awareness of the medical staff about this challenge. Finally,

organizations should be held accountable for diagnosing, treating, and adequately reporting pain in children.

To conclude, we need to keep in mind that within the humanitarian context, children are often the most vulnerable population, often without the ability to express their pain and stress. Therefore, it is our obligation through good clinical practice and ethics to reduce agony in children as part of their medical care.

Data availability statement. No data are associated with this study.

Acknowledgements. No funding was received.

Competing interest. All authors declare no conflict of interest.

References

1. **Integrating Palliative Care and Symptoms into Response to Humanitarian Emergencies and Crises Guidebook.** WHO, 2018. Accessed October 13, 2022. <https://who.int/publications/i/item/9789241514469>
2. Rosa WE, Grant I, Knaul FM, et al. The value of alleviating suffering and dignifying death in war and humanitarian crises. *Lancet*. 2022;**399**(10334):1447–1450.
3. **IASP Announces Revised Definition of Pain**, 2023. Accessed January 13, 2023. <https://www.iasp-pain.org/publications/iasp-news/iasp-announces-revised-definition-of-pain/>
4. Mansfield KE, Sim J, Jordan JL, et al. A systematic review and meta-analysis of the prevalence of chronic widespread pain in the general population. *Pain*. 2016;**157**(1):55.
5. Stovner LJ, Hagen K, Linde M, et al. The global prevalence of headache: an update, with analysis of the influences of methodological factors on prevalence estimates. *J Headache Pain*. 2022;**23**(1):34.
6. Meucci RD, Fassa AG, Faria NM. Prevalence of chronic low back pain: systematic review. *Revista de Saude Publica*. 2015;49.
7. Doherty M, Power L, Petrova M, et al. Illness-related suffering and need for palliative care in Rohingya refugees and caregivers in Bangladesh: a cross-sectional study. *PLoS Med*. 2020;**17**(3):e1003011.
8. Joury E, Meer R, Chedid JC, et al. Burden of oral diseases and impact of protracted displacement: a cross-sectional study on Syrian refugee children in Lebanon. *Br Dent J*. 2021;1–5.
9. Klas J, Grzywacz A, Kulszo K, et al. Challenges in the medical and psychosocial care of the paediatric refugee—a systematic review. *Int J Res Pub Health*. 2022;**19**(17):10656.
10. Levy E, Alkan M, Shaul S, et al. Medical conditions and treatment in a transit camp in Serbia for Syrian, Afghani, and Iraqi migrants. *J Int Humanit Action*. 2017;**2**:1–6.
11. Naaman O, Yulevich A, Sweed Y. Syria civil war pediatric casualties treated at a single medical center. *J Pediatr Surg*. 2020;**55**(3):523–529.
12. Schnall AH, Hanchey A, Nakata N, et al. Disaster-related shelter surveillance during the Hurricane Harvey response—Texas 2017. *Disaster Med Public Health Prep*. 2020;**14**(1):49–55.
13. Schneider M, Pautex S, Chappuis F. What do humanitarian emergency organizations do about palliative care? A systematic review. *Med Conflict Surviv*. 2017;**33**(4):263–272.
14. Birnie KA, Chambers CT, Fernandez CV, et al. Hospitalized children continue to report undertreated and preventable pain. *Pain Res Manag*. 2014;**19**(4):198–204.
15. Eccleston C, Fisher E, Howard RF, et al. Delivering transformative action in paediatric pain: a Lancet Child & Adolescent Health Commission. *Lancet Child Adolesc Health*. 2021;**5**(1):47–87.
16. Grout RW, Thompson-Fleming R, Carroll AE, et al. Prevalence of pain reports in pediatric primary care and association with demographics, body mass index, and exam findings: a cross-sectional study. *BMC Pediatr*. 2018;**18**:1.
17. Hewes HA, Dai M, Mann NC, et al. Prehospital pain management: disparity by age and race. *Prehosp Emerg Care*. 2018;**22**(2):189–197.
18. Izsak E, Moore JL, Stringfellow K, et al. Prehospital pain assessment in pediatric trauma. *Prehosp Emerg Care*. 2008;**12**(2):182–186.

19. Moultrie F, Shriver A, Hartley C, et al. A universal right to pain relief: balancing the risks in a vulnerable patient population. *Lancet Child Adolesc Health*. 2019;3(2):62–64.
20. Murphy A, McCoy S, O'Reilly K, et al. A prevalence and management study of acute pain in children attending emergency departments by ambulance. *Prehosp Emerg Care*. 2016;20(1):52–58.
21. Sleeman KE, De Brito M, Etkind S, et al. The escalating global burden of serious health-related suffering: projections to 2060 by world regions, age groups, and health conditions. *Lancet Glob Health*. 2019;7(7):e883–892.
22. Knaul FM, Farmer PE, Krakauer EL, et al. Alleviating the access abyss in palliative care and pain relief—an imperative of universal health coverage: the Lancet Commission report. *Lancet*. 2018;391(10128):1391–1454.
23. Markenson D. Have we forgotten about the needs of children?. *Disaster Med Public Health Prep*. 2014;8(3):188–190.
24. Zieliński J, Morawska-Kochman M, Zatoński T. Pain assessment and management in children in the postoperative period: A review of the most commonly used postoperative pain assessment tools, new diagnostic methods and the latest guidelines for postoperative pain therapy in children. *Adv Clin Exp Med*. 2020;29(3):365–374.
25. Williams DM, Rindal KE, Cushman JT, et al. Barriers to and enablers for prehospital analgesia for pediatric patients. *Prehosp Emerg Care*. 2012;16(4):519–526.
26. Awuah WA, Ng JC, Mehta A, et al. Vulnerable in silence: paediatric health in the Ukrainian crisis. *Ann Med Surg*. 2022;104369.
27. Betancourt TS, Keegan K, Farrar J, et al. The intergenerational impact of war on mental health and psychosocial wellbeing: lessons from the longitudinal study of war-affected youth in Sierra Leone. *Confl Health*. 2020;14(1):1–8.
28. Jabbar SA, Zaza HI. Impact of conflict in Syria on Syrian children at the Zaatari refugee camp in Jordan. *Early Child Dev Care*. 2014;184(9-10):1507–1530.
29. Meiqari L, Hoetjes M, Baxter L, et al. Impact of war on child health in northern Syria: the experience of Médecins Sans Frontières. *Eur J Pediatr*. 2018;177:371–380.
30. Ramirez D, Haas SA. The long arm of conflict: how timing shapes the impact of childhood exposure to war. *Demography*. 2021;58(3):951–974.
31. Raskoff SZ, Thurm A, Miguel HO, et al. Pain research and children and adolescents with severe intellectual disability: ethical challenges and imperatives. *Lancet Child Adolesc Health*. 2023;7(4):288–296.
32. Friedrichsdorf SJ, Giordano J, Desai Dakoji K, et al. Chronic pain in children and adolescents: diagnosis and treatment of primary pain disorders in head, abdomen, muscles and joints. *Children*. 2016;3(4):42.
33. Ayede AI. Neonatal pain management in sub-Saharan Africa. *Lancet Child Adolesc Health*. 2020;4(10):713–714.
34. Hamdan-Mansour AM, Abdel Razeq NM, AbdulHaq B, et al. Displaced Syrian children's reported physical and mental wellbeing. *Child Adolesc Ment Health*. 2017;22(4):186–193.
35. Nelson S, Miller JV, Timmers I, et al. Paediatric chronic pain as a catalyst for toxic stress. *Lancet Child Adolesc Health*. 2022;6(10):671–672.
36. Fischer AA. Pressure algometry over normal muscles. Standard values, validity and reproducibility of pressure threshold. *Pain*. 1987;30(1):115–126.
37. Quinn BL, Sheldon LK, Cooley ME. Pediatric pain assessment by drawn faces scales: a review. *Pain Manag Nurs*. 2014;15(4):909–918.
38. Zhou H, Albrecht MA, Roberts PA, et al. Consistency of pediatric pain ratings between dyads: an updated meta-analysis and metaregression. *Pain Rep*. 2022;7(5):e1029.
39. Press J, Gidron Y, Maimon M, et al. Effects of active distraction on pain of children undergoing venipuncture: who benefits from it?. *Pain Clinic*. 2003;15(3):261–269.