


Results: The severity of the injuries after shooting were assessed using the Abbreviated Injury Scale and did not differ significantly between the animals. All survived the first 60 min. The production of ROS significantly was lower in the group that had received hydrocortisone (see figure). Administration of U0126 reduced the synthesis of ROS by about two-thirds in both groups, independently of time.

Conclusion: Early treatment with hydrocortisone after trauma inhibits the synthesis of ROS. The results also suggest that the ERK branch of the MAP-kinase signalling cascade is an important transduction pathway in trauma. The capacity of selective ERK inhibitors to modulate the post-traumatic inflammatory response is promising, but compelling evidence of their usefulness in vivo still is lacking.

References: Duncia JV, et al: MEK inhibitors: The chemistry and biological activity of U0126, its analogs, and cyclization products. *Bioorg Med Chem Lett* 1998;8:2839–2844.

Keywords: glucocorticoids; gunshot; MAPK; reactive O₂ species; trauma

Prehosp Disast Med 2002;17:s3.



Impaired Recognition of Novelty in Rats with Mild Hypothermia Observed in Body or Brain

Myhrer T, Try Valo E, Haugen AH

Norwegian Defense Research Establishment, Division for Protection and Materiel, Kjeller, Norway

Introduction: Optimal cognitive function is dependent on normal body/brain temperature in both humans and animals. Relatively low temperatures have been linked to cognitive deficits in humans (33°C) and in rats (30°C). A link between body and brain temperatures has been reported for the rat. The purpose of the present study was two-fold. First, it was to examine whether the two categories of mild hypothermia (32–34°C and 35–37°C) might result in cognitive impairment in a novelty test previously shown to be sensitive to subtle brain lesions. Second, it was investigated whether recording body temperature and brain temperature might yield parallel results.

Materials and Methods: Male Wistar rats were provided with a transmitter in the peritoneum for monitoring body temperature or a thermistor probe in the frontal cortex for monitoring brain temperature. The rats were cooled in water (14–16°C) for 6–8 min., and then were allowed to dry for 30 min before behavioral testing. Following a pre-training procedure, the rats were twice given a novelty test in which optimal performance requires the detection of a discrete novel object.

Results: It was found that both levels of hypothermia resulted in a decreased ability to detect environmental changes; the deepest hypothermia (32–34°C) produced the most pronounced effects. A very close correspondence was noted between body and brain temperatures and behavioral effects. In

normothermic control animals, the body and brain temperatures virtually were identical in the standardized test situation.

Conclusion: It is evident that a modest decline in body/brain temperature of 1–3°C is sufficient to impair cognitive functions.

Keywords: brain; cognition; hypothermia; temperature
Prehosp Disast Med 2002;17:s 4.

Reproductive Health Characteristics in the Danish Gulf War Study

Guldager B

Danish Armed Forces Health Services

Introduction: An increased prevalence of neuropsychological, gastrointestinal, and skin symptoms^{1–7} has been found in Danish Gulf War Veterans (GWV). The hypothesis was tested that the male reproductive system also could have been affected by various exposures.

Methods: A cross-sectional study was performed during the period of 1997 to 1998, involving 661 male subjects deployed in the Gulf between 02 August 1990 and 31 December 1997. A control group of 215 Danish military men was selected with random matching by age and type of work. All participants were interviewed based on a previously completed comprehensive questionnaire, and underwent clinical and para-clinical examinations. A venous blood sample was drawn to determine serum concentrations of the follicle-stimulating hormone, the luteinizing hormone, testosterone, serum hormone binding globulin, and inhibin B.

Results: No differences were found between GWV and controls with respect to any of the reproductive hormones, nor with respect to fertility or the prevalence of spontaneous abortions, congenital diseases, or malformations among the offspring. Also, cohabitational characteristics were similar. These results are in agreement with later unpublished data from an explorative, tentative study looking at specific diagnoses in children of GWV and in those of controls. Only self-assessed sexual problems were more prevalent among GWV, 12%, as compared with 3.7% among controls, $p < 0.001$. A statistically significant association was found for self-reported sexual problems and psychological exposures related to specific traumatic events during deployment.

Conclusion: The biological reproductive health of male GWV seemed to be unaffected by their engagement in the post-war peacekeeping mission.⁸

References

1. *Dan Med Bull* 1999;46(5):416–419.
2. *Ugeskr Laeger* 1999;161(39):5423–5428.
3. *Dan Med Bull* 1999;46(5):420–423.
4. *Dan Med Bull* 1999;46(5):423–427.
5. *Ugeskr Laeger* 2000;162(49):6687–6691.
6. *APMIS* 2001; 109(7–8):517–524.
7. *Dan Med Bull* 2001;48(1):29–32.
8. *Scand J Sex* 2001;1(4):43–55.

Keywords: health, reproductive; Gulf War; peacekeeping; resources

Prehosp Disast Med 2002;17:s4.

Exercise-Induced Bronchial Obstruction and Weather

Latvala J,¹ Karjalainen J,² Remes J¹

1. Finnish Institute of Occupational Health, Oulu, Finland
2. Central Military Hospital, Helsinki, Finland

Introduction: Exercise-induced asthma (EIA) is a common respiratory disorder. The mechanism of airway narrowing is not simple, but airway drying and cooling evidently play a major role. The aim of this study was to investigate how closely the airway narrowing observed in EIA depends on ambient temperature and humidity, and whether the dependence on humidity is greater than on temperature.

Material and Methods: The population consisted of 1,809 army conscripts admitted to the Central Military Hospital in Helsinki, in order to assess their fitness for military service or to diagnose asthma. The maximum decrease in peak expiratory flow (DPEF%) was measured 30 minutes after an eight-minute outdoor running test, with the heart rate at 85% of the predicted maximum. The dependence of DPEF% on temperature and humidity was analysed by a linear regression model. Statistical significance was tested using the t-test.

Results: The dependence of DPEF% on temperature and humidity was negative and significant statistically ($p < 0.001$). The regression coefficient with respect to absolute air humidity was -1.026 and with respect to air temperature was -0.327 . The difference was significant statistically.

- DPEF% = $23.907 - 1.026 \times$ absolute air humidity, $p < 0.001$.

- DPEF% = $20.347 - 0.327 \times$ air temperature, $p < 0.001$.

The DPEF% was dependent most clearly on temperature and humidity when the ambient temperature was below -4°C and humidity below $4\text{gH}_2\text{O/kg}$.

Conclusion: Air temperature, and more obviously air humidity, have a significant influence on airway narrowing in EIA, and should be taken into account when EIA tests are performed on runners in the outdoors.

Keywords: asthma, exercise-induced; humidity; peak expiratory flow rate; temperature; weather

Prehosp Disast Med 2002;17:s5.

Cold-Allodynia and Alterations in Thin- and Thick-Fibre Function after Cold Injuries in Military Coastal Rangers

Jorum E, Opstad PK

Laboratory of Neurophysiology, Department of Neurology, National Hospital, Oslo; Norwegian Defense Research Establishment, Division for Material and Protection, Kjeller, Norway

Introduction: Cold allodynia and hyperalgesia are typical findings in patients with neuropathic pain of different origins. The neurophysiological mechanisms are unknown. Cold allodynia and hyperalgesia also are observed as sequelae in subjects who previously have been exposed to cold injuries.

Material and Methods: The subjects investigated for this study were 26 coastal rangers and officers who participated in a 5-day military training course, in small, open vessels along the coast of Helgeland in January 1998. All soldiers were exposed to the same type of cold: wet and cloudy the first 2 days, followed by cold and wind the remaining days. The first investigation of the soldiers was performed 2–3 months after the cold injury. All soldiers had a thorough clinical investigation, including a neurography, an investigation of thin fiber function by Thermotest (Somedic), a measurement of subcutaneous fat, and maximal oxygen uptake.

Results: Six subjects were without symptoms and had normal neurological and neurophysiological findings. The other 20 had different types of sensory disturbances. Compared to a normal control group from the Norwegian Military Academy,

the rest of the subjects had neurophysiological findings of both thick and thin fiber neuropathy. There was no significant alteration in the heat pain threshold. The most significant and interesting finding was the reduction in the cold pain threshold in the feet (cold allodynia). A one and two year follow-up has demonstrated that the thin and thick fiber neuropathy healed completely, whereas cold allodynia still persisted in some of the subjects. Ischaemic blockade of the nerve fibers indicates that the cold allodynia is mediated through the Ad-fibers.

Conclusion: Soldiers with persistent symptoms after 2 years experienced pain when exposed to cold at higher temperature than normal. Cold allodynia, as demonstrated by the Thermotest, is the neurophysiological finding of relevance to this symptom. Therefore, assessment of thin-fiber function is of great value in evaluating patients with sequelae following cold injuries.

Definition: Allodynia is the pain response to stimuli that normally are not painful.

Keywords: allodynia; cold; hyperalgesia; injuries; neuropathy; sequelae; thermotest

Prehosp Disast Med 2002;17:s5.

Stress-Related Anterior Lower Leg Pain: Evaluation with MRI and Intracompartmental Pressure Measurement

Kiuru MJ,^{1,2} Mäntysaari MJ,¹ Pibljämäki HK,³ Ahovuo JA²

1. Research Institute of Military Medicine, Helsinki, Finland
2. Department of Radiology, Helsinki University Central Hospital, Helsinki, Finland
3. Department of Surgery, Central Military Hospital, Helsinki, Finland

Purpose: To assess the feasibility of MRI and anterior tibial compartment pressure measurement in evaluating the causes of stress-related anterior lower leg pain.

Materials and Methods: This series was composed of 24 consecutive conscripts suffering from stress-related, anterior, lower leg pain. Twenty patients had bilateral symptoms. The patients underwent clinical examination, MR imaging, and intracompartmental pressure measurement. The intracompartmental pressure was measured in the anterior tibial compartment using a standardized protocol. Intra-compartmental pressure values higher than 40 mmHg were considered pathological.

Results: As compared with the intracompartmental pressure measurement, 50% of the patients had a false-positive clinical diagnosis of chronic exertional compartment syndrome (CECS). The duration of symptoms was longer in the patients with CECS than in those without ($p < 0.01$). MR imaging revealed abnormal findings in 35 symptomatic legs, the remaining 9 symptomatic legs were normal. In 33 legs, there was a stress fracture, and two legs had traction periostitis. There were no significant differences in the bone findings between the patients with CECS and those without ($p > 0.05$).

Conclusions: Stress-related anterior lower leg pain may result from CECS, stress fracture, and traction periostitis. The clinical diagnosis is not reliable. CECS and stress fracture or traction periostitis may occur separately or simultaneously.

Keywords: compartmental pressures; leg; MRI; pain; periostitis; stress fracture

Prehosp Disast Med 2002;17:s6.