

DERIVATIVES OF THETA FUNCTIONS AS TRACES OF PARTITION EISENSTEIN SERIES

TEWODROS AMDEBERHAN , KEN ONO  AND AJIT SINGH 

Abstract. In his “lost notebook,” Ramanujan used iterated derivatives of two theta functions to define sequences of q -series $\{U_{2t}(q)\}$ and $\{V_{2t}(q)\}$ that he claimed to be quasimodular. We give the first explicit proof of this claim by expressing them in terms of “partition Eisenstein series,” extensions of the classical Eisenstein series $E_{2k}(q)$, defined by

$$\lambda = (1^{m_1}, 2^{m_2}, \dots, n^{m_n}) \vdash n \quad \mapsto \quad E_\lambda(q) := E_2(q)^{m_1} E_4(q)^{m_2} \cdots E_{2n}(q)^{m_n}.$$

For functions $\phi : \mathcal{P} \mapsto \mathbb{C}$ on partitions, the *weight $2n$ partition Eisenstein trace* is

$$\mathrm{Tr}_n(\phi; q) := \sum_{\lambda \vdash n} \phi(\lambda) E_\lambda(q).$$

For all t , we prove that $U_{2t}(q) = \mathrm{Tr}_t(\phi_U; q)$ and $V_{2t}(q) = \mathrm{Tr}_t(\phi_V; q)$, where ϕ_U and ϕ_V are natural partition weights, giving the first explicit quasimodular formulas for these series.

§1. Introduction and statement of results

In his “lost notebook”, Ramanujan considered the [12, p. 369] two sequences of q -series:

$$U_{2t}(q) = \frac{1^{2t+1} - 3^{2t+1}q + 5^{2t+1}q^3 - 7^{2t+1}q^6 + \cdots}{1 - 3q + 5q^3 - 7q^6 + \cdots} = \frac{\sum_{n \geq 0} (-1)^n (2n+1)^{2t+1} q^{\frac{n(n+1)}{2}}}{\sum_{n \geq 0} (-1)^n (2n+1) q^{\frac{n(n+1)}{2}}}, \quad (1.1)$$

$$V_{2t}(q) = \frac{1^{2t} - 5^{2t}q - 7^{2t}q^2 + 11^{2t}q^5 + \cdots}{1 - q - q^2 + q^5 + \cdots} = \frac{\sum_{n=-\infty}^{\infty} (-1)^n (6n+1)^{2t} q^{\frac{n(3n+1)}{2}}}{\sum_{n=-\infty}^{\infty} (-1)^n q^{\frac{n(3n+1)}{2}}}, \quad (1.2)$$

and he offered identities such as

$$\begin{aligned} U_0 &= 1, & U_2 &= E_2, & U_4 &= \frac{1}{3}(5E_2^2 - 2E_4), & U_6 &= \frac{1}{9}(35E_2^3 - 42E_2E_4 + 16E_6), \dots \\ V_0 &= 1, & V_2 &= E_2, & V_4 &= 3E_2^2 - 2E_4, & V_6 &= 15E_2^3 - 30E_2E_4 + 16E_6, \dots \end{aligned}$$

where $E_2(q), E_4(q)$, and $E_6(q)$ are the usual Eisenstein series

$$E_2 := 1 - 24 \sum_{n=1}^{\infty} \sigma_1(n) q^n, \quad E_4 := 1 + 240 \sum_{n=1}^{\infty} \sigma_3(n) q^n, \quad \text{and} \quad E_6 := 1 - 504 \sum_{n=1}^{\infty} \sigma_5(n) q^n,$$

and where $\sigma_v(n) := \sum_{d|n} d^v$. He made the following claim:

“In general U_{2t} and V_{2t} are of the form $\sum K_{\ell, m, n} E_2^\ell E_4^m E_6^n$, where $\ell + 2m + 3n = t$.”

Received July 11, 2024. Revised August 6, 2024. Accepted October 25, 2024.

2020 Mathematics subject classification: Primary 11F03, 05A17; Secondary 11M36.

Keywords: partition Eisenstein series, quasimodular forms.

© The Author(s), 2025. Published by Cambridge University Press on behalf of Foundation Nagoya Mathematical Journal. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted re-use, distribution, and reproduction in any medium, provided the original work is properly cited.



Berndt, Chan, Liu, Yee, and Yesilyurt [5], [6] proved this claim using Ramanujan's identities [13]

$$D(E_2) = \frac{E_2^2 - E_4}{12}, \quad D(E_4) = \frac{E_2 E_4 - E_6}{3}, \quad \text{and} \quad D(E_6) = \frac{E_2 E_6 - E_4^2}{2}, \quad (1.3)$$

where $D := q \frac{d}{dq}$. However, their results are not explicit. Indeed, Andrews and Berndt (see p. 364 of [3]) proclaim that "...it seems extremely difficult to find a general formula for all $K_{\ell, m, n}$."

We offer a solution to the general problem of obtaining the first explicit formulas for U_{2t} and V_{2t} . We note that Ramanujan's claim is that U_{2t} and V_{2t} are weight $2t$ quasimodular forms, as the ring of quasimodular forms is the polynomial ring (for example, see [10])

$$\mathbb{C}[E_2, E_4, E_6] = \mathbb{C}[E_2, E_4, E_6, E_8, E_{10}, \dots],$$

and so our goal is to obtain explicit formulas in terms of the classical sequence of Eisenstein series (for example, see Chapter 1 of [11] or Thm. 12.17 of [4])

$$E_{2k}(q) := 1 - \frac{4k}{B_{2k}} \sum_{n=1}^{\infty} \sigma_{2k-1}(n) q^n, \quad (1.4)$$

where B_{2k} is the $2k$ th Bernoulli number. We express Ramanujan's q -series as explicit "traces of partition Eisenstein series."

As an important step toward this goal, we first derive generating functions for his series. In terms of Dedekind's eta-function $\eta(q) := q^{\frac{1}{24}} \prod_{n=1}^{\infty} (1 - q^n)$ and Jacobi–Kronecker quadratic characters, we have the following result.

THEOREM 1.1. *As a power series in X , the following are true.*

(1) *If $\chi_{-4}(\cdot) = \left(\frac{-4}{\cdot}\right)$, then we have*

$$\sum_{t \geq 0} (-1)^t U_{2t}(q) \cdot \frac{X^{2t+1}}{(2t+1)!} = \frac{1}{2\eta(q)^3} \cdot \sum_{n \in \mathbb{Z}} \chi_{-4}(n) q^{\frac{n^2}{8}} \sin(nX).$$

(2) *If $\chi_{12}(\cdot) = \left(\frac{12}{\cdot}\right)$, then we have*

$$\sum_{t \geq 0} (-1)^t V_{2t}(q) \cdot \frac{X^{2t}}{(2t)!} = \frac{1}{2\eta(q)} \cdot \sum_{n \in \mathbb{Z}} \chi_{12}(n) q^{\frac{n^2}{24}} \cos(nX).$$

REMARK 1.2. Theorem 1.1 represents two special cases of Lemma 2.1, which pertains to arbitrary theta functions. Using Theorem 1.1, we obtain Theorem 3.4 that gives two further identities for these particular generating functions as infinite products in trigonometric functions.

These generating functions shall offer the connection to traces of partition Eisenstein series. To make this precise, we recall that a *partition of n* is any nonincreasing sequence of positive integers $\lambda = (\lambda_1, \lambda_2, \dots, \lambda_s)$ that sum to n , denoted $\lambda \vdash n$. Equivalently, we use the notation $\lambda = (1^{m_1}, \dots, n^{m_n}) \vdash n$, where m_j is the multiplicity of j . For such λ , we define

the weight $2n$ partition Eisenstein series¹

$$\lambda = (1^{m_1}, 2^{m_2}, \dots, n^{m_n}) \vdash n \quad \longmapsto \quad E_\lambda(q) := E_2(q)^{m_1} E_4(q)^{m_2} \dots E_{2n}(q)^{m_n}. \quad (1.5)$$

The Eisenstein series $E_{2k}(q)$ corresponds to the partition $\lambda = (k)$, as we have $E_{(k^1)}(q) = E_{2k}(q)$ ¹.

To define partition traces, suppose that $\phi : \mathcal{P} \mapsto \mathbb{C}$ is a function on partitions. For each positive integer n , its partition Eisenstein trace is the weight $2n$ quasimodular form

$$\text{Tr}_n(\phi; q) := \sum_{\lambda \vdash n} \phi(\lambda) E_\lambda(q). \quad (1.6)$$

Such traces arise in recent work on MacMahon’s sums-of-divisors q -series (see Thm. 1.4 of [1]).

For partitions $\lambda = (1^{m_1}, \dots, n^{m_n}) \vdash n$, we require the following functions:

$$\phi_U(\lambda) := 4^n (2n + 1)! \cdot \prod_{k=1}^n \frac{1}{m_k!} \left(\frac{B_{2k}}{(2k)(2k)!} \right)^{m_k}, \quad (1.7)$$

$$\phi_V(\lambda) := 4^n (2n)! \cdot \prod_{k=1}^n \frac{1}{m_k!} \left(\frac{(4^k - 1)B_{2k}}{(2k)(2k)!} \right)^{m_k}. \quad (1.8)$$

Ramanujan’s series are weighted traces of partition Eisenstein series of these functions.

THEOREM 1.3. *If t is a positive integer, then the following are true.*

- (1) *We have that $U_{2t}(q) = \text{Tr}_t(\phi_U; q)$.*
- (2) *We have that $V_{2t}(q) = \text{Tr}_t(\phi_V; q)$.*

EXAMPLES. Here we offer examples of Theorem 1.3.

- (1) By direct calculation, we find for $t = 3$ that

$$\phi_U((3^1)) = 16/9, \quad \phi_U((1^1, 2^1)) = -42/9, \quad \text{and} \quad \phi_U((1^3)) = 35/9.$$

This reproduces Ramanujan’s identity

$$\text{Tr}_3(\phi_U; q) = \frac{1}{9}(16E_6 - 42E_2E_4 + 35E_2^3) = U_6.$$

- (2) By direct calculation, we find for $t = 4$ that

$$\begin{aligned} \phi_V((4^1)) &= -272, \phi_V((1^1, 3^1)) = 448, \phi_V((2^2)) = 140, \phi_V((1^2, 2^1)) \\ &= -420, \text{ and } \phi_V((1^4)) = 105. \end{aligned}$$

Therefore, we have that

$$\text{Tr}_4(\phi_V; q) = -272E_8 + 448E_2E_6 + 140E_4^2 - 420E_2^2E_4 + 105E_2^4 = V_8.$$

- (3) Similar calculations give expressions for the weight 14 quasimodular forms U_{14} and V_{14}

$$\begin{aligned} U_{14} &= \frac{1}{405}(25025E_2^7 - 210210E_2^5E_4 + 420420E_2^3E_4^2 - 168168E_2E_4^3 + 400400E_2^4E_6 \\ &\quad - 960960E_2^2E_4E_6 + 192192E_4^2E_6 + 366080E_2E_6^2 - 720720E_2^3E_8 + 864864E_2E_4E_8 \end{aligned}$$

¹ These E_λ should not be confused with the partition Eisenstein series introduced by Just and Schneider [9], which are semi-modular instead of quasimodular.

$$\begin{aligned}
 & -329472E_6E_8 + 1048320E_2^2E_{10} - 419328E_4E_{10} - 1061376E_2E_{12} + 552960E_{14}), \\
 V_{14} = & 135135E_2^7 - 1891890E_2^5E_4 + 6306300E_2^3E_4^2 - 4204200E_2E_4^3 + 5045040E_2^4E_6 \\
 & - 20180160E_2^2E_4E_6 + 6726720E_4^2E_6 + 10762752E_2E_6^2 - 12252240E_2^3E_8 \\
 & + 24504480E_2E_4E_8 - 13069056E_6E_8 + 23831808E_2^2E_{10} - 15887872E_4E_{10} \\
 & - 32195072E_2E_{12} + 22368256E_{14}.
 \end{aligned}$$

Then, using the identities

$$E_{14} = E_2^4E_6, \quad E_{12} = \frac{250}{691}E_6^2 + \frac{441}{691}E_4^3, \quad E_{10} = E_4E_6, \quad E_8 = E_4^2,$$

we get

$$\begin{aligned}
 U_{14} = & \frac{1}{27}(-3648E_4^2E_6 - 17920E_2E_6^2 + 19320E_2E_4^3 - 300300E_2^3E_4^2 \\
 & + 400400E_2^4E_6 - 210210E_2^5E_4 + 87360E_2^2E_4E_6 + 25025E_2^7), \\
 V_{14} = & 138048E_4^2E_6 - 885248E_2E_6^2 - 246792E_2E_4^3 - 5945940E_2^3E_4^2 \\
 & + 5045040E_2^4E_6 - 1891890E_2^5E_4 + 3651648E_2^2E_4E_6 + 135135E_2^7.
 \end{aligned}$$

REMARK 1.4. The coefficients $\phi_V(\lambda)$ are always integers. The first author and other collaborators have found a combinatorial interpretation and a proof this fact, which will appear in a forthcoming work.

In view of these results, it is natural to pose the following problem.

PROBLEM. Determine and characterize further functions $\phi : \mathcal{P} \mapsto \mathbb{C}$ for which $\{\text{Tr}_t(\phi; q)\}$ is a natural and rich family of weight $2t$ quasimodular forms.

To prove these results, we make use of the Jacobi Triple Product identity, special q -series, exponential generating functions for Bernoulli numbers, and properties of Pólya’s cycle index polynomials. In Section 2, we derive a general result for q -series of the form (1.1) and (1.2) (see Lemma 2.1), which gives Theorem 1.1 as special cases. In Section 3, we prove Theorem 1.3 using these results and properties of Pólya’s cycle index polynomials.

§2. Generating functions for Ramanujan-type q -series

Theorem 1.1 gives two special cases of general generating functions associated to formal theta functions for Dirichlet characters. If χ modulo N is a Dirichlet character, then let

$$\Theta(\chi; q) := \sum_{n=1}^{\infty} \chi(n)n^{a_\chi}q^{n^2}, \tag{2.1}$$

where we let

$$a_\chi := \begin{cases} 0 & \text{if } \chi \text{ is even,} \\ 1 & \text{if } \chi \text{ is odd.} \end{cases} \tag{2.2}$$

Then, in analogy with Ramanujan’s U_{2t} and V_{2t} (see (1.1) and (1.2)), we let

$$R_{2t}(\chi; q) := \frac{D^t(\Theta(\chi; q))}{\Theta(\chi; q)} = \frac{\sum_{n=1}^{\infty} \chi(n)n^{2t+a_\chi}q^{n^2}}{\Theta(\chi; q)}. \tag{2.3}$$

LEMMA 2.1. Assuming the notation above, as a power series in X we have

$$\sum_{t=0}^{\infty} (-1)^t R_{2t}(\chi; q) \cdot \frac{X^{2t+1}}{(2t+1)!} = \frac{1}{2i\Theta(\chi; q)} \sum_{n \in \mathbb{Z}} \chi(n) q^{n^2} n^{a_\chi-1} \cdot e^{inX}.$$

REMARK 2.2. Lemma 2.1 holds for periodic functions $\chi : \mathbb{Z} \rightarrow \mathbb{C}$ that are either even or odd.

Proof. By direct calculation, we have that

$$\frac{1}{2i} \sum_{n \in \mathbb{Z}} \chi(n) q^{n^2} n^{a_\chi-1} \cdot e^{inX} = \frac{1}{2i} \sum_{n=1}^{\infty} q^{n^2} n^{a_\chi-1} (\chi(n) e^{inX} + (-1)^{a_\chi-1} \chi(-n) e^{-inX}).$$

For all χ , we have that $(-1)^{a_\chi-1} \chi(-n) = -\chi(n)$, and so this reduces to

$$\frac{1}{2i} \sum_{n=1}^{\infty} \chi(n) q^{n^2} n^{a_\chi-1} (e^{inX} - e^{-inX}) = \sum_{n=1}^{\infty} \chi(n) q^{n^2} n^{a_\chi-1} \sin(nX).$$

Using the Taylor series for $\sin(nX)$, this gives (after change of summation)

$$\begin{aligned} \frac{1}{2i} \sum_{n \in \mathbb{Z}} \chi(n) q^{n^2} n^{a_\chi-1} \cdot e^{inX} &= \sum_{n=1}^{\infty} \chi(n) n^{a_\chi-1} q^{n^2} \sum_{t=0}^{\infty} (-1)^t \cdot \frac{(nX)^{2t+1}}{(2t+1)!} \\ &= \sum_{t=0}^{\infty} (-1)^t \cdot \frac{X^{2t+1}}{(2t+1)!} \cdot D^t(\Theta(\chi; q)). \end{aligned}$$

Thanks to (2.3), we obtain the claimed generating function by dividing through by $\Theta(\chi; q)$. □

Proof of Theorem 1.1. To prove claim (1), we consider $\chi_{-4}(n) := (\frac{-4}{n})$, which is the only odd character modulo 4. In this case, we have $a_{\chi_{-4}} = 1$, and so Lemma 2.1 gives

$$\sum_{t=0}^{\infty} (-1)^t R_{2t}(\chi_{-4}; q) \cdot \frac{X^{2t+1}}{(2t+1)!} = \frac{1}{2i\Theta(\chi_{-4}; q)} \sum_{n \in \mathbb{Z}} \chi_{-4}(n) q^{n^2} \cdot e^{inX}.$$

Furthermore, Jacobi’s classical identity (for example, see p. 17 of [11]) implies that

$$\eta(q^8)^3 = \Theta(\chi_{-4}; q) = \sum_{n=0}^{\infty} (-1)^n (2n+1) q^{(2n+1)^2}.$$

Therefore, we have that

$$\sum_{t=0}^{\infty} (-1)^t R_{2t}(\chi_{-4}; q) \cdot \frac{X^{2t+1}}{(2t+1)!} = \frac{1}{2i\eta(q^8)^3} \sum_{n \in \mathbb{Z}} \chi_{-4}(n) q^{n^2} \cdot e^{inX}.$$

Claim (1) follows by letting $q \rightarrow q^{\frac{1}{8}}$, replacing the complex exponential in terms of trigonometric functions, followed by taking the real part.

To prove claim (2), we note that Euler’s Pentagonal Number Theorem (see p. 17 of [11]) implies that

$$\eta(q^{24}) = \sum_{n=1}^{\infty} \chi_{12}(n) q^{n^2},$$

where $\chi_{12}(n) = \left(\frac{12}{n}\right)$ is the unique primitive character with conductor 12. Therefore, $a_\chi = 0$, and so Lemma 2.1 gives

$$\sum_{t=0}^{\infty} (-1)^t R_{2t}(\chi_{12}; q) \cdot \frac{X^{2t+1}}{(2t+1)!} = \frac{1}{2i\eta(q^{24})} \sum_{n \in \mathbb{Z}} \chi_{12}(n) \frac{q^{n^2}}{n} \cdot e^{inX}.$$

Claim (2) follows by letting $q \rightarrow q^{\frac{1}{24}}$, then differentiating in X , followed by taking the real part as in (1). □

§3. Proof of Theorem 1.3

Here we prove Theorem 1.3 using the generating functions in Theorem 1.1. We apply Pólya’s cycle index polynomials and the exponential generating function for Bernoulli numbers.

3.1 Bernoulli numbers

Here we recall a convenient generating function for Bernoulli numbers, and we refer the reader to [8, 1.518.1]. If $\text{sinc}(X) := \sin X/X$, then we have

$$\frac{1}{\text{sinc}(X)} = \exp \left(- \sum_{k \geq 1} \frac{(-4)^k B_{2k}}{(2k)(2k)!} \cdot X^{2k} \right). \tag{3.1}$$

3.2 Pólya’s cycle index polynomials

We require Pólya’s cycle index polynomials in the case of symmetric groups (for example, see [14]). Namely, recall that given a partition $\lambda = (\lambda_1, \dots, \lambda_{\ell(\lambda)}) \vdash t$ or $(1^{m_1}, \dots, t^{m_t}) \vdash t$, we have that the number of permutations in \mathfrak{S}_t of cycle type λ is $t!/z_\lambda$, where $z_\lambda := 1^{m_1} \dots t^{m_t} m_1! \dots m_t!$. The *cycle index polynomial* for the symmetric group \mathfrak{S}_t is given by

$$Z(\mathfrak{S}_t) = \sum_{\lambda \vdash t} \frac{1}{z_\lambda} \prod_{i=1}^{\ell(\lambda)} x_{\lambda_i} = \sum_{\lambda \vdash t} \prod_{k=1}^t \frac{1}{m_k!} \left(\frac{x_k}{k}\right)^{m_k}. \tag{3.2}$$

We require the following well-known generating function in t -aspect.

LEMMA 3.1. (Example 5.2.10 of [14]) *As a power series in y , the generating function for the cycle index polynomials satisfies*

$$\sum_{t \geq 0} Z(\mathfrak{S}_t) y^t = \exp \left(\sum_{k \geq 1} x_k \frac{y^k}{k} \right).$$

EXAMPLE 3.2. Here are the first few examples of Pólya’s cycle index polynomials:

$$Z(\mathfrak{S}_1) = x_1, Z(\mathfrak{S}_2) = \frac{1}{2!}(x_1^2 + x_2), Z(\mathfrak{S}_3) = \frac{1}{3!}(x_1^3 + 3x_1x_2 + 2x_3).$$

3.3 Some power series identities

We consider the infinite series factors of the functions in Theorem 1.1.

LEMMA 3.3. *As a power series in X , the following are true.*

(1) *We have that*

$$\frac{q^{-\frac{1}{8}}}{2} \cdot \sum_{n \in \mathbb{Z}} \chi_{-4}(n) \cdot q^{\frac{n^2}{8}} \sin(nX) = \sin X \cdot \prod_{j \geq 1} (1 - q^j)(1 - 2 \cos(2X)q^j + q^{2j}).$$

(2) We have that

$$\begin{aligned} & \frac{q^{-\frac{1}{24}}}{2} \sum_{n \in \mathbb{Z}} \chi_{12}(n) \cdot q^{\frac{n^2}{24}} \cos(nX) \\ &= \cos X \prod_{n \geq 1} (1 - q^n)(1 + 2 \cos(2X)q^n + q^{2n})(1 - 2 \cos(4X)q^{2n-1} + q^{4n-2}). \end{aligned}$$

Proof. Both claims follow from the Jacobi Triple Product Identity (see Th. 2.8 of [2])

$$\sum_{n \in \mathbb{Z}} (-1)^n q^{\frac{n^2}{2}} z^n = \prod_{j \geq 1} (1 - q^j)(1 - q^{j-\frac{1}{2}}z)(1 - q^{j-\frac{1}{2}}z^{-1}). \tag{3.3}$$

To prove (1), we make the substitutions $2i \sin X = e^{iX}(1 - e^{-2iX})$ and $z = q^{\frac{1}{2}}e^{2iX}$ to obtain

$$\frac{1}{2i} \sum_{n \in \mathbb{Z}} (-1)^n q^{\frac{n^2+n}{2}} e^{(2n+1)iX} = \sin X \prod_{j \geq 1} (1 - q^j)(1 - q^j e^{2iX})(1 - q^j e^{-2iX}).$$

To obtain claim (1), we note the following simple reformulation

$$\frac{1}{2i} \sum_{n \in \mathbb{Z}} (-1)^n q^{\frac{n^2+n}{2}} e^{(2n+1)iX} = \frac{q^{-\frac{1}{8}}}{2i} \cdot \sum_{n \in \mathbb{Z}} \chi_{-4}(n) \cdot q^{\frac{n^2}{8}} e^{inX},$$

and then take the real part of both sides.

We now turn to claim (2). Thanks to [7, (1.1)], we obtain

$$\begin{aligned} \sum_{n \in \mathbb{Z}} q^{n(3n+1)}(z^{3n} - z^{-3n-1}) &= \prod_{n \geq 1} (1 - q^{2n})(1 - zq^{2n}) \left(1 - \frac{q^{2n-2}}{z}\right) \\ &\quad \times (1 - z^2q^{4n-2}) \left(1 - \frac{q^{4n-2}}{z^2}\right). \end{aligned}$$

By replacing $q \rightarrow q^{\frac{1}{2}}, z \rightarrow -z^2$, factoring out $1 + z^{-2}$ and multiplying through by z , we get

$$\begin{aligned} \sum_{n \in \mathbb{Z}} (-1)^n q^{\frac{n(3n+1)}{2}} \frac{z^{6n+1} + z^{-6n-1}}{z + z^{-1}} &= \prod_{n \geq 1} (1 - q^n)(1 + z^2q^n) \left(1 + \frac{q^n}{z^2}\right) \\ &\quad \times (1 - z^4q^{2n-1}) \left(1 - \frac{q^{2n-1}}{z^4}\right). \end{aligned}$$

After letting $z = e^{iX}$, we pair up conjugate terms to get

$$\begin{aligned} & \sum_{n \in \mathbb{Z}} (-1)^n q^{\frac{n(3n+1)}{2}} \cos(6n + 1)X \\ &= \cos X \prod_{n \geq 1} (1 - q^n)(1 + 2 \cos(2X)q^n + q^{2n})(1 - 2 \cos(4X)q^{2n-1} + q^{4n-2}). \end{aligned}$$

The left hand side of the expression above equals the infinite sum in Lemma 3.3 (2). □

To prove Theorem 1.3, we also require the following power series identities.

THEOREM 3.4. *The following identities are true.*

(1) *As power series in X , we have*

$$\sum_{t \geq 0} (-1)^t U_{2t}(q) \cdot \frac{X^{2t+1}}{(2t+1)!} = \sin X \cdot \prod_{j \geq 1} \left[1 + \frac{4(\sin^2 X)q^j}{(1-q^j)^2} \right].$$

(2) *As power series in X , we have*

$$\sum_{t \geq 0} (-1)^t V_{2t}(q) \cdot \frac{X^{2t}}{(2t)!} = \cos X \cdot \prod_{j \geq 1} \left[1 - \frac{4(\sin^2 X)q^j}{(1+q^j)^2} \right] \left[1 + \frac{4(\sin^2 2X)q^{2j-1}}{(1-q^{2j-1})^2} \right].$$

Proof. We first prove claim (1). By combining Theorem 1.1 (1) and Lemma 3.3 (1), we obtain

$$\sum_{t \geq 0} (-1)^t U_{2t}(q) \cdot \frac{X^{2t+1}}{(2t+1)!} = \sin X \cdot \prod_{j \geq 1} \frac{(1 - 2 \cos(2X)q^j + q^{2j})}{(1 - q^j)^2}.$$

A straightforward algebraic manipulation with $-2 \cos(2X) = -2 + 4 \sin^2 X$ yields

$$\sum_{t \geq 0} (-1)^t U_{2t}(q) \cdot \frac{X^{2t+1}}{(2t+1)!} = \sin X \cdot \prod_{j \geq 1} \left[1 + \frac{4(\sin^2 X)q^j}{(1 - q^j)^2} \right].$$

Now we turn to claim (2). By combining Theorem 1.1 (2) and Lemma 3.3 (2), we obtain

$$\begin{aligned} \sum_{t \geq 0} (-1)^t V_{2t}(q) \cdot \frac{X^{2t}}{(2t)!} &= \cos X \prod_{n \geq 1} (1 + 2 \cos(2X)q^n + q^{2n})(1 - 2 \cos(4X)q^{2n-1} + q^{4n-2}) \\ &= \cos X \prod_{k \geq 1} (1 + q^k)^2 (1 - q^{2k-1})^2 \prod_{j \geq 1} \left[1 - \frac{4(\sin^2 X)q^j}{(1 + q^j)^2} \right] \\ &\quad \left[1 + \frac{4(\sin^2 2X)q^{2j-1}}{(1 - q^{2j-1})^2} \right]. \end{aligned}$$

The proof now follows from the simple identity

$$\prod_{k=1}^{\infty} (1 + q^k) = \prod_{k=1}^{\infty} \frac{1}{1 - q^{2k-1}}.$$

□

3.4 Proof of Theorem 1.3

For each positive odd integer j , we consider the Lambert series

$$\mathbf{S}_j(q) := \sum_{m \geq 1} \frac{m^j q^m}{1 - q^m} = \frac{B_{j+1}}{2(j+1)} - \frac{B_{j+1}}{2(j+1)} E_{j+1}(q). \tag{3.4}$$

This expression in terms of the $E_{j+1}(q)$ follows from (1.4). The proof of Theorem 1.3 boils down to deriving expressions for the power series in Theorem 3.4 in terms of the $\mathbf{S}_j(q)$.

Proof of Theorem 1.3. We first prove claim (1) regarding Ramanujan’s U_{2t} series. The key fact underlying the proof is the following power series identity.

$$\sum_{t \geq 0} (-1)^t U_{2t}(q) \cdot \frac{X^{2t+1}}{(2t+1)!} = \sin X \cdot \exp \left(-2 \sum_{r \geq 1} \frac{\mathbf{S}_{2r-1}(q)}{(2r)!} (-4X^2)^r \right). \tag{3.5}$$

Thanks to Theorem 3.4 (1), this identity will follow from

$$\exp \left(-2 \sum_{r \geq 1} \frac{\mathbf{S}_{2r-1}(q)}{(2r)!} (-4X^2)^r \right) = \prod_{j \geq 1} \left[1 + \frac{4(\sin^2 X)q^j}{(1-q^j)^2} \right]. \tag{3.6}$$

To establish (3.6), we compute the following double-sum in two different ways. First, we use the Taylor expansion of $\cos(y)$ and then interchange the order of summation to get

$$\sum_{j,k \geq 1} \frac{q^{kj} \cos(2kX)}{k} = \sum_{r \geq 0} \frac{(-4X^2)^r}{(2r)!} \sum_{k \geq 1} k^{2r-1} \sum_{j \geq 1} q^{kj}.$$

By combining the geometric series and the Taylor series for $\log(1 - Y)$ with (3.4), we obtain

$$\sum_{j,k \geq 1} \frac{q^{kj} \cos(2kX)}{k} = \sum_{r \geq 0} \frac{(-4X^2)^r}{(2r)!} \sum_{k \geq 1} \frac{k^{2r-1} q^k}{1 - q^k} = -\log(q)_\infty + \sum_{r \geq 1} \frac{\mathbf{S}_{2r-1}(q)}{(2r)!} (-4X^2)^r, \tag{3.7}$$

where $(q)_\infty := \prod_{n=1}^\infty (1 - q^n)$ is the q -Pochhammer symbol.

On the other hand, using $2 \cos \theta = e^{i\theta} + e^{-i\theta}$ and Taylor expansion of $\log(1 - Y)$, we find that

$$\begin{aligned} \sum_{j,k \geq 1} \frac{q^{kj} \cos(2kX)}{k} &= \frac{1}{2} \sum_{j \geq 1} \left(\sum_{k \geq 1} \frac{(e^{2iX} q^j)^k}{k} + \sum_{k \geq 1} \frac{(e^{-2iX} q^j)^k}{k} \right) \\ &= -\frac{1}{2} \sum_{j \geq 1} \log [1 - 2(\cos 2X)q^j + q^{2j}]. \end{aligned}$$

After straightforward algebraic manipulation, we get

$$\sum_{j,k \geq 1} \frac{q^{kj} \cos(2kX)}{k} = -\log(q)_\infty - \frac{1}{2} \log \left(\prod_{j \geq 1} \left[1 + \frac{4(\sin^2 X)q^j}{(1-q^j)^2} \right] \right). \tag{3.8}$$

Identity (3.6) follows by comparing (3.7) and (3.8), and in turn confirms (3.5).

We now investigate the exponential series in (3.5). Thanks to (3.4), followed by an application of (3.1), we obtain

$$\begin{aligned} &\exp \left(-2 \sum_{k \geq 1} \frac{(-4)^k \mathbf{S}_{2k-1}(q)}{(2k)!} X^{2k} \right) \\ &= \exp \left(- \sum_{k \geq 1} \frac{(-4)^k B_{2k}}{(2k)(2k)!} X^{2k} \right) \cdot \exp \left(\sum_{k \geq 1} \frac{B_{2k} \cdot E_{2k}(q)}{(2k)(2k)!} (-4X^2)^k \right) \end{aligned}$$

$$= \frac{X}{\sin X} \cdot \exp \left(\sum_{k \geq 1} \frac{B_{2k} \cdot E_{2k}(q)}{2(2k)!} \frac{(-4X^2)^k}{k} \right).$$

We recognize this last expression in the context of Pólya’s cycle index polynomials. Namely, Lemma 3.1 gives the identity (here $\lambda = (1^{m_1}, \dots, t^{m_t} \vdash t)$)

$$\exp \left(\sum_{k \geq 1} Y_k \frac{w^k}{k} \right) = \sum_{t \geq 0} \left(\sum_{\lambda \vdash t} \prod_{k=1}^t \frac{1}{m_k!} \left(\frac{Y_k}{k} \right)^{m_k} \right) w^t,$$

which we apply with $Y_k = \frac{B_{2k} \cdot E_{2k}(q)}{2(2k)!}$ and $w = -4X^2$. This gives

$$\begin{aligned} \sum_{t \geq 0} (-1)^t U_{2t}(q) \frac{X^{2t+1}}{(2t+1)!} &= \sin X \cdot \exp \left(-2 \sum_{k \geq 1} \frac{\mathbf{S}_{2k-1}(q)}{(2k)!} (-4X^2)^k \right) \\ &= \sin X \cdot \frac{X}{\sin X} \cdot \sum_{t \geq 0} \left(\sum_{\lambda \vdash t} \prod_{k=1}^t \frac{1}{m_k!} \left(\frac{B_{2k} \cdot E_{2k}(q)}{(2k)(2k)!} \right)^{m_k} \right) (-4X^2)^t. \end{aligned}$$

By comparing the coefficients of X^{2t+1} , we find that

$$(-1)^t \frac{U_{2t}(q)}{(2t+1)!} = (-4)^t \sum_{\lambda \vdash t} \prod_{k=1}^t \frac{1}{m_k!} \left(\frac{B_{2k} \cdot E_{2k}(q)}{(2k)(2k)!} \right)^{m_k},$$

which in turn, thanks to (1.7), proves Theorem 1.3 (1).

We now turn to claim (2) regarding Ramanujan’s V_{2t} whose proof is analogous to the proof of (1). The main difference follows from the need for the generalized Lambert series

$$\mathbf{A}_{2r-1}(q) := \sum_{k \geq 1} \frac{(-1)^{k-1} k^{2r-1} q^k}{1 - q^k} = \mathbf{S}_{2r-1}(q) - 4^r \mathbf{S}_{2r-1}(q^2). \tag{3.9}$$

The expression in \mathbf{S}_{2r-1} is straightforward. For the sake of brevity, we note that calculations analogous to the proof of (3.6) give the identity

$$\prod_{n \geq 1} \left[1 - \frac{4(\sin^2 X)q^n}{(1 + q^n)^2} \right] = \exp \left(2 \sum_{r \geq 1} \frac{\mathbf{A}_{2r-1}(q)(-4X^2)^r}{(2r)!} \right),$$

as well as

$$\begin{aligned} \prod_{n \geq 1} \left[1 + \frac{4(\sin^2 2X)q^{2n-1}}{(1 - q^{2n-1})^2} \right] &= \prod_{n \geq 1} \left[1 + \frac{4(\sin^2 2X)q^n}{(1 - q^n)^2} \right] \cdot \prod_{n \geq 1} \left[1 + \frac{4(\sin^2 2X)q^{2n}}{(1 - q^{2n})^2} \right]^{-1} \\ &= \exp \left(-2 \sum_{r \geq 1} \frac{\mathbf{S}_{2r-1}(q)(-16X^2)^r}{(2r)!} \right) \exp \left(2 \sum_{r \geq 1} \frac{\mathbf{S}_{2r-1}(q^2)(-16X^2)^r}{(2r)!} \right). \end{aligned}$$

Therefore, combining these two expressions with (3.9), Theorem 3.4 (2) gives

$$\sum_{t \geq 0} (-1)^t V_{2t}(q) \cdot \frac{X^{2t}}{(2t)!} = \cos X \cdot \exp \left(-2 \sum_{r \geq 1} \frac{\mathbf{S}_{2r-1}(q)}{(2r)!} (-4(2X)^2)^r \right) \exp \left(2 \sum_{r \geq 1} \frac{\mathbf{S}_{2r-1}(q)}{(2r)!} (-4X^2)^r \right).$$

As in the proof of (1), we recognize generating functions for Pólya’s cycle index polynomials. Namely, by applying (3.1) and Lemma 3.1 we obtain

$$\begin{aligned} \sum_{t \geq 0} \frac{(-1)^t V_{2t}(q) X^{2t}}{(2t)!} &= \cos X \cdot \frac{2X}{\sin(2X)} \cdot \exp \left(\sum_{k \geq 1} \frac{4^k B_{2k} \cdot E_{2k}(q)}{2(2k)!} \frac{(-4X^2)^k}{k} \right) \\ &\quad \times \frac{\sin X}{X} \cdot \exp \left(\sum_{k \geq 1} \frac{-B_{2k} \cdot E_{2k}(q)}{2(2k)!} \frac{(-4X^2)^k}{k} \right) \\ &= \exp \left(\sum_{k \geq 1} \frac{(4^k - 1) B_{2k} \cdot E_{2k}(q)}{2(2k)!} \frac{(-4X^2)^k}{k} \right) \\ &= \sum_{t \geq 0} \left(\sum_{\lambda \vdash t} \prod_{k=1}^t \frac{1}{m_k!} \left(\frac{(4^k - 1) B_{2k} \cdot E_{2k}(q)}{(2k)(2k)!} \right)^{m_k} \right) (-4X^2)^t. \end{aligned}$$

By comparing the coefficients of X^{2t} , we deduce that

$$V_{2t}(q) = 4^t (2t)! \sum_{\lambda \vdash t} \prod_{k=1}^t \frac{1}{m_k!} \left(\frac{(4^k - 1) B_{2k} \cdot E_{2k}(q)}{(2k)(2k)!} \right)^{m_k},$$

which thanks to (1.8) completes the proof of Theorem 1.3 (2). □

Acknowledgment. The authors thank the referee, George Andrews and Wei-Lun Tsai for comments on earlier drafts of this paper. The second author thanks the Thomas Jefferson Fund and the NSF (DMS-2002265 and DMS-2055118). The third author is grateful for the support of a Fulbright Nehru Postdoctoral Fellowship.

Data availability statement. Data sharing not applicable to this article as no datasets were generated or analyzed during the current study.

REFERENCES

- [1] T. Amdeberhan, K. Ono and A. Singh, *MacMahon’s sums-of-divisors and allied q-series*, Adv. Math. **452** (2024), Article 109820.
- [2] G. E. Andrews, *Theory of Partitions*, Cambridge University Press, Cambridge, 1998.
- [3] G. E. Andrews and B. C. Berndt, *Ramanujan’s Lost Notebook, Part II*, Springer, New York, 2009.
- [4] T. M. Apostol, *Introduction to Analytic Number Theory*, Springer-Verlag, New York, 1976.
- [5] B. C. Berndt, S. H. Chan, Z.-G. Liu and H. Yesilyurt, *A new identity for $(q; q)_\infty^{10}$ with an application to Ramanujan’s partition congruence modulo 11*, Q. J. Math. **55** (2004), 13–30.
- [6] B. C. Berndt and A. J. Yee, *A page on Eisenstein series in Ramanujan’s lost notebook*, Glasg. Math. J. **45** (2003), 123–129.
- [7] L. Carlitz and M. V. Subbarao, *A simple proof of the quintuple product identity*, Proc. Amer. Math. Soc. **32** (1972), no. 1, 42–44.

- [8] I. S. Gradshteyn and I. M. Ryzhik, *Table of integrals, series, and products. Translated from the Russian. Translation edited and with a preface by Victor Moll and Daniel Zwillinger*, 8th updated and revised edition, Elsevier/Academic Press, Amsterdam, 2015.
- [9] M. Just and R. Schneider, *Partition Eisenstein series and semi-modular forms* *Partition Eisenstein series and semi-modular forms*, Res. Number Theory **7** (2021), Paper No. 61.
- [10] M. Kaneko and D. Zagier, *A generalized Jacobi theta function and quasimodular forms*, in *The Moduli Space of Curves (Texas Island, 1994)*, Progr. Math., 129, Birkhäuser Boston, Boston, MA, 1995, pp. 165–172.
- [11] K. Ono, *The Web of Modularity: Arithmetic of the Coefficients of Modular Forms and q -Series*, CBMS Regional Conference Series in Mathematics, vol. 102, The American Mathematical Society, Providence, RI, 2004.
- [12] S. Ramanujan, *The Lost Notebook and Other Unpublished Papers*, Narosa Publishing House, New Delhi; Berlin, New York, 1988. Springer-Verlag, Reprinted 2008.
- [13] S. Ramanujan, *On certain arithmetical functions*, Trans. Camb. Phil. Soc. **22** (1916), 159–184.
- [14] R. P. Stanley, *Enumerative Combinatorics, Vol. 2*, Cambridge Stud. Adv. Math., 62 Cambridge University Press, Cambridge, 1999.

Tewodros Amdeberhan
Department of Mathematics, Tulane University
New Orleans
LA 70118
USA
tamdeber@tulane.edu

Ken Ono (Corresponding Author)
Department of Mathematics, University of Virginia
Charlottesville
VA 22904
USA
ko5wk@virginia.edu

Ajit Singh
Department of Mathematics, University of Virginia
Charlottesville
VA 22904
USA
ajit18@iitg.ac.in