

- 3 Wykes T, Steel C, Everitt B, Tarrier N. Cognitive behaviour therapy for schizophrenia: effect sizes, clinical models, and methodological rigor. *Schizophr Bull* 2008; **34**: 523–37.
- 4 Hogarty GE, Kornblith SJ, Greenwald P, DiBarry AL, Cooley S, Ulrich RF, Carter M, Flesher S. Three years trials of personal therapy with schizophrenics living with or independent of family. I: Description of study and effects on relapse rates. *Am J Psychiatry* 1997; **154**: 1504–13.
- 5 Hogarty GE, Greenwald P, Ulrich RF, Kornblith SJ, DiBarry AL, Cooley S, Carter M, Flesher S. Three years trials of personal therapy with schizophrenics living with or independent of family. II: Effects on adjustment of patients. *Am J Psychiatry* 1997; **154**: 1514–24.

Philippa A. Garety, Department of Psychology, Institute of Psychiatry, King's College London, De Crespigny Park, London SE5 8AF. Email: philipa.garety@iop.kcl.ac.uk; **David G. Fowler**, Department of Psychology and Psychiatry, School of Medicine, University of East Anglia, Norwich; **Daniel Freeman**, Department of Psychology, Institute of Psychiatry, King's College London; **Paul Bebbington**, Mental Health Sciences, University College London; **Graham Dunn**, Health Medical Research Group, Community Based Medicine, University of Manchester; **Elizabeth Kuipers**, Department of Psychology, Institute of Psychiatry, King's College London, UK

doi: 10.1192/bjp.193.4.345a

Corrections

Efficacy of antidepressants in juvenile depression: meta-analysis. *BJP*, 193, 10–17. Page 12, Fig. 1: some minor errors (not affecting the data reported) occurred in the published version of this figure. The correct version appears below:




Fig. 2 Forest plot of rate ratios (RR, with 95% CI, on the logarithmic scale) of responses to drug or placebo in 30 randomised double-blind placebo-controlled comparisons of rates of 'response' to antidepressants v. placebo, with overall pooled RR (1.22; 95% CI 1.15–1.31; blue diamond), based on meta-analysis.

Squares represent trials of serotonin reuptake inhibitors (SRIs; 12 trials); circles represent tricyclic antidepressants (TCAs; 14 trials) and other types of antidepressants (4 trials); the size of the data point is proportional to weight defined by study participant number and measurement variance.

Community treatment orders are not a good thing. *British Journal of Psychiatry*, 193, 96–100. Page 98, col. 2: Mary O'Hagan's name was misspelt. The relevant sentence should read: In the words of Mary O'Hagan, who initiated the service user movement in New Zealand and was the first chair of the World Network of Users and Survivors in Psychiatry, 'community treatment orders are oppressive and corrupting – it's tragic that other countries are following Australia and New Zealand's example' (M. O'Hagan, personal communication, 2007).

Computerised cognitive-behavioural therapy for depression: systematic review. *British Journal of Psychiatry*, 193, 181–184. The first sentence of the Acknowledgements (p. 183) should read: This project was funded by the NIHR Health Technology Assessment Programme (project ref. 04/01/01) and commissioned on behalf of NICE. It has been published in full in Health Technology Assessment, Vol. 10, No. 33.

doi: 10.1192/bjp.193.4.346