

## EV1246

### In a period of economical crisis who is at risk for attempted suicide?

I. Coppola<sup>1,\*</sup>, D. Marangon<sup>2</sup>, C. Gramaglia<sup>1</sup>, C. Delicato<sup>1</sup>, S. Di Marco<sup>1</sup>, E. Gattoni<sup>1</sup>, A. Venesia<sup>1</sup>, G.C. Avanzi<sup>3</sup>, L. Castello<sup>3</sup>, F. Bert<sup>4</sup>, M.R. Gualano<sup>4</sup>, R. Siliquini<sup>4</sup>, E. Torre<sup>1</sup>, P. Zeppego<sup>1</sup>

<sup>1</sup> Università del Piemonte Orientale, Translational Medicine, Novara, Italy

<sup>2</sup> SC Psichiatria, AOU Maggiore della Carità, Translational Medicine, Novara, Italy

<sup>3</sup> Emergency Medicine, Università del Piemonte Orientale, Translational Medicine, Novara, Italy

<sup>4</sup> Università degli Studi di Torino, Public Health and Paediatric Sciences, Torino, Italy

\* Corresponding author.

**Introduction** According to Durkheim's theory, suicide attempts increase during periods of rapid economic changes and decrease again when stability is achieved, due to the social integration in the community guaranteed by occupational status. Since 2009 in Europe, it was observed an increase in the suicide rates, in line with the increase of unemployment. Although unemployment is considered a risk factor for suicide, it was observed a noticeable increase in suicide rates among Italian employed men during the last years, probably related to the specific economic scenario and to emerging difficulties in keeping an income to adequately maintain one's own family standard of living.

**Aim** To study attempted suicides with a specific focus and attention on the current Italian economic scenario.

**Methods** Determinants of emergency room visits for psychiatric reasons in patients aged > 16 years were studied prospectively from 2008 to 2015 at the AOU Maggiore della Carità Hospital, Novara, Italy. Each patient was assessed by an experienced psychiatrist with a clinical interview and socio-demographic and clinical features were gathered. Statistical significance was set at  $P \leq 0.05$ .

**Results** Preliminary results suggest an increase in suicide attempts according to the rise of the unemployment rate in Novara. Although in absolute figures most attempters are unemployed, it seems that suicide attempts are more frequent in the group of employed subjects, which includes also people under redundancy fund. This subgroup may be considered halfway between employed and unemployed attempters.

**Conclusions** Data collection and statistical analyses are still ongoing. Implications for clinical practice will be discussed.

**Disclosure of interest** The authors have not supplied their declaration of competing interest.

<http://dx.doi.org/10.1016/j.eurpsy.2016.01.2231>

## EV1247

### Dermal reactivity and resilience in a sample of depressed patients

I. Coppola<sup>1,\*</sup>, E. Gattoni<sup>1</sup>, C. Gramaglia<sup>1</sup>, S. Di Marco<sup>1</sup>, C. Delicato<sup>1</sup>, D. Marangon<sup>2</sup>, A. Feggi<sup>1</sup>, M. Martelli<sup>1</sup>, C. Guerriero<sup>1</sup>, L.H. Thorell<sup>3</sup>, M. Sarchiapone<sup>4</sup>, E. Torre<sup>1</sup>, P. Zeppego<sup>1</sup>

<sup>1</sup> Università del Piemonte Orientale, Translational Medicine, Novara, Italy

<sup>2</sup> SC Psichiatria, AOU Maggiore della Carità, Translational Medicine, Novara, Italy

<sup>3</sup> Linköping University and Emotra AB, Clinical and Experimental Medicine, Svedalen, Sweden

<sup>4</sup> Università del Molise, Medicine and Health Science, Campobasso, Italy

\* Corresponding author.

**Background** Mental disorders are considered a risk factor for suicide: for example, the lifetime risk of suicide is estimated to be 4%–8% in people with mood disorders. The literature suggests that a

history of suicide attempts is significantly related to electrodermal hyporeactivity and that patients with violent suicide attempters exhibit faster habituation of the electrodermal response to repeated neutral tones than patients with non-violent attempts. The impact of depressive symptoms on suicidal ideation may be moderated by resilience. Resilience refers to the ability to maintain or regain mental health despite experiencing adversity. In bipolar and depressive disorders, resilience may influence severity of episodes, frequency of relapse and response to treatment.

**Aim** The objective of our study was to assess the possible correlation among attempted suicide, resilience and dermal reactivity in a sample of depressed patients.

**Methods** We recruited patients with depressive disorders and bipolar depression; data about socio-demographic, clinical features, severity of attempted suicide and suicidal risk were gathered. Patients filled in the Resilience Scale for Adult (RSA). Moreover, for each patient we performed the Edor test in order to assess dermal reactivity.

**Results** Data collection is still ongoing. We expect to find lower levels of resilience in those patients who are hyporeactive and attempted suicide. Moreover, we hypothesize that suicide attempts in the group of hyporeactive patients would be characterized by planning and greater severity. Clinical implications will be discussed.

**Disclosure of interest** The authors have not supplied their declaration of competing interest.

<http://dx.doi.org/10.1016/j.eurpsy.2016.01.2232>

## EV1248

### How does someone who weighs so little feel like a burden to everyone?

F. Coutinho\*, I. Brandão, E. Pereira

Centro Hospitalar de São João, Clínica de Psiquiatria e Saúde Mental, Porto, Portugal

\* Corresponding author.

**Introduction** Anorexia nervosa (AN) is one of the most lethal psychiatric disorders, which is explained partially by starvation related health problems, but also because of high suicide rates. One of the proposed theories to explain a suicide attempt is the interpersonal theory of suicide (IPTS), with its three essential variables: feelings of thwarted belongingness, perceived burdensomeness and the acquired capacity for suicide.

**Objective** To review the literature about suicidality in AN and to present a case report of a patient with restrictive AN who has committed suicide.

**Methods** To review of the literature using the database Medline, through PubMed, with the keywords “anorexia nervosa”, “eating disorders” and “suicide”.

**Results** Suicide attempts and completed suicide are highly prevalent among patients with AN, and some authors suggest that 20–40% of deaths in AN are due to suicide. Recently, the IPTS has been proposed as a mean to explain increased suicidality in AN patients.

**Conclusion** We present a case report about a patient with a long standing AN who has committed suicide after leaving a goodbye note describing strong feelings of perceived burdensomeness directed to her family. It is of utmost importance to continue the study about the phenomenon of suicide in AN to be able to prevent this tragic outcome.

**Disclosure of interest** The authors have not supplied their declaration of competing interest.

<http://dx.doi.org/10.1016/j.eurpsy.2016.01.2233>