

The Journal of Laryngology and Otology

EDITED BY
WALTER HOWARTH
ASSISTANT EDITOR
G. H. BATEMAN
WITH THE COLLABORATION OF
V. E. NEGUS R. G. MACBETH

Contents

SOME OBSERVATIONS ON ATTIC DISEASE AND ITS TREATMENT	A. R. DINGLEY and ✓ E. ZWIEFACH ✓
OTOGENIC MENINGITIS	K. McLAY ✓
ALLERGIC RHINITIS AND INFECTION IN CHILDHOOD	J. PEPYS ✓
THE CASE FOR A NITROGEN GAS SECRETORY FUNCTION OF THE EPITHELIAL LINING OF THE MIDDLE-EAR CLEFT IN THE PRODUCTION OF THE "AIR" IN THE CLEFT	W. STIRK ADAMS ✓
CLINICAL RECORDS—	
A CASE OF TUBERCULOUS INFECTION OF THE MAXILLARY SINUS	M. J. GIBSON AND ✓ J. H. PRAIN ✓
NEUROFIBROMA OF PHARYNX	J. P. STEWART ✓
OSTEOMA OF THE FRONTO-ETHMOIDAL SINUSES WITH PARTICULAR REFERENCE TO THEIR EFFECT ON THE FRONTO-NASAL DUCT	GILBERT H. HOWELLS ✓ and H. MORUS JONES ✓
CLINICAL NOTE—	
CORTISONE IN FENESTRATION SURGERY	C. A. HUTCHINSON ✓
GENERAL NOTES	

London
Headley Brothers Ltd
109 Kingsway WC2

The Journal of Laryngology and Otology

(Founded in 1887 by MORELL MACKENZIE and NORRIS WOLFENDEN)

EDITED BY
WALTER HOWARTH

ASSISTANT EDITOR
G. H. BATEMAN

WITH THE COLLABORATION OF
V. E. NEGUS R. G. MACBETH

1. Original Articles are accepted on the condition that they have not been published elsewhere.

2. Manuscripts should be typewritten, on one side only of the paper, and well spaced.

The Harvard system of recording references should be used, e.g. GREEN, C., and BROWN, D. (1951) *J. Laryng.*, 65, 33. Abbreviations of Journals should follow the style recommended in *World Medical Periodicals*, published by World Health Organization 1952.

It is most important that authors should verify personally the accuracy of every reference before submitting a paper for publication.

3. Galley proofs and engraver's proofs of illustrations are sent to the author. Corrections, which should be kept to a minimum, must be clearly marked, and no extra matter added.

4. Illustration blocks are provided free up to the limit of £5 per article; beyond this authors are expected to pay half the cost.

5. Orders for reprints should be sent when returning galley proofs, and for this purpose special forms are supplied.

6. Authors of original communications on Oto-Laryngology in other journals are invited to send a copy, or two reprints, to the *Journal of Laryngology*. If they are willing, at the same time, to submit their own abstract (in English, French, Italian, or German) it will be welcomed.

7. Editorial communications may be addressed to THE EDITOR, *Journal of Laryngology*, c/o HEADLEY BROTHERS LTD, 109 Kingsway London WC 2.

8. The annual subscription is three guineas sterling (U.S.A. \$10,) post free, and is payable in advance.

9. Single copies will be on sale at 7s. 6d. each; copies of parts up to Vol. LXIII may be purchased at 4s. each.

10. All subscriptions, advertising and business communications should be sent to the publishers, HEADLEY BROTHERS LTD, 109 Kingsway London WC 2.

United States of America

Orders for this *Journal* may be sent either through a local bookseller, or to Messrs G. E. STECHERT & CO., 31-33 East 10th Street, New York, or they may be sent direct to the publishers, HEADLEY BROTHERS LTD, 109 Kingsway London WC 2, England.

Please mention *The Journal of Laryngology* when replying to advertisements

FOR THE EFFECTIVE LOCAL TREATMENT OF SINUSITIS

Penicillin

the potent and penetrating antibacterial
action of 1,500 units of penicillin per ml.
Activity in the presence of pus

plus

Paredrinex

rapid, efficient, and safe vasoconstrictor
that by shrinking oedematous and engorged
tissues enables the penicillin to reach
all affected areas in adequate concentration

equals

PENDEX

*for effective local penicillin therapy
in upper respiratory tract infections,
sinusitis—rhinitis—nasopharyngitis.*

*Available, on prescription
only, in 15 ml. ($\frac{1}{2}$ -oz.) bottles.
Detailed information on
request.*

For cost to N.H.S., please see M. & J. list of costs dated October, 1953.

MENLEY & JAMES, LIMITED, COLDHARBOUR LANE, LONDON, S.E.5

for Smith Kline & French International Co., owner of the trade marks 'Pendex' and 'Paredrinex'
PKP113

Please mention *The Journal of Laryngology* when replying to advertisements

HEARING RE-EDUCATION

without the use of

HEARING AIDS



A further report upon the work of the ArdenTe Audiotherapy Unit working in association with Victor L. Browd, M.D., Adjunct Professor of Otolaryngology, New York Polyclinic Medical School and Hospital.

A PATIENT with, for example, an average hearing loss of 55 to 60 decibels is usually brought to the level of those having a 30 to 35 decibel loss by the amplification of a hearing aid. Experience shows that his hearing performances only partially reflect this acoustic gain. A programme of hearing re-education is then superimposed and additional hearing is regularly acquired thereby. What, then, is to prevent a patient with a 30 to 35 decibel average loss from also acquiring hearing if given the same re-education without a hearing aid? Patients with a 25 and those with a 20 decibel average loss have even greater possibilities. What is to prevent these from being carried to higher levels, and with greater ease, than any patient fitted with a hearing aid? In other words, the proposal to give re-education of hearing without a hearing aid involves no more than this: that the same hearing re-education programme regularly employed with benefit in cases in which a patient has first been brought up to a certain level by a hearing aid be given without a hearing aid in cases in which the patient is already at such a level.

A report, analysis and interpretation of the results of 50 hard-of-hearing persons so rehabilitated will gladly be forwarded upon request to:

THE SECRETARY
THE ARDENTE AUDIO THERAPY UNIT
21/23 Wigmore Street, London, W.1
Telephone: LANGham 8251 (5 lines)

Please mention *The Journal of Laryngology* when replying to advertisements

POST-TONSILLECTOMY COMFORT

Immediate pain relief—Speedier Convalescence

The pain of traumatized tissues following tonsillectomy, demands its own relief—and points the need for analgesia that quickly reaches the irritated area.

ASPERGUM provides 'salivary analgesia' through the simple act of chewing — it brings pain-relieving acetylsalicylic acid into *intimate* and *prolonged* contact with the tonsillar

region, seldom reached even intermittently by gargling. The rhythmic stimulation of muscular action also aids in relieving local spasticity & stiffness: more rapid tissue repair is promoted. Each pleasantly flavoured chewing gum tablet provides 3½ grains acetylsalicylic acid, permitting frequent use. Particularly suitable for children.

Aspergum

for more than two decades a dependable
and welcome aid to patient-comfort

Ethically promoted in packages of 16 tablets and moisture proof bottles of 36 and 250

WHITE LABORATORIES LTD., 428, SOUTHCROFT ROAD, LONDON, S.W.16

THE LARYNGOSCOPE

A Monthly Journal

devoted to the Diseases of the
EAR, NOSE AND THROAT

*Official organ for the American Laryngological,
Rhinological and Otological Society*

Price \$14.00 per year

Canada \$13.00 per year

MAX A. GOLDSTEIN, M.D.
FOUNDER


THEODORE E. WALSH, M.D.
EDITOR

640 SOUTH KINGSHIGHWAY
SAINT LOUIS 10, MO.

Please mention *The Journal of Laryngology* when replying to advertisements

The New KEELER OTOSCOPE

This latest addition to the **KEELER** range of instruments embodies good design and a quite remarkable performance that will interest every otologist



SPECIAL FEATURES

- Five all-NYLON specula, sizes 2.5, 3.5, 4.5, 5.5 and 8 mm. Unbreakable, easily sterilised, more comfortable than metal and affording a better seal when used with Siegle attachment
- Lateral adjustments for lamp and magnifier to permit a shadowless field of view
- Intense, homogeneous illumination
- Ample space for surgical manipulation
- Screw clamp for rapid and positive location of specula
- Lightweight, robust, well-balanced and inexpensive
- Supplied with spare lamp, specula and Siegle attachment complete in case
- Also incorporated in our range of E.E.N.T. Diagnostic Sets

39 Wigmore Street
LONDON, W.1
England



617 S. 52nd Street
Philadelphia, 43
Pa., U.S.A.

Please mention *The Journal of Laryngology* when replying to advertisements

AMPLIVOX MODEL 61 THE CLINICAL AUDIOMETER OF INTERNATIONAL REPUTE

"Messrs. Amplivox were among the first firms to produce an audiometer in this country, and their larger model is one of the best instruments of the kind now made in the world at a competitive price."—The LANCET, 23-12-50.

- Eleven exact test frequencies 125-12,000 c.p.s.
- Simplified hearing loss dial. The same set of figures is read for both bone and air conduction at all frequencies, and for speech.
- Bone conduction tests can be made from 125-4,000 c.p.s.
- Masking Tone calibrated in decibels, permitting accurate control of masking.
- Double Air receivers enable test tones to be switched instantly from ear to ear.
- Speech test circuit monitors speech level, permitting accurate measurement of hearing loss for speech.
- Loudness Balance Control establishes presence of recruitment in monaural deafness.
- Automatic voltage compensator.

Recruitment Test Set accessory establishes presence of recruitment by amplitude modulation, enabling each ear to be tested independently.

Speech Turntable, English made P.B. and Harvard Spondee Records available.

Full details are available from the manufacturers who will gladly arrange demonstrations if required.



The Basic Instrument of Modern Otology

ACCURATE • COMPLETE • SIMPLE TO OPERATE

AMPLIVOX LTD., 2 BENTINCK ST., LONDON, W.1 (Welbeck 2591)

Please mention *The Journal of Laryngology* when replying to advertisements

ADVERTISEMENTS

CONTENTS

	PAGE
SOME OBSERVATIONS ON ATTIC DISEASE AND ITS TREATMENT. A. R. Dingley and E. Zwiefach (London)	127
OTOGENIC MENINGITIS. K. McLay (Edinburgh)	140
ALLERGIC RHINITIS AND INFECTION IN CHILDHOOD. J. Pepys (London)	147
A CASE FOR A NITROGEN GAS SECRETORY FUNCTION OF THE EPITHELIAL LINING OF THE MIDDLE-EAR CLEFT IN THE PRODUCTION OF THE "AIR" IN THE CLEFT. W. Stirk Adams (Birmingham)	152
CLINICAL RECORDS—	
A Case of Tuberculous Infection of the Maxillary Sinus. M. J. Gibson and J. H. Prain (Dundee)	170
Neurofibroma of Pharynx. J. P. Stewart (Edinburgh)	174
Osteoma of the Fronto-Ethmoidal Sinuses with particular reference to their effect on the Fronto-Nasal Duct. Gilbert H. Howells and H. Morus Jones (London)	177
CLINICAL NOTE—	
Cortisone in Fenestration Surgery. C. A. Hutchinson (Salisbury)	189
GENERAL NOTES	191

For Advertisement space in this Journal apply to :
HEADLEY BROTHERS Ltd, 109 Kingsway London WC 2

Please mention *The Journal of Laryngology* when replying to advertisements