

support for dissemination/implementation functions, cross-institutional collaboration/networking, and leadership composition. **RESULTS/ANTICIPATED RESULTS:** In total, 52% of hubs will renew under the new PAR in the next few years, providing an incentive to demonstrate dissemination capacity (although hubs will likely lag in operationalizing these activities until they are funded). A third of hubs (34%) represent more than one academic/research institution, and almost 80% of hubs have more than one clinical affiliate. To accommodate these different levels of institutional complexity, broad diffusion will require multi-modal, locally adapted dissemination efforts. Only 25% of hubs have capacity to undertake additional dissemination activities, and only 27% provide formal D&I support, suggesting that additional capacity/support will be needed to operationalize the CTSA dissemination mission. In total, 30% of hubs participate in cross-institutional collaboration/networking, so many may not have existing norms/tools supporting inter-institutional collaboration, but 77% include leadership from outside the School of Medicine, facilitating effective intra-institutional dissemination. **DISCUSSION/SIGNIFICANCE OF IMPACT:** Understanding more about CTSA hubs as both adopters and transmitters of innovation can facilitate strategic use of these sites as a built-in dissemination network to amplify the reach and impact of clinical innovation and improve population health. Based on this initial analysis, the CTSA network does not appear to be fully primed for broad, rapid dissemination of innovation across its sites. In-depth interviews are being conducted to investigate CTSA hubs' perceptions of their dissemination capacity and roles as adopters and transmitters of innovation.

2224

### Determining if intestinal commensal bacteria enhance the frequency of reassortment of an enteric, segmented virus, reovirus

Matthew Lanahan, Andrea Erickson<sup>1</sup> and Julie Pfeiffer<sup>1</sup>

<sup>1</sup> Department of Molecular Microbiology, University of Texas Southwestern Medical Center, Dallas, TX, USA

**OBJECTIVES/SPECIFIC AIMS:** The overall goal is to determine if intestinal commensal bacteria play a role in enteric virus evolution. We will use reovirus, an enteric segmented virus, to investigate specific goals. First, we will determine if specific bacterial species enhance the coinfection frequency of 2 separate strains of reovirus. Second, we will determine if the presence/absence of different bacterial species in the microbiota of mice results in different reovirus reassortment frequencies. Finally, we will discover if reassortant reovirus is present in human populations. **METHODS/STUDY POPULATION:** My first goal is to determine if specific bacterial species enhance the coinfection frequency of 2 strains of reovirus. In our lab, we have a panel of commensal intestinal bacterial strains, as well as a number of lab adapted bacterial strains. We will use this panel of bacteria to determine if reovirus binds to different species of bacteria using a binding assay involving radiolabeled virus. Additionally, we will determine if specific species of bacteria alter the coinfection frequency through a Flow cytometry based assay. This will involve mixing virus with bacteria, infecting cells in culture, and straining for reovirus proteins for flow cytometry. Our second goal is to determine if specific bacteria promote reassortment of reovirus in a mouse model of infection. To do this, we will use gnotobiotic techniques to create mice harboring different intestinal bacteria populations. Mice will be infected with 2 strains of reovirus, and then feces and organs will be collected. Progeny virus will be subjected to a plaque assay on 2 different types of cells. The first type of cells will be normal cells in culture in which all viable viruses will form plaques. The second will be a cell line that stably expresses siRNAs against specific reovirus segments in which only specific reassortants will form plaques. These 2 plaque assays will be used to quantify the total number of viruses present and the total number of reassortant viruses present. Additionally, SDS-PAGE and RT-PCR will be used to confirm reassortants. Our third goal is to determine if reassortant reovirus is present in infected humans. To do this, I will obtain feces from reovirus-infected children and isolate reovirus. One specific reovirus reassortant is known to propagate in dual-infected mice. I will use the plaque assay technique to determine if this reassortant is also present in humans. To determine if other reassortants are present, I will use RT-PCR and SDS-PAGE. **RESULTS/ANTICIPATED RESULTS:** Based on previous studies with other enteric viruses, we suspect that specific bacterial species bind reovirus strains with different efficiencies. It is likely that a number of bacterial species will promote coinfection. The bacterial strains that binds both reovirus strains at a high efficiency will likely enhance coinfection by the greatest amount. It is likely that mice harboring different bacterial populations will produce different reovirus reassortment frequencies. We predict that bacteria that enhance reovirus coinfection in vitro should also enhance reovirus reassortment in our mouse model. Therefore, mice specifically lacking bacteria that promote coinfection should have significantly lower amounts of reassortant reovirus. It will be important to control for the overall amount of replication within mice with different microbiotas, as this will affect the basal reassortment frequency. We suspect

that reovirus reassortants are present in humans. Work done both in vitro and in mouse models indicates that reassortment happens at high frequencies. Additionally, one specific reassortant commonly propagates in mice due to an enhanced cellular attachment phenotype. Therefore, we predict that this reassortant also commonly emerges after coinfection and reassortment in humans. **DISCUSSION/SIGNIFICANCE OF IMPACT:** Segmented viruses, such as influenza and rotavirus, are important human pathogens. Viral reassortment poses a unique threat to humans, as it enables new viruses to emerge and cause pandemics or epidemics. However, little is known about what factors promote viral reassortment. This study will provide insight into a novel mechanism of segmented virus evolution.

2421

### Development and validation of a translational rat model of neonatal abstinence syndrome

Lisa Brents<sup>1</sup>, Bryce A. Griffin<sup>1</sup>, Caitlin Caperton<sup>1</sup>, Lauren Russell<sup>1</sup>, Christian Cabanlong<sup>1</sup>, Catheryn Wilson<sup>1</sup>, Kyle Urquhart<sup>1</sup>, Brad Martins<sup>1</sup>, Amy L. Patton<sup>1</sup>, Alexander W. Alund<sup>1</sup>, S. Michael Owens<sup>2</sup>, William E. Fantegrossi<sup>2</sup> and Jeffery H. Moran<sup>1</sup>

<sup>1</sup> PinPoint Testing, LLC and University of Arkansas for Medical Sciences; <sup>2</sup> University of Arkansas Translational Research Institute

**OBJECTIVES/SPECIFIC AIMS:** Rodent models can be used to study neonatal abstinence syndrome (NAS), but the applicability of findings from the models to NAS in humans is not well understood. The objective of this study was to develop a rat model of norbuprenorphine-induced NAS and validate its translational value by comparing blood concentrations in the norbuprenorphine-treated pregnant rat to those previously reported in pregnant women undergoing buprenorphine treatment. **METHODS/STUDY POPULATION:** Pregnant Long-Evans rats were implanted with 14-day osmotic minipumps containing vehicle, morphine (positive control), or norbuprenorphine (0.3–3 mg/kg/d) on gestation day 9. Within 12 hours of delivery, pups were tested for spontaneous or precipitated opioid withdrawal by injecting them with saline (10 mL/kg, i.p.) or naltrexone (1 or 10 mg/kg, i.p.), respectively, and observing them for well-validated neonatal withdrawal signs. Blood was sampled via indwelling jugular catheters from a subset of norbuprenorphine-treated dams on gestation day 8, 10, 13, 17, and 20. Norbuprenorphine concentrations in whole blood samples were quantified using LC/MS/MS. **RESULTS/ANTICIPATED RESULTS:** Blood concentrations of norbuprenorphine in rats exposed to 1–3 mg/kg/d of norbuprenorphine were similar to levels previously reported in pregnant women undergoing buprenorphine treatment. Pups born to dams treated with these doses exhibited robust withdrawal signs. Blood concentrations of norbuprenorphine decreased across gestation, which is similar to previous reports in humans. **DISCUSSION/SIGNIFICANCE OF IMPACT:** These results suggest that dosing dams with 1–3 mg/kg/day norbuprenorphine produces maternal blood concentrations and withdrawal severity similar to those previously reported in humans. This provides evidence that, at these doses, this model is useful for testing hypotheses about norbuprenorphine that are applicable to NAS in humans.

2525

### Development of human cell-based screening assays to detect subject-specific drug-response variability

Francesca Stillitano, Joshua Mayourian, Jaydev Dave, Jean-Sébastien Hulot and Roger J. Hajjar

Icahn School of Medicine at Mount Sinai

**OBJECTIVES/SPECIFIC AIMS:** The goals of this study are to develop a human-based screening assay for testing individual drug reactions and investigate the mechanism underlying susceptibility to develop dILQT. **METHODS/STUDY POPULATION:** We derived iPSC-CMs from 10 subjects with a high sensitivity to Sotalol (high-S group) and 10 subjects with no changes in QT interval after administration of the same drug (low-S group). Multielectrode array (MEA) was used to measure field potential duration, a surrogate to the QT interval in the electrocardiogram, in iPSC-CMs under basal conditions and in response to increasing concentrations of Sotalol. Transcriptomic profiling of iPSC-CMs from high-S Versus low-S groups was performed using RNA-sequencing. A parameter sensitivity analysis was performed on the Paci *et al.* iPSC-CM mathematical model to further support the lead hits identified via RNA-sequencing. **RESULTS/ANTICIPATED RESULTS:** Cardiac differentiation resulted in the generation of iPSC-CMs with appropriate cardiac channel