

1 **Nature's contributions to people in drylands**

2

3 David Eldridge¹, Chenxu Wang², Yanxu Liu^{2*}, Jingyi Ding², Yan Li², Xutong Wu², Changjia
4 Li²

5

6 1. Centre for Ecosystem Sciences, School of Biological, Earth and Environmental Sciences,
7 University of New South Wales, Sydney, NSW, 2052, Australia

8

9 2. State Key Laboratory of Earth Surface Processes and Resource Ecology, Faculty of
10 Geographical Science, Beijing Normal University, Beijing, China

11

12 * corresponding author: email: yanxuliu@bnu.edu.cn

13

14 *Abstract*

15

16 Humans depend heavily on nature. Drylands are home to 2.5 billion people, but the extent to
17 which nature contributes to people (NCP) in drylands has been little explored. We examined
18 the global contribution of nature to people, aiming to compare drylands and non-drylands. We
19 predicted a lower contribution in drylands than non-drylands, largely because of the sparser
20 population densities (peoples' needs) and more degraded status of natural resources (lower
21 potential contribution). Consistent with expectation, nature's contribution was about 30%
22 lower in drylands, with significantly lower values for drylands in Asia, Oceania, Africa and
23 South America, but no difference for Europe and North America. Differences were due
24 mainly to lower contributions from material and regulating contributions, i.e., the regulation
25 of air-quality, climate, water quantity and flow, soil protection and the supply of woody
26 material, and potentially, lower use by people in drylands. Predicted declines in rainfall and

1

This peer-reviewed article has been accepted for publication but not yet copyedited or typeset, and so may be subject to change during the production process. The article is considered published and may be cited using its DOI.

10.1017/dry.2024.2

This is an Open Access article, distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives licence (<http://creativecommons.org/licenses/by-nc-nd/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is unaltered and is properly cited. The written permission of Cambridge University Press must be obtained for commercial re-use or in order to create a derivative work.

27 increasing temperature are likely to place increasing pressure on nature to contribute to
28 human well-being in drylands. A better understanding of nature's contributions to people
29 would improve our ability to allocate limited resources and achieve sustainable development
30 in drylands.

31

32 **Keywords:** regulating services, environmental quality, global population, rangelands, arid
33 and semiarid

34

35 **Introduction**

36

37 The physical, social, cultural, and spiritual wellbeing of humans is highly dependent on
38 nature (Diaz et al., 2018; Hill et al., 2021). Nature encompasses not only organisms and their
39 ecosystems, but ecological and evolutionary processes on Earth, resulting in both positive and
40 negative consequences for humans and their quality of life (IPES, 2019). Nature's
41 contribution to people (NCP) has been defined as 'all the contributions, both positive and
42 negative, of living nature (diversity of organisms, ecosystems, and their associated ecological
43 and evolutionary processes) to people's quality of life (Diaz et al. 2018). Nature's
44 contributions can be organised broadly into three categories: material contributions (e.g., food
45 and energy), non-material contributions (e.g., recreation, spiritual services and experiences),
46 and regulating contributions (e.g., clean air and good water quality). Material and regulating
47 contributions are similar to the elements captured in the ecosystem services paradigm,
48 whereas non-material contributions include attributes that relate to the quality of life, belief
49 systems, or nature-based experiences (Diaz et al., 2018; Hill et al., 2021).

50

51 Implicit in the NCP concept therefore is not only what nature can provide, but also what
52 people need; social justice, spiritual beliefs and their links to the natural environment (Pascua
53 et al., 2017), regardless of whether these needs are realised (Chaplin Kramer et al., 2019).

54 The magnitude of these contributions, therefore, would be expected to be greater where
55 people have the strongest association with nature and the greatest needs (Chaplin Kramer et
56 al., 2019). We would expect, therefore, stronger contributions to human societies where
57 people have a closer connection with nature or where they derive a living from the land. Such
58 associations would be expected to be greater in drylands, where 90% of the human
59 population have relatively low standards of living (Reynolds et al., 2007). Yet, previous
60 global assessments (e.g., Liu et al., 2023) suggest that it is in arid and semi-arid environments

61 and developing countries, mostly drylands, where ecosystem degradation fails to sustain the
62 needs of people, i.e., areas with the greatest benefit gaps (*sensu* Hill et al., 2021) compared
63 with wealthy, less marginalised communities.

64
65 Despite global assessments of NCP (Chaplin-Kramer et al., 2019), we have a relatively
66 limited understanding of how nature contributes to the lives and welfare of people in
67 drylands. Drylands are important because they account for almost 40% of the terrestrial land
68 area, and are home to about 38% of the world's population (2.5 billion people; Huang et al.,
69 2017). Moreover, large areas of drylands are devoted to primary production, particularly
70 fodder production for livestock (Právělie, 2016). This makes them more vulnerable to
71 environmental changes associated with increasing aridity (Huang et al., 2017) than other,
72 more mesic environments (Berdugo et al., 2022). Many people in drylands are marginalised,
73 often in areas of political conflict (Global Conflict Tracker, 2023), with low standards of
74 living and sometimes poor nutrition (Právělie 2016). Areas where overall contributions are
75 low have been shown to be associated with transitional climates (Liu et al., 2023), reflecting
76 ongoing degradation and declines in nature itself, through for example, land clearing,
77 desertification, and atmospheric pollution. This suit of physical and environmental conditions
78 likely places drylands at a greater risk of famine and global tragedy. Exploring how nature
79 can contribute to people in drylands is critical if we are to balance the competing needs of
80 people and the natural environment, a more equitable human society, and work towards
81 achieving sustainable development of drylands.

82
83 Here we report on a study where we used environmental, social and biological data as
84 surrogates of 18 contributions that nature can make to people in drylands (Hill et al., 2021).
85 These contributions comprise seven regulating, six material, and five non-material categories.
86 Drylands are defined as areas where the ratio of evaporation to average annual precipitation
87 exceeds 0.65 (MEA, 2005), including dry subhumid, semi-arid, arid and hyper-arid areas.
88 Previous studies have focussed on nature's contributions at global scales (Chaplin-Kramer et
89 al., 2019; Hill et al., 2021) or explored the service provision of drylands rather than
90 associating provision and people's need (Maestre et al., 2022). Traditional ecosystem service
91 approaches have focussed only on regulating and provisioning contributions such as primary
92 production, carbon, and food, but neglect the actual non-material needs of dryland people.

93

94 Our study links the provision of tangible goods and services with human needs, endeavouring
95 to provide insights into the connection between potential contributions from nature and the
96 capacity of people in drylands to use these contributions. We asked the following two
97 questions: First, does the average contribution to people in drylands differ from that in non-
98 drylands, and if so, what is the nature of the difference in these contributions? We posit that
99 drylands would have a lower overall (average) contribution. Our rationale is that drylands are
100 less densely populated, and their environmental resources more degraded and susceptible to
101 global changes than non-drylands (Hill et al., 2021). Second, have drylands exhibited greater
102 temporal declines in contributions over the past decades (1992 – 2018) than non-drylands?
103 We would expect the affirmative, given the generally greater declines in environmental
104 quality in drylands than non-drylands over the past half century, though this could be masked
105 by a larger population size and therefore greater human need over that period.

106

107 **Methods**

108

109 We used the datasets of Liu et al., (2023; see Supplementary Text S1), and the assessment are
110 briefly described as follows. A general simplified flow chart illustrating the process of
111 calculating nature's contribution to people using Habitat creation and maintenance (NCP1) as
112 an example is presented in Figure 1.

113

114 *Spatial datasets*

115

116 We reclassified the European Space Agency Climate Change Initiative-Land Cover (ESA
117 CCI-LC) product (European Space Agency 2018), as the core data indicating the change of
118 nature for NCP assessment. Complex ecosystem classifications reduce the accuracy of some
119 NCP calculations, so ecosystem classes need to be consolidated and reclassified to harmonize
120 different terminologies. Reclassification and consolidation can simplify different
121 terminologies prior to analyses, such as combining multiple forests into a single forest class.
122 This process simplifies the computational steps and permits a more rapid assessment of NCP.
123 We used 20 spatial datasets to make the 18 NCP assessments. Most raster datasets had spatial
124 resolutions finer than 10 km, providing sufficient pixels for each sub-basin unit. Maps of
125 richness of mammals, birds and amphibians (Jenkins et al., 2013) at a resolution of 10 km
126 were downloaded from BiodiversityMapping.org. The Global Inventory Modeling and
127 Mapping Studies (GIMMS) provided the vegetation leaf area index (LAI)3g product at a

128 spatial resolution of 1/12 arc degrees (Zhu et al., 2013). The global human settlement layer
129 (GHSL) was downloaded from the Joint Research Center (JRC) and included grids for built-
130 up areas, population, and settlements (Corbane et al., 2019). The gross primary production
131 (GPP) dataset was estimated using a revised light use efficiency model, with a spatial
132 resolution of 0.05 arc degrees (Zheng et al., 2020). The vectorized Global Mangrove Watch
133 (GMW) datasets were transformed into 1 km spatial resolution data (Bunting et al., 2018) and
134 evapotranspiration (ET) was a synthesized product with a 1 km spatial resolution (Elnashar et
135 al., 2021). The MODIS (Terra Moderate Resolution Imaging Spectroradiometer Land Water
136 Mask (MOD44W) Version 6 data product was accessed from the Land Processes Distributed
137 Active Archive Center (LP DAAC) with a spatial resolution of 250 m (Carroll et al., 2017).
138 Annual streamflow maps were obtained from the FLO1K dataset at a spatial resolution of 1
139 km (Barbarossa et al., 2018).

140

141 A pesticide risk score, based on the most popular active pesticide ingredient, was at a spatial
142 resolution of 1/12 arc degrees (Tang et al., 2021). The soil erosion score was evaluated based
143 on studies by Liu et al., (2019) at a spatial resolution of 1/12 arc degrees. The Harmonized
144 World Soil Database was at a spatial resolution of 1 km (Fischer et al., 2008). Slope and
145 elevation data were obtained from the Shuttle Radar Topography Mission digital elevation
146 model at a resolution of 3 arc seconds (Jarvis et al., 2008). The aridity index (AI) was
147 determined as the relationship between precipitation and evapotranspiration and mapped at a
148 resolution of 30 arc seconds (Trabucco and Zomer 2019). Floodplain data were at a 250 m
149 resolution (Nardi et al., 2019). Data on the yield and aggregated value of crop production
150 were derived from the Spatial Production Allocation Model dataset in 2010 (SPAM2010) at a
151 spatial resolution of 1/12 arc degrees (Yu et al., 2020). The “best crop” map that indicated the
152 maximum achievable bioenergy yields was derived from the dataset of lignocellulosic
153 bioenergy crops at a spatial resolution of 0.5 arc degrees (Li et al., 2020). Aboveground
154 carbon biomass density data were derived from a 2010 harmonized map at a spatial resolution
155 of 300 m (Spawn et al., 2010). Nighttime light data were obtained from a harmonized dataset
156 from two satellites at a spatial resolution of 30 arc seconds (Li et al., 2020). The locations of
157 natural and mixed world heritage sites were obtained from WHC.UNESCO.org. The vector
158 road dataset was downloaded from the Socioeconomic Data and Applications Center
159 (SEDAC) and named Global Roads Open Access Data Set, Version 1 (gROADSv1; SEDAC
160 2013). We applied the HydroBasin level 06 in the HydroATLAS database to take advantage
161 of the nested sub-basins at multiple scales for regionalization (Linke et al., 2019). To

162 accommodate the spatial resolution of the various spatial datasets described above, the units
163 smaller than 500 km² were merged into adjacent largest units in order to include more than
164 four pixels of 1/12 arc degree raster data in a basin. This resulted in a database of 15204 basin
165 units.

166

167 *Spatial assessment*

168

169 The assessment of NCPs uses an indicator-based approach with two indicators: 1) nature's
170 potential contribution, and 2) nature's actual contribution to people (Table 1). Nature's
171 potential contribution relates to the potential to provide resources, services, knowledge or
172 inspiration. For example, nature contributes to the regulation of crop pests (NCP10) by
173 supporting a diverse community of birds (Mayne et al., 2023). This contribution depends on
174 whether a given basin unit supports crops that require this pest regulation or whether there are
175 people that can benefit from this pest regulation. Although the potential contribution may be
176 large, the actual contribution may be zero, due to an absence of people or crops, for example,
177 the inability of people to use products derived from nature. Because actual human
178 requirements from nature generally increase as population size increases, we set population
179 as static so that we could observe changes in NCPs driven by nature changes alone, i.e., in the
180 absence of population increase. Put simply, increases in human requirements could lead to an
181 increased NCP assessment, which could mask any potential threats of natural ecosystem loss,
182 and lead to perverse landscape management outcomes (Chaplin-Cramer et al., 2019).

183

184 The indicator framework was used to calculate a globally rapid assessment of all NCPs to
185 identify spatiotemporal heterogeneity of the distribution rather than simulating a defined
186 value for biophysical units. In order to develop a rapid assessment framework, no more than
187 three global parameters were used for each NCP, except for hazard regulation (NCP9), which
188 required four parameters to adequately parameterize. Details of the procedures and datasets
189 used to calculate each NCP are given in Supplementary Text S1 and Figure 1. The lowest
190 values of the parameters were assigned a value of 0, and the threshold value of 1 was set as
191 the 90th percentile values of each originally assessed NCP value in 1992. All the value
192 exceeding the threshold should be assigned as 1. By min-max normalization, the normalized
193 value of every NCP was in the range of 0–1. Note that we did not change people's need
194 between the 1992 and 2018 (see Supplementary Text S1).

195

196 Linear models (Bates et al., 2015) were used to examine differences in mean NCP values
197 between drylands and non-drylands in relation to 1) six continents, 2) individual
198 contributions, and 3) between 1992 and 2018. We tested for the correlation between the value
199 of each NCP and population size using Pearson's r . Analytical tests were performed in the R
200 statistical software (R Core Team 2021) prior to linear modelling to ensure that the data met
201 the necessary assumptions implicit in linear modelling.

202

203 **Results**

204

205 *Nature generally contributes less to people in drylands*

206

207 The global average value of nature's contribution to people was about 30% lower in dryland
208 than non-drylands ($\chi^2 = 47.3$, $df = 1$, 114, $P < 0.001$, Fig. S1), consistent with our prediction.
209 Nature's contributions to drylands were significantly lower for Africa, Asia, Oceania and
210 South America, but there were no differences for Europe or North America (dryland/non-
211 dryland by continent interaction: $\chi^2 = 15.2$, $df = 5$, 114, $P = 0.009$; Fig. 2). There was a small
212 (albeit non-significant; $P > 0.21$) decline in nature's contribution to people with increasing
213 continent size for drylands, but not for non-drylands. We also found evidence of an increase
214 in the magnitude of NCP with increasing population size, particularly for air quality
215 regulation (NCP3), food (NCP12), medicine (NCP14) and learning/inspiration (NCP15;
216 Table S1). Identity (NCP17) declined strongly in both drylands and non-drylands with
217 increasing population size (Table S1).

218

219 We then focused on the average contribution to people across global drylands for different
220 locations, i.e., the average value across all 18 contributions and considered both potential
221 (Fig. 3a) and actual (Fig. 3b) contribution. We found extensive areas of low actual NCP in
222 North Africa (Algeria, Libya, Niger, Mauritania, Mali, Chad, Egypt, northern Sudan,
223 northern Ethiopia), West Africa (Namibia and South Africa), the west coast of South
224 America (northern Chile and Patagonia), much of inland central Australia, the Arabian
225 Peninsula, western Eurasia (Afghanistan, Iran, Turkmenistan), and west-central China and
226 Russia. Conversely, high values were more insular, and occurred in southern India, north-
227 eastern China, the Iberian Peninsula, western Turkey, south-eastern South Africa, north-
228 western USA, and a narrow strip in north-eastern Brazil and coastal Eastern Australia (Figs.
229 3a and 3b). Although potential and actual contributions were spatially similar overall, actual

230 contributions were greater for the Iberian Peninsula, the Indian subcontinent, and the eastern
231 side of the Eurasian drylands (Figs. 3a and 3b).

232

233 There were, however, large differences between drylands and non-drylands for specific
234 contributions. For example, drylands contributed less to six of the 18 NCP categories i.e.,
235 regulation of air quality (NCP3), climate (NCP4), water quantity and flow (NCP6), soil
236 protection (NCP8; mainly in Asia and Oceania, Table S2), woody material (NCP13) and
237 options (NCP18; Fig. 4).

238

239 *Spatio-temporal changes in nature's contributions*

240

241 We found a general decline in NCP between 1992 and 2018 across all contributions and for
242 both drylands and non-drylands ($-0.47 \pm 0.71\%$, mean \pm SE) but this masked the changes in
243 some contributions. For example, the average contribution by nature declined more in
244 drylands than non-drylands for 10 contributions: habitat (NCP1), pollination (NCP2), oceans
245 (NCP5), water quality (NCP7), soil protection (NCP8), hazard regulation (NCP9), pest
246 regulation (NCP10) and bioenergy (NCP11), medicine (NCP14) and experience (NCP16), but
247 increased for climate (NCP4), water quality/flow (NCP6), food (NCP12), woody material
248 (NCP13) and options (NCP18; Table 2, Table S2), again consistent with our second
249 prediction.

250

251 We also detected some spatial changes over the 26 years. The value of climate regulation in
252 drylands increased in north-central and southern Africa, northern and south-western
253 Australia, northern India, western Iran and western USA, but declined in central Australia
254 and western China (Figs. 5 and 6). For water quantity/flow regulation, we detected increases
255 in north-central Africa, the Arabian Peninsula, northern Australia, much of mainland China,
256 India and Iran, but declines were evident in northern and southern Africa, central, northern
257 and eastern Australia, the Iberian Peninsula, the western USA, and the west coast of South
258 America. Similarly, there were some spatial declines in the value of pest regulation (NCP10)
259 in drylands across extensive areas of Africa, the western USA and the western coast of South
260 America, western Iran, northern India, central China, and large areas of Africa and Central
261 Australia.

262

263 **Discussion**

264

265 We used an indicator framework to compare nature's contribution to people in drylands with
266 non-drylands. Unsurprisingly, the magnitude of this contribution was about 30% lower in
267 drylands. These differences, however, were inconsistent across continents, with significantly
268 lower values for drylands in Asia, Oceania, Africa and South America, but no difference in
269 Europe and North America. Furthermore, we identified some hotspots of low contribution in
270 North Africa, the Arabian Peninsula, central Australia, and west-central China, and high
271 values in southern India, north-eastern China, the Iberian Peninsula, eastern Australia, and the
272 north-west coast of the United States of America. Finally, potential and actual NCP values
273 were similar, except for the heavily populated areas in Spain, India and China. Our results are
274 consistent with the understanding that nature's contribution to people is likely to be lower
275 where the quality of the natural ecosystem or its capacity to produce is low (Chaplin-Cramer
276 et al., 2019) and in sparsely populated drylands where the capacity of people to use nature's
277 products is low (Brauman et al., 2020). Our results also suggest that the magnitude of
278 nature's contribution globally will decline as drylands expand at the expense of non-drylands.

279

280 *A spatial understanding of nature's contribution to people in global drylands*

281

282 Within those continents with a lower drylands contribution, we found that the reduction in
283 contribution was due largely to a reduction in the magnitude of regulating contributions such
284 as climate (NCP4), water quantity and flow (NCP6), soil protection (NCP8) and the
285 production of woody material (NCP13; Fig. 4), reflecting a generally stronger reliance upon
286 primary resources by drylands in contrast to non-drylands (Brauman et al., 2020, Hill et al.,
287 2021).

288

289 Three dryland areas characterised by low levels of regulating contributions and sparse
290 populations densities are North Africa (e.g., Algeria, Tunisia, Libya, and Egypt), the Arabian
291 Peninsula, and central Australia (Fig. 3). Low levels of climate regulation (NCP4) across
292 these three areas result from the sparse forest and limited mid- and groundstorey cover (< 5%
293 Maestre et al., 2021) dominated by short stature woody perennials and low stature herbaceous
294 biomass (Fischer and Turner, 1978; Stafford Smith and Morton, 1990; Le Houerou, 2000;
295 Brinkmann et al., 2011), but they often support a high plant species diversity (Maestre et al.,
296 2021). Intense browsing and grazing by livestock, the dominant land use in drylands, reduces
297 plant cover (e.g., Brinkmann et al., 2009), thus reducing the potential for capture of

298 greenhouse gases and increasing climate-driven consequences for humans (Brouman et al.,
299 2020). Vegetation cover and biomass are also critical parameters that influence the generation
300 of aerosols, which are high over the Arabian Peninsula (Tandule et al., 2022) and North
301 Africa (Gherboudj et al., 2017). It is unsurprising, therefore, that these three regions have a
302 relatively lower capacity to support stable soils (NCP8) or extensive wood production
303 (NCP13). The potential to produce wood suitable for saw milling (NCP13) is also low due to
304 the predominance of lower stature vegetation (shrublands at the expense of forests), highly
305 variable precipitation, and high evapotranspiration (Stafford Smith and Morton 1990). The
306 only substantial difference in Europe was the lower value for woody material (NCP13) in
307 drylands than non-drylands (Table S2), reflecting the dominance of short stature xerophytic
308 shrubs with low potential for forestry in the drylands of southern Spain, southern Italy and
309 west-central Poland. Importantly, yields of woody material are likely to decline due to
310 increased risk of droughts and wildfires in Europe exacerbated by changing climates (Górriz-
311 Mifsud et al., 2000).

312

313 Large areas of North Africa remote from coastal influences are mapped as having low actual
314 values of water quantity and flow regulation (NCP6, Fig. 3b). Many North African countries
315 face severe environmental challenges due to water scarcity (Hamed et al., 2018), which
316 compromise agricultural industries that rely heavily on water supply (Radhouane, 2013).
317 Surface and groundwater sources are sparsely distributed in North Africa and the Arabian
318 Peninsula (Siebert et al., 2015), and surface water is scarce in Central Australia, where it is
319 held for only short periods in isolated depressions and ephemeral waterways (Brim Box et al.,
320 2022). Consequently, most perennial vegetation is dependent entirely on groundwater (Eamus
321 et al., 2006). Large areas of the Arabian Peninsula also lack surface water but have the
322 capacity to access aquifers recharged from sporadic river flooding (UNDP/RBAS 2013).
323 Overall, these three example drylands are more sensitive to increasing dryness associated
324 with climate change than non-drylands.

325

326 Implicit in the NCP concept is population size, and therefore potential contribution to people.
327 We found generally positive relationships between NCP values and population size (Table
328 S2), consistent with our understanding that population size and ecosystem production are
329 positively correlated (Luch, 2007). Our three focal drylands are all relatively sparsely
330 populated, with densities of 0.1, 1 and about 4 people km⁻² for Central Australia, the Arabian
331 Peninsula and North Africa, respectively (Gapminder-Systems Globalis, 2022). Values of

332 some NCPs (e.g., air quality, food production, medicine, pest regulation, and learning
333 inspiration) were significantly related to population density in both drylands and non-
334 drylands (Table S2). However identity (NCP17) declined with increasing population size,
335 possibly reflecting the alienation of traditional knowledge at large spatial scales or where
336 populations are changing rapidly (Darvill and Lindo, 2015).

337

338 Average contribution values for drylands in two continents, Europe and North America were
339 similar to values in non-drylands. North American and European (southern Spain, Sicily)
340 drylands are densely populated, have relatively large GDPs, and highly mechanised primary
341 production (Al Shamsi et al., 2018; Bauer and Iles, 2023; Martinez-Valderrama et al., 2024).
342 For example, the drylands in Almeria, on the Iberian Peninsula in Spain support a mixture of
343 wooded Mediterranean forest and grassland located within a matrix of industrial agriculture
344 such as greenhouses and irrigated agriculture (Ghafraoui et al., 2023) and support a moderate
345 population density of about 80 persons km⁻². This is reflected in the high value of NCP12
346 (food) in drylands (Table S2). Extensive areas of farmland in Spain are located near Córdoba
347 and Seville, the most developed locations since antiquity (Martinez-Valderrama et al., 2023),
348 and this area is regarded as a food bowl for Europe (Ayuda and Pinilla, 2021). Furthermore,
349 desert regions of Almeria are highly iconic and display unique landscape features ('badlands'
350 Zgłobicki et al., 2021) that many city dwellers will not normally experience. People prefer
351 these natural, albeit highly eroded, landscape more than greenhouses. This likely reflects the
352 high value that the population places on natural landscapes and landscape diversity, which
353 should be reflected in learning and inspiration (NCP15) and identity (NCP16). North
354 American drylands are also highly developed, support large urban centres, and include iconic
355 desert environments with extensive natural and mixed ecosystems (NCP1) with potential for
356 bioenergy production (NCP11; Nabhan et al., 2020), and areas that are accessible to people
357 for experience of the natural world (NCP16, Table S2).

358

359 *Greater decline in nature's contribution to people in drylands*

360

361 The magnitude of nature's contribution has declined markedly over the past half-century
362 (e.g., Broumann et al., 2020; Liu et al., 2023), and results between 1992 and 2018 indicate a
363 substantially greater declines in drylands than non-drylands (Table 2. Figs. 5 & 6).
364 Importantly, the greatest declines were for pollination (NCP2, 65% decline), soil protection
365 (NCP8, 56%), hazard regulation (NCP9, 42%), pest regulation (NCP10, 54%), medicine

366 (NCP14, 59%), and experience (NCP16, 42%). Potential contributions have declined for
367 virtually all regulating contributions, e.g., plant pollination and pest regulation (Potts et al.,
368 2016), and most declines have been due to a loss in environmental quality (e.g. Liu et al.,
369 2023). Non-material declines are also evident, for example, with increased urbanisation
370 removing local communities and indigenous people from their connections with the land and
371 natural environments (Soga and Gaston, 2016).

372

373 Many of nature's contributions, particularly material contributions, are based on vegetation-
374 related proxies. One might expect, therefore, a generally lower contribution in drylands than
375 non-drylands, though this was not always the case (e.g., Fig. 4). Improvements in database
376 quality and the availability of more specialised data on different contributions at finer spatial
377 scales in drylands should lead to a more reliable assessment of the relative differences
378 between drylands and non-drylands, particularly if new proxies are more closely aligned to
379 particular contributions. It is clear that the benefits accruing from nature are likely to be
380 greatest where nature is most intact (Chaplin-Kramer et al., 2019), suggesting that areas
381 suffering from environmental degradation will contribute less. The consequences of
382 increasing aridity are that nature's contributions to drylands will continue to decline,
383 particularly for dryland types that are most susceptible to changing climates. Distinct dryland
384 sub-types are likely to respond differently to climate change (e.g., hyper-arid compared with
385 dry subhumid) simply because nature's contribution depends on both the potential
386 contribution (which is dependent on vegetation and therefore rainfall) and realised
387 contribution (lower population sizes and therefore lower demand for material, non-material
388 and cultural contributions). Thus, more detailed assessment of different dryland subtypes
389 would likely reveal how increasing global dryness might alter nature's contributions. Our
390 results indicate that any declines in the environmental quality of drylands will have not only
391 environmental implications, but will impact human health (medicine) and the physical and
392 psychological experiences that humans derive from nature.

393

394 *Conclusions*

395

396 We used relatively predictable, intuitive, yet simple proxies to calculate nature's contribution
397 to people in drylands. We acknowledge, however, that our capacity to improve these
398 estimates is hampered by the lack of available databases at the scale commensurate with
399 drylands and non-drylands, and/or the lack of more nuanced information that is more closely

400 aligned with a given contribution. This is particularly relevant for non-material contributions
401 that relate to belief systems or personal experiences. Thus, our assessments can only be based
402 on global databases and remotely sensed, broad-scale proxies. Advances in remote sensing
403 technologies and access to databases at finer spatial scales should allow us to refine our
404 assessment of nature's contribution in drylands, across large areas where data are sparsely
405 distributed. Nonetheless, our study demonstrates that lower contributions to people in
406 drylands can be attributed to declining quality of environmental resources in natural systems
407 (Liu et al., 2023; Table S1). The value of these attributes declines with declining rainfall and
408 increasing dryness, yet their value (realised and potential) also increases with increasing
409 population pressure. Predicted large-scale increases in aridity, combined with marked
410 population increase and therefore accelerated land degradation (Právělie, 2016) are likely to
411 place increasing pressure on nature to contribute to the physical wellbeing and function of
412 drylands, its biota and people.

413

414 **Conflict of interest**

415 The authors declare that they have no conflict of interest.

416

417 **Acknowledgments**

418 This work was supported by the Hermon Slade Foundation, the National Natural Science
419 Foundation of China (41991235, 41991232), and the Fundamental Research Funds for the
420 Central Universities in China.

421

422 **Author contribution**

423 David Eldridge and Yanxu Liu designed the research and Chenxu Wang performed the
424 analyses. David Eldridge drafted the manuscript. Yan Li, Jingyi Ding, Changjia Li, and
425 Xutong Wu contributed to the design and editing of the manuscript.

426

427 **References**

428

429 Al Shamsi, KB, Compagnoni, A, Timpanaro, G, Cosentino, S, Guarnaccia, P (2018) A
430 sustainable organic production model for “Food Sovereignty” in the United Arab Emirates
431 and Sicily-Italy. Sustainability 10, 620 doi:10.3390/su10030620

432

433 Ayude, MI, Pinilla, V (2021) Agricultural exports and economic development in Spain during
434 the first wave of globalisation, *Scandinavian Economic History Review* 69, 199-216, DOI:
435 [10.1080/03585522.2020.1786450](https://doi.org/10.1080/03585522.2020.1786450)
436

437 Barbarossa, V, Huijbregts, M, Beusen, A, Beck, HE, King, H, Schipper, AM. (2018) FLO1K,
438 global maps of mean, maximum and minimum annual streamflow at 1 km resolution from
439 1960 through 2015. *Scientific Data* 5, 180052. <https://doi.org/10.1038/sdata.2018.52>
440

441 Bates, D, Mächler, M, Bolker, B, Walker, S (2015) Fitting linear mixed-effects models
442 Using lme4. *Journal of Statistical Software* 67. <https://doi.org/10.18637/jss.v067.i01>
443

444 Baur, P, Iles, A (2023) Replacing humans with machines: a historical look at technology
445 politics in California agriculture. *Agriculture and Human Values* 40, 113–140.
446

447 Berdugo, M, Gaitán, JJ, Delgado-Baquerizo, M, Crowther, TW, Dakos, V (2022) Prevalence
448 and drivers of abrupt vegetation shifts in global drylands. *Proceedings of the National*
449 *Academy of Sciences USA* 119(43):e2123393119. doi: 10.1073/pnas.2123393119.
450

451 Brauman, KA, Garibaldo, LA, Polansky, S et al., (2020) Global trends in nature's
452 contributions to people. *Proceedings of the National Academy of Sciences USA*
453 DOI/10.1073/pnas.2010473117
454

455 Brim Box, J, Leiper, I, Nano, C, Stokeld, D, Jobson, P, Tomlinson, A, Cobban, D, Bond, T,
456 Randall, D. and Box, P. (2022) Mapping terrestrial groundwater-dependent ecosystems in
457 arid Australia using Landsat-8 time-series data and singular value decomposition. *Remote*
458 *Sensing in Ecology and Conservation* 8, 464-476.
459

460 Brinkmann, K, Dickhoefer, U, Schlecht, E, & Buerkert, A. (2011) Quantification of
461 aboveground rangeland productivity and anthropogenic degradation on the Arabian Peninsula
462 using Landsat imagery and field inventory data. *Remote Sensing of Environment* 115, 465–
463 474.
464

465 Brinkmann, K, Patzelt, A, Dickhoefer, U, Schlecht, E, & Buerkert, A. (2009) Vegetation
466 patterns and diversity along an altitudinal and a grazing gradient in the Jabal al Akhdar
467 mountain range of northern Oman. *Journal of Arid Environments* 73, 1035–1045.
468

469 Bouman, T, Verschoor, M, Albers, C.J, Böhm, G, Fisher, S.D, Poortinga, W, Whitmarsh, L,
470 and Steg, L. (2020) When worry about climate change leads to climate action: How values,
471 worry and personal responsibility relate to various climate actions. *Global Environmental*
472 *Change* 62, 102061, <https://doi.org/10.1016/j.gloenvcha.2020.102061>.
473

474 Bunting, P, Rosenqvist, A, Hilarides, L, Lucas, RM, Thomas, N, Tadono, T, Worthington, TA,
475 Spalding, M, Murray, NJ, Rebelo, L-M. (2022) Global Mangrove Extent Change 1996–2020:
476 Global Mangrove Watch Version 3.0. *Remote Sensing*, 14, 3657.
477 <https://doi.org/10.3390/rs14153657>
478

479 Carroll, ML, DiMiceli, CM, Townshend, JRG, Sohlberg, RA, Elders, AI, Devadiga, S, Sayer,
480 AM, Levy RC (2017) Development of an operational land water mask for MODIS Collection
481 6, and influence on downstream data products, *International Journal of Digital Earth* 10, 2,
482 207-218, DOI: 10.1080/17538947.2016.1232756
483

484 Center for International Earth Science Information Network - CIESIN - Columbia University,
485 and Information Technology Outreach Services - ITOS - University of Georgia. 2013. Global
486 Roads Open Access Data Set, Version 1 (gROADSv1) Palisades, New York: NASA
487 Socioeconomic Data and Applications Center (SEDAC).
488 <https://doi.org/10.7927/H4VD6WCT>. Accessed 3 March 2022.
489

490 Chaplin-Kramer, R. et al., (2019) Global modeling of nature’s contributions to people.
491 *Science* 366, 255-258
492

493 Darvill, R, Lindo, Z. (2015) Quantifying and mapping ecosystem service use across
494 stakeholder groups: Implications for conservation with priorities for cultural values.
495 *Ecosystem Services* 13, 153-161.
496

497 Diaz, S. et al., (2018) Assessing nature’s contributions to people. *Science* 359, 270–272.
498

499 Eamus, D, Froend, R, Loomes, R, Hose, G. & Murray, B. (2006) A functional methodology
500 for determining the groundwater regime needed to maintain the health of groundwater-
501 dependent vegetation. *Australian Journal of Botany* 54, 97–114.

502

503 Elnashar, A, Wang, L, Wu, B, Zhu, W, and Zeng, H. (2018) Synthesis of global actual
504 evapotranspiration from 1982 to 2019. *Earth System Science Data* 13, 447–480.

505

506 Fischer, G, Nachtergaele, F., Prieler, S, van Velthuizen, HT, Verelst, L, Wiberg, D. (2008)
507 *Global Agro-ecological Zones Assessment for Agriculture (GAEZ 2008)*. IIASA, Laxenburg,
508 Austria and FAO, Rome, Italy.

509

510 Fischer, RA, Turner, NC (1978) Plant Productivity in the Arid and Semiarid Zones. *Annual*
511 *Review of Plant Physiology* 29, 277–317.

512

513 Gapminder - Population v7 (2022); Gapminder - Systema Globalis (2022); HYDE (2017);
514 United Nations - World Population Prospects – with major processing by Our World in Data.

515

516 El Ghafroui, Y, Quintas-Soriano, C, Pacheco-Romero, M, Murillo-López, B.E, and Castro,
517 A. (2023) Diverse values of nature shape human connection to dryland landscapes in Spain.
518 *Journal of Arid Environments* 216, 105023

519

520 Gherboudj, I, NaseemaBeegum, S. & Ghedira, H. (2017) Identifying natural dust source
521 regions over the Middle-East and North-Africa: Estimation of dust emission potential. *Earth-*
522 *Science Reviews* 165, 342–355.

523

524 Global Conflict Tracker (2023) Center for Preventative Action, Council on Foreign
525 Relations. <https://www.cfr.org/global-conflict-tracker/?category=us>.

526

527 Górriz-Mifsud, E, Ameztegui, A, González, J.R, Trasobares, A. (2022) Climate-Smart
528 Forestry Case Study: Spain. In: Hetemäki, L, Kangas, J, Peltola, H. (eds) *Forest Bioeconomy*
529 *and Climate Change. Managing Forest Ecosystems, Vol 42*. Springer, Cham.

530 https://doi.org/10.1007/978-3-030-99206-4_13

531

532 Hamed, Y, Hadji, R, Redhaounia, B. et al., (2018) Climate impact on surface and
533 groundwater in North Africa: a global synthesis of findings and recommendations. Euro-
534 Mediterranean Journal of Environmental Integrity 3, 25 [https://doi.org/10.1007/s41207-018-](https://doi.org/10.1007/s41207-018-0067-8)
535 [0067-8](https://doi.org/10.1007/s41207-018-0067-8)
536

537 Hill, R, Díaz, S, Pascual, U, Stenseke, M, Molnár, Z, Van Velden, J. (2021) Nature's
538 contributions to people: Weaving plural perspectives. *One Earth* 4, 910-915.
539

540 Huang, J, Li, Y, Fu, C, Chen, F, Fu, Q, Dai, A, Shinoda, M, Ma, Z, Guo, W, Li, Z, Zhang, L,
541 Liu, Y, Yu, H, He, Y, Xie, Y, Guan, X, Ji, M, Lin, L, Wang, S, Yan, H, Wang, G. (2017)
542 Dryland climate change: Recent progress and challenges. *Reviews in Geophysics* 55, 719–
543 778.
544

545 IPBES (2019) Global assessment report on biodiversity and ecosystem services of the
546 Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services. E.S.
547 Brondizio, J. Settele, S, Díaz, and H.T. Ngo (editors) IPBES secretariat, Bonn, Germany.
548 <https://doi.org/10.5281/zenodo.3831673>
549

550 Jarvis, A, Reuter, HI, Nelson, A, Guevara, E. (2008) Hole-Filled SRTM for the Globe Version
551 4. CGIAR-CSI SRTM 90 m Database. <https://srtm.csi.cgiar.org>
552

553 Jenkins, CN, Pimm, SL, Joppa, LN. (2013) Global patterns of terrestrial vertebrate diversity
554 and conservation. *Proceedings of the National Academy of Sciences USA*. 110 (28) E2602-
555 E2610 <https://doi.org/10.1073/pnas.1302251110>
556

557 Kashyap, R, Grover, S, Puri, H, Kaur, S, Singh, J, Sandhu, KS, Kaur, G, Kharva, H, Kaur, S,
558 Kaur, B. (2023) Insect and Pest Management for Sustaining Crop Production Under
559 Changing Climatic Patterns of Drylands. In: Naorem, A, Machiwal, D. (eds) *Enhancing*
560 *Resilience of Dryland Agriculture Under Changing Climate*. Springer, Singapore.
561 https://doi.org/10.1007/978-981-19-9159-2_21
562

563 Le Houerou, HN. (2000) Restoration and rehabilitation of arid and semiarid mediterranean
564 ecosystems in North Africa and West Asia: A Review. *Arid Soil Research and Rehabilitation*,
565 14, 3–14.

566

567 Li, W, Ciais, P, Stehfest, E, van Vuuren, D, Popp, A, Arneth, A, Di Fulvio, F, Doelman, J,
568 Humpenöder, F, Harper, AB, Park, T, Makowski, D, Havlik, P, Obersteiner, M, Wang, J,
569 Krause, A, Liu, W. (2020) Mapping the yields of lignocellulosic bioenergy crops from
570 observations at the global scale. *Earth Systems Science Data* 12, 789–804.

571

572 Linke, S, Lehner, B, Ouellet Dallaire, C, Ariwi, J, Grill, G, Anand, M, Beames, P, Burchard-
573 Levine, V, Maxwell, S, Moidu, H, Tan F, Thieme, M. (2019) Global hydro-environmental
574 sub-basin and river reach characteristics at high spatial resolution. *Scientific Data* 6, 283
575 DOI: 10.1038/s41597-019-0300-6

576

577 Liu, Y, Fu, B, Wang, S, Rhodes, J.R, Li, Y, Zhao, W, Li, C, Zhou, S, and Wang, C. (2023)
578 Global Assessment of Nature’s Contribution to People. *Science Bulletin* 68, 424-435.

579

580 Luck, GW. (2007) The relationships between net primary productivity, human population
581 density and species conservation. *Journal of Biogeography* 34, 201-212.

582

583 Maestre, F.T, Benito, B.M, Berdugo, M, Concostrina-Zubiri, L, Delgado-Baquerizo, M,
584 Eldridge, D.J, Guirado, E, Gross, N, Kéfi, S, Le Bagousse-Pinguet, Y, Ochoa-Hueso, R. and
585 Soliveres, S. (2021) Biogeography of global drylands. *New Phytologist* 231, 540-558.

586

587 Maestre, F.T., Le Bagousse-Pinguet, Y., Delgado-Baquerizo, M., Eldridge, D.J., Saiz, H.,
588 Berdugo, M., et al., (2022) Grazing and ecosystem service delivery in global drylands.
589 *Science* 378 (6622): 915-920.

590

591 Martínez-Valderrama, J, Gartzia, R., Olcina, J, Guriado, E, Ibanez, J, Maestre, FT. (2024)
592 Uberizing agriculture in drylands: a few enriched, everyone endangered. *Water Resources*
593 *Management* 38, 193-114.

594

595 Mayne, SJ, King, DI, Andersen, JC, Elkinton, JS. (2023) Crop-specific effectiveness of birds
596 as agents of pest control. *Agriculture, Ecosystems & Environment*, 348, 108395,
597 <https://doi.org/10.1016/j.agee.2023.108395>

598

599 Millennium Ecosystem Assessment, 2005. Drylands Systems". Chapter 22 in: Ecosystems
600 and Human Wellbeing: Current State and Trends, Volume 1. Island Press.
601

602 Nabhan, GP, Riordan, EC, Monti, L, Rea, AM, Wilder, BT, Ezcurra, E, Mabry, JB, Aronson,
603 J, Barron-Gafford, GA, García, JM, Búrquez, A, Crews, TE, Mirocha, P, Hodgson, WC.
604 (2020) An Aridamerican model for agriculture in a hotter, water scarce world. *Plants, People,*
605 *Planet 2*, 627–639. But don't expect you
606

607 Nardi, F, Annis, A, Di Baldassarre, G, Vivoni, ER, Grimaldi, S. (2019) GFPLAIN250m, a
608 global high-resolution dataset of Earth's floodplains. *Scientific Data* 6, 180309.
609 <https://doi.org/10.1038/sdata.2018.309>
610

611 Pascua, P, McMillen, H, Ticktin, T, Vaughan, M, Winter, K. (2017) Beyond services: A
612 process and framework to incorporate cultural, genealogical, place-based, and indigenous
613 relationships in ecosystem service assessments. *Ecosystem Services* 26, 465-475
614

615 Potts, SG, Imperatriz-Fonseca, V, Ngo HT, Aixen, MA, Biesmeijer, JC, Breeze, TD, Dicks,
616 LV, Garibaldi, LA, Hill, R, Settele, J, Vanbergen, AJ. (2016) Safeguarding pollinators and
617 their values to human well-being. *Nature* 540: 220-229
618

619 Právělie, R. (2016) Drylands extent and environmental issues. A global approach. *Earth-*
620 *Science Reviews* 161, 259-278.
621

622 R Core Team (2021). R: A language and environment for statistical computing. R Foundation
623 for Statistical Computing, Vienna, Austria. URL <https://www.R-project.org/>.
624

625 Radhouane, L (2013) Climate change impacts on North African countries and on some
626 Tunisian economic sectors. *J Agri Environ Intern Dev (JAEID)* 107, 101–113.
627

628 Reynolds, JF, Stafford Smith, MD, Lambin II, EF, Turner, BK, Mortimore, M, Batterbury,
629 SPJ, Downing, T, Dowlatabadi, H, Fernández, RJ, Herrick, JE, Huber-Sannwald, E, Jiang, H,
630 Leemans, R, Lynam, T, Maestre, FT, Ayarza, M, Walker, B. (2007) Global desertification:
631 building a science for dryland development. *Science* 316, 847–851.
632

633 Siebert, S, Kummu, M, Porkka, M, Döll, P, Ramankutty, N, Scanlon, BR. (2015) A global
634 data set of the extent of irrigated land from 1900 to 2005. *Hydrology and Earth Systems*
635 *Science* 19, 1521–1545.

636

637 Soga, M, Gaston, KJ. (2016) Extinction of experience: The loss of human-nature interactions.
638 *Frontiers in Ecology and Environment* 14, 94-101

639

640 Spawn, S.A., Sullivan, C.C., Lark, T.J., Gibbs, HK. (2020) Harmonized global maps of above
641 and belowground biomass carbon density in the year 2010. *Scientific Data* 7, 112.
642 <https://doi.org/10.1038/s41597-020-0444-4>)

643

644 Stafford Smith, DM, Morton, SR. (1999) A framework for arid Australia. *Journal of Arid*
645 *Environments* 18, 255-278.

646

647 Still, CJ, Berry, JA, Collatz, GJ, DeFries, RS. (2003) Global distribution of C₃ and C₄
648 vegetation: Carbon cycle implications, *Global Biogeochemical Cycles* 17, 1006,
649 [doi:10.1029/2001GB001807](https://doi.org/10.1029/2001GB001807).

650

651 Tandule, CR, Gogoi, MM, Kotalo, RG, Suresh Babu, S. (2022) On the net primary
652 productivity over the Arabian Sea due to the reduction in mineral dust deposition. *Scientific*
653 *Reports* 7761. <https://doi.org/10.1038>

654

655 Tang, F.H.M., Lenzen, M., McBratney, A, Maggi, F. (2021) Risk of pesticide pollution at the
656 global scale. *Nature Geoscience* 14, 206–210.

657

658 Trabucco, A, Zomer, R. (2019) Global Aridity Index and Potential Evapotranspiration (ET0)
659 Climate Database v2. figshare. Fileset. <https://doi.org/10.6084/m9.figshare.7504448.v3>)

660

661 UNDP/RBAS (2013) *Water Governance in the Arab Region - Managing Scarcity and*
662 *Securing the Future*. United Nations Development Program, Regional Bureau for Arab States,
663 New York, 10017, USA. 182 pp.

664

665 Yu, QY, You, LZ, Wood-Sichra, U, Ru, Y, Joglekar, AKB, Fritz, S, Xiong, W, Lu, M, Wu,
666 W, Yang, P. (2020) A cultivated planet in 2010-part 2: The global gridded agricultural-
667 production maps. *Earth System Science Data* 12, 3545-3572

668

669 Zgólbicki, W., Poesen, J., Joshi, V., Sóle-Benet, A., De Geeter, S. (2021). Gullies and
670 Badlands as Geoheritage Sites. In: Singh, R., Wei, D., Anand, S. (eds) *Global Geographical
671 Heritage, Geoparks and Geotourism. Advances in Geographical and Environmental Sciences.*
672 Springer, Singapore. https://doi.org/10.1007/978-981-15-4956-4_9

673

674 Zheng, Y, Shen, RQ, Wang, Y, Li, X, Liu, S, Liang, S, Chen, JM, Ju, W, Zhang, Li, Yuan,
675 W. (2020) Improved estimate of global gross primary production for reproducing its long-
676 term variation, 1982-2017. *Earth System Science Data* 12, 2725-2746.

677

678

679 Zhu, Z, Bi, J, Pan, Y, Ganguly, S, Anav, A, Xu, L, Samanta, A, Piao, S, Nemani, RR, Myneni,
680 RB. (2013) Global Data Sets of Vegetation Leaf Area Index (LAI)3g and Fraction of
681 Photosynthetically Active Radiation (FPAR)3g Derived from Global Inventory Modeling and
682 Mapping Studies (GIMMS) Normalized Difference Vegetation Index (NDVI3g) for the
683 Period 1981 to 2011. *Remote Sensing* 5, 927-948.

684

685 **Table 1.** Description of nature’s actual and potential contribution to people (adapted from Liu et al., 2023). The data sources indicate the source
 686 of information used to assess both the potential and actual contribution to people, as well as the parameter (weighted parameter) used in the
 687 calculation of actual contribution.

NCP	Description (after Diaz et al. 2018)	Contribution type	Potential contribution to people	Actual contribution to people	Weighted parameter	Data sources
NCP1: habitat	Habitat creation and maintenance	Regulating	Natural and mix ecosystems: potential natural habitats	Actual animal biodiversity (mammals, birds and amphibians)	Animal biodiversity	Amphibian, mammal and bird richness BiodiversityMapping.org
NCP2: crop pollination	Pollination and dispersal of seeds and other propagules	Regulating	Mix ecosystems: key place of seed dispersal to cropland	Production for cross-pollinated crops: yield of crops required pollination	Production for cross-pollinated crops	Extent of natural vegetation within 3 km buffer of cropland
NCP3: air quality regulation	Regulation of air quality	Regulating	Vegetation Leaf Area Index in natural and mix ecosystems: potential pollution entraining vegetation	Built-up land requiring pollution entrainment: actual emission from human habitat required entrainment	Built-up land and vegetation leaf area index	LAI data from Global Inventory Modeling and Mapping Studies (GIMMS)
NCP4: climate regulation	Regulation of climate	Regulating	Gross primary productivity in perennial vegetation: carbon sequestration	Default: not valued because of the global scale requirement	Gross primary productivity	Gross primary productivity (GPP) databases

NCP5: ocean acidification regulation	Regulation of ocean acidification	Regulating	Amount of mangrove forest in coast: key place of long-term carbon sink from ocean	Default: not valued because of the global scale requirement	Distribution of mangroves	Mangrove distribution data (1996 – 2016)
NCP6: water quantity and flow regulation	Regulation of freshwater quantity location and timing	Regulating	Evapotranspiration in natural and mix ecosystems: participation of ecosystem in water cycle	Streamflow: actual requirement for flow regulation by ecological processes including evapotranspiration	Evapotranspiration and streamflow	Terrestrial evapotranspiration data
NCP7: water quality regulation	Regulation of freshwater and coastal water quality	Regulating	Natural ecosystems surrounded rivers: natural capacity on decontamination	Nonpoint source pollution indicated by pesticide risk: actual requirement for decontamination	Water location and pesticide risk	Permanent water bodies within the MOD44B database
NCP8: soil protection	Formation protection and decontamination of soils and sediments	Regulating	Soil retention of natural ecosystems: potential amount of soil retention	Soil fertility indicated by organic carbon: actual contribution of fertility retention	Soil retention amount and organic carbon	Harmonized World Soil Database in conjunction with Revised Universal Soil Loss Equation (RUSLE) modelling
NCP9: hazard regulation	Regulation of hazards and extreme events	Regulating	Natural ecosystems reducing landslide, desertification, flood and storm tides	Value of crop productions: agricultural value benefit from hazard prevention	Distribution of drylands and floodplains, slope, and crop production value	Combination of Global- Aridity_ET0 database, GFPLAIN 250 m dataset, and VP CROP A production database

NCP10: pest regulation	Regulation of detrimental organisms and biological processes	Regulating	Bird biodiversity in mix ecosystems: pest enemy diversity for agricultural production	Value of crop productions: actual value of crops prevented from pest	Bird biodiversity and crop production value	Aggregated value of production dataset
NCP11: bioenergy	Energy	Material	Shrub, grass and mix ecosystems: potential land for bioenergy plants in high probability	Potential lignocellulosic bioenergy crops: score of bioenergy production could be harvested	Lignocellulosic bioenergy crops	Lignocellulosic bioenergy crops dataset
NCP12: food	Food and feed	Material	Cultivated and mix ecosystems: potential land for food production	Yield of production for food crops: actual yield of food crops	Yield of food crops	Food crop yield data from the global synergy cropland layer in the SPAM dataset
NCP13: wood material	Materials, companionship and labour	Material	Forest ecosystems: potential land for logging	Aboveground biomass carbon density: actual yield of logging	Aboveground biomass carbon density	Aboveground carbon biomass datasets
NCP14: medicine	Medical, biochemical and genetic resources	Material	Diversity of natural and mix ecosystems: species diversity indicated by landscape diversity	Rural population: local people potentially using native herbal medicine	Natural landscape diversity and rural population	Shannon's landscape diversity index and global population databases
NCP15 learning and inspiration	Learning and inspiration	Non-material (cultural)	The diversity of ecosystem: diversity of nature,	Social development indicated by nighttime light:	Landscape diversity and nighttime light	Shannon's landscape diversity index and nighttime light databases

			include artificial landscape	people's requirement in a developing society		
NCP16: experience	Physical and psychological experiences	Non-material (cultural)	Density of natural and mix World Heritage sites: proximity of unique natural landscape	Accessibility indicated by road density: people's accessibility to get the unique experience	Density of natural and mixed world heritages and road density	World heritage database
NCP17: identity	Supporting identities	Non-material (cultural)	Change rate of landcover: landscape stability	Population on the changed landscape: actual amount of people within identity shaping	Rates of land cover and population changes	European Space Agency Climate Change Initiative Land Cover datasets
NCP18: options	Maintenance of options	Mixture of all three	Diversity of the other 17 NCPs	Diversity of nature to provide future benefits	Shannon's diversity index	Shannon's diversity index of NCPs 1 to 17

688

689

690

691


692 **Table 2.** Percentage change in NCP for drylands and non-drylands between 1992 and 2018
693 and the dryland trend.

NCP code	Description	Dryland	Non-dryland	Dryland trend
NCP1	Habitat	-1.19	-0.89	Greater decline
NCP2	Crop pollination	-2.02	0.70	Greater decline
NCP3	Air quality regulation	-0.63	-0.73	Lower decline
NCP4	Climate regulation	11.11	1.10	Increase
NCP5	Ocean acidification regulation	-5.35	-4.32	Greater decline
NCP6	Water quantity and flow regulation	4.03	-0.20	Increase
NCP7	Water quality regulation	-1.86	-1.20	Greater decline
NCP8	Soil protection	-2.28	-1.01	Greater decline
NCP9	Hazard regulation	-2.62	-1.52	Greater decline
NCP10	Pest regulation	-5.43	-2.52	Greater decline
NCP11	Bioenergy	-0.48	1.53	Greater decline
NCP12	Food	2.95	2.31	Increase
NCP13	Wood material	1.11	-0.67	Increase
NCP14	Medicine	-1.47	-0.61	Greater decline
NCP15	Learning and inspiration	2.38	2.42	Lower increase
NCP16	Experience	-1.32	-0.77	Greater decline
NCP17	Identity	-3.32	-5.44	Lower decline
NCP18	Options	0.53	0.41	Increase

694

695


696 Fig. 1



697

698


699 Fig. 2



700

701 Nature's contribution to people (mean ± SE) for drylands and non-drylands by continent. Asterisks
702 indicate a significant difference between dryland and non-dryland at $P < 0.05$ (based on linear
703 modelling). The actual number of dryland and non-dryland basins are Asia: 2,241 dryland and 2,774
704 non-dryland basins; Africa: 2,456 and 888; Europe: 285 and 924; North America: 677 and 2,274; South
705 America: 577 and 1,204; Oceania: 848 and 56. For analytical purposes, analyses are based on 18
706 observations, i.e., the average values for each NCP type.


707



709

710 Mean (a) potential and (b) actual contribution of nature to people in global drylands (average of 18 NCP
711 categories).

712




714

715 Mean contribution of each of the 18 NCP categories for drylands and non-drylands. Asterisks indicate
 716 a significant difference in contribution value between dryland and non-dryland at $P < 0.05$. Analyses
 717 are based on linear modelling.

718

719 Fig. 5




720

721 Spatiotemporal change in NCP between 1992 and 2018 for NCP1 to NCP8 for drylands

722

723 Fig. 6



724

725 Spatiotemporal change in NCP between 1992 and 2018 for NCP9 to NCP18 for drylands

726