

---

## INSTRUCTIONS TO CONTRIBUTORS

The *Journal of Helminthology* publishes papers on all aspects of animal parasitic helminths, particularly those of medical or veterinary importance, but only in exceptional circumstances will systematic or taxonomic studies be acceptable.

Manuscripts, which must be in English or French (with an English summary) should be accompanied by a letter signed by *all* the authors and should be addressed to:

The Editor, Journal of Helminthology,  
London School of Hygiene and Tropical Medicine,  
Winches Farm Field Station,  
395 Hatfield Road,  
St Albans, Herts AL4 0XQ.  
England.

Two copies of a typescript, on size A4 paper with double spacing, should be submitted. Papers should be preceded by a short abstract and will normally have the following sections: brief Introduction; Materials and Methods; Results; Discussion; Acknowledgements; References. However, the form of the paper may vary, depending on its subject matter; recent past issues should be consulted for a suitable form. Research Notes should also be preceded by a brief abstract. Illustrations should be drawn in Indian ink, preferably not more than double the final size. Care should be taken that all illustrations fit into the format of the Journal. The maximum size an illustration will be printed is 12.0×20.0 cm. Where many separate drawings are made, some indication of how they may be grouped to make a corporate plate without undue wastage of space, should be indicated. Some indication of scale (preferably a scale bar) should normally be given on the figure. Photocopies of illustrations should be enclosed for refereeing purposes. Lettering and numbering, which must be of a high standard, should be added by the author, with due regard for subsequent reduction.

Photographs should be glossy prints of the same size as they are to appear in the Journal (maximum size 12.0×20.0 cm). Composite prints must be mounted and can have the separate photographs abutting; they will then have a separating line inserted by the printers. All figures and letters on photographs must be inserted by the author.

Information should not be repeated in the text and in tables or figures. The legends to tables and to figures should be sufficiently detailed for the information to be understood without reference to the text.

References should be given in alphabetical order with the full title of the journal. The following are examples:

DUKE, B. O. L. (1971) The ecology of onchocerciasis in man and animals. In: *Ecology and Physiology of parasites* (editor, A. M. Fallis) pp. 213–222. Adam Hilger Ltd.: London.

JAMES, C. & WEBBE, G. (1973) A comparison of Egyptian and East African strains of *Schistosoma haematobium*. *Journal of Helminthology*, **47**, 49–59.

25 offprints are provided free of charge; additional copies may be ordered at the proof stage.

# Contents

	Pages
The ability of <i>Aedes aegypti</i> mosquitoes to survive and transmit infective larvae of <i>Brugia pahangi</i> over successive blood meals. S. W. LINDSAY and D. A. DENHAM	159-168
The antifilarial activity of a novel group of organic arsenicals upon <i>Brugia pahangi</i> . D. A. DENHAM, S. L. OXENHAM, I. MIDWINTER and E. A. H. FRIEDHEIM	169-172
Fractionation of <i>Fasciola hepatica</i> tegument antigens and their application to the serodiagnosis of experimental fascioliasis by the enzyme-linked immunosorbent assay. G. V. HILLYER and A. E. SERRANO	173-178
Immunological aspects in the rejection process of <i>Hymenolepis muris-sylvaticae</i> from CFLP and NMRI mice. C. VAN HAEREN and P. H. DE RYCKE	179-186
Chitin synthesis in zygotes of <i>Ascaris suum</i> . P. DUBINSKÝ, M. RYBOŠ, L. TURČEKOVÁ and E. OSSIKOVSKI	187-192
Nervous system in <i>Olveria indica</i> , a rumen paramphistome (Digenea) of bovines, as revealed by non-specific esterase staining. N. MISHRA and V. TANDON	193-199
Studies on the enzyme phenol oxidase in a freshwater monogenetic trematode. A. BHAGAVATHIAMMAI and K. RAMALINGAM	201-209
Anthelmintic activity of triclabendazole. (Research Note). G. C. COLES	210-212
Experimental infection of the baboon ( <i>Papio cynocephalus</i> ) with <i>Echinococcus granulosus</i> of camel, cattle, sheep and goat origin from Kenya. C. N. L. MACPHERSON, J. E. ELSE and M. SULEMAN	213-217
Immunity to tapeworms: acquired resistance to <i>Hymenolepis citelli</i> in the mouse. S. T. O. ALGHALI	219-233
Phenol oxidase activity of a monogenean, <i>Paramazocraes thrissocles</i> . T. THANGARAJ, A. VINAYAKAM and K. NELLAIAPPAN	234-238
Detection of circulating immune complexes in the sera of dogs infected with <i>Dirofilaria immitis</i> by Clq-binding enzyme-linked immunosorbent assay. K. MATSUMURA, Y. KAZUTA, R. ENDO, K. TANAKA and T. INOUE	239-243
The comparative efficacy of oxfendazole administered as bolus and suspension to naturally infected sheep in Greece. C. A. HIMONAS and J. THEODORIDES	245-249
The ability of single sex infections of <i>Schistosoma japonicum</i> to induce resistance to reinfection in mice. N. A. MOLONEY, P. HINCHCLIFFE and G. WEBBE	250-254
Announcements	200, 218, 244