

O T I T I S M E D I A



**DECONGESTION OSMOSIS
WITH REMEDIAL ANALGESIA**

DECONGESTION SUCCESSFULLY ACHIEVED by the addition of Ephedrine Sulphate which acting in synergy with the other ingredients produces shrinkage of the mucosa, promotes drainage from the middle ear and a rapid control of pain. The bactericidal constituents of Auralgicin cover a wide range of micro-organisms including those likely to be present in otitis media, thus avoiding the danger of masking.

AURALGICIN D.D.A.
(BENGER)


Each ml. contains

Phenazonum	0.050 g.	Papaveretum	0.025 g.
EPHEDRINE SULPH.	0.01 g.	Chlorbutol	0.010 g.
Pot. Hydroxyquinolin Sulph.			0.001
Glycer ad 1 ml.			

FOR EXTERNAL APPLICATION

BENGER'S LTD., HOLMES CHAPEL, CHESHIRE

Please mention *The Journal of Laryngology* when replying to advertisements



Instruments used in the practice of Laryngology, Rhinology
and Otology

Mayer & Phelps

MANUFACTURERS
of Surgical Instruments
and Aseptic Hospital
Furniture

Established over
Eighty Years

CHIRON HOUSE

32-34 New Cavendish St., London, W.1

DOWN BROS. and MAYER & PHELPS have amalgamated
The personal managements remain as heretofore

DOWN BROS. and MAYER & PHELPS LTD.