

This is a “preproof” accepted article for Weed Science. This version may be subject to change in the production process, and does not include access to supplementary material.

DOI: 10.1017/wet.2024.103

## **Weed Management in Early Planted Soybean Early Planted Soybean Weed Management as Effected by Herbicide Application Rate and Timing**

Logan R. Miller<sup>1</sup>, Christopher A. Landau<sup>2</sup>, Martin M. Williams II<sup>3</sup>, and Aaron G Hager<sup>4</sup>

<sup>1</sup>Graduate Research Assistant (0009-0005-7469-955X), Department of Crop Sciences, University of Illinois, IL, USA; <sup>2</sup>Postdoctoral Research Agronomist, United States Department of Agriculture-Agricultural Research Service (USDA-ARS), Urbana, IL, USA; <sup>3</sup>Research Ecologist, USDA-ARS, Urbana, IL, USA; <sup>4</sup>Professor, Department of Crop Sciences, University of Illinois, IL, USA

### **Author for correspondence:**

Aaron G. Hager, Professor, Department of Crop Sciences, University of Illinois, N-321 Turner Hall, 1102 S. Goodwin Ave, Urbana, IL 61801. (Email: hager@illinois.edu)

This is an Open Access article, distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives licence (<http://creativecommons.org/licenses/by-nc-nd/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is unaltered and is properly cited. The written permission of Cambridge University Press must be obtained for commercial re-use or in order to create a derivative work.

## Abstract

The opportunity to increase soybean yield has prompted Illinois farmers to plant soybean earlier than historical norms. Extending the growing season with an earlier planting date might alter the relationship between soybean growth and weed emergence timings, potentially altering the optimal herbicide application timings to minimize crop yield loss due to weed interference and ensure minimal weed seed production. The objective of this research was to examine various herbicide treatments applied at different timings and rates to assess the effect on weed control and yield in early planted soybean. Field experiments were conducted in 2021 at three locations across central Illinois to determine effective chemical strategies for weed management in early planted soybean. Preemergence (PRE) treatments consisted of a *S*-metolachlor plus metribuzin premix applied at planting or just prior to soybean emergence at 1/2x (883 + 210 g ai ha<sup>-1</sup>) or 1x (1,766 + 420 g ai ha<sup>-1</sup>) label recommended rates. Postemergence (POST) treatments were applied when weeds reached 10 cm tall and consisted of 1x rates of glufosinate (655 g ai ha<sup>-1</sup>) plus glyphosate (1,260 g ae ha<sup>-1</sup>) plus ammonium sulfate, with or without pyroxasulfone at a 1/2x (63 g ai ha<sup>-1</sup>) or 1x (126 g ai ha<sup>-1</sup>) rate. Treatments comprised of both a full rate of PRE followed by (fb) a POST resulted in the greatest and most consistent weed control at the final evaluation timing. The addition of pyroxasulfone to POST treatments did not consistently reduce late-season weed emergence. The lack of a consistent effect by pyroxasulfone could be attributed to suppression of weeds by soybean canopy closure due to earlier soybean development. The full rate of PRE extended the timing of POST application 2 to 3 weeks for all treatments at all locations except Urbana. Full-rate PRE treatments also reduced the time between the POST application and soybean canopy closure. Overall, a full rate PRE reduced early season weed interference and minimized soybean yield loss due to weed interference.

**Nomenclature:** Glufosinate; glyphosate; metribuzin; pyroxasulfone; *S*-metolachlor; soybean; *Glycine max* (L.) Merr.

**Keywords:** soil-residual herbicide; soybean weed suppression

## Introduction

Improvements in soybean genetics, seed treatments, and planting technology and equipment have all combined to increase soybean yield and profitability over the last several decades. Concomitant with these advances has been a shift to earlier soybean planting. Early soybean planting has become an increasingly common practice with farmers across central Illinois and the US Midwest (USDA ESMIS 2024). The reason for earlier planting is to increase soybean growth prior to the summer solstice which can lead to increased yield (Wilcox and Frankenberger, 1987). Illinois, the leading soybean producing state, alone accounted for 15.8% (4.53 million hectares) of soybean planted in the US in 2022 (USDA NASS 2022, 2023). Considering the economic value and dominance of soybean as a cash crop in Illinois, evaluating weed management in an early-planted soybean environment is prudent.

There are concerns with planting soybean early, such as inadequate crop stands (Oplinger and Philbrook 1992) and increased disease incidence (Hamman et al. 2002). Weed control is another concern, and there are insufficient data to formulate recommendations for managing weeds in early-planted soybean despite extensive research on weed control practices in soybean (especially chemical options).

Preemergence (PRE) herbicides are valuable components of an integrated weed management program. PRE herbicides reduce early season weed interference and often extend the time available to control weeds later in the growing season with a postemergence (POST) herbicide (Corrigan and Harvey 2000). The commercialization of glyphosate-resistant soybean in 1996 substantially reduced use of PRE herbicides (Shaner 2000). This greatly increased selection pressure on the weed communities with POST herbicides (including glyphosate), which led to the evolution of glyphosate resistance in weeds (Duke 2018). As a result, utilization of PRE herbicides in soybean has regained popularity to manage widespread resistance to many POST soybean herbicides.

In central Illinois, early soybean planting is generally considered to begin the first week of April, while historically farmers waited to plant soybean until late April and May. Prior to 2020, Illinois farmers on average planted less than 10% of the soybean crop by April 30, whereas the average hectares planted early from 2020 to 2024 was 28% (USDA ESMIS 2024). Weed control is crucial early in the growing season as soybean are vulnerable to yield loss from weed interference (Cowan et al. 1998; Van Acker et al. 1993). Waterhemp [*Amaranthus tuberculatus*

(Moq.) Sauer], a summer annual weed common throughout the US soybean growing regions, can reduce soybean yield up to 43% (Hager et al. 2002). However, the relative timing of crop and weed emergence may change with planting date, which might necessitate adjustment of herbicide application timing for early-planted soybean. Furthermore, the weed community might change with early-planted soybean, with increased prevalence of early emerging summer annual species such as common lambsquarters and giant ragweed (Werle et al. 2014).

Including herbicides with soil-residual activity with the POST herbicide can extend control of later emerging weed species such as waterhemp, thereby reducing soil seedbank replenishment and reducing selection pressure on future herbicide applications (Gonzini et al 1999; Koger et al 2007). The concept of integrating a split PRE application in soybean may provide enhanced crop safety and extend residual weed control.

Due to lack of data in early-planted soybean, questions regarding the necessity of PRE and/or POST herbicides, along with questions about application rates, persist. The objectives of this research were to: 1) evaluate the need for PRE and POST herbicides in early-planted soybean; and 2) determine the appropriate application rates and timings for PRE and POST herbicides. The knowledge gained will allow weed management practitioners to formulate research-based weed management recommendations in early-planted soybean.

## **Materials and Methods**

### ***Site Selection***

Field experiments were conducted in 2021 at three locations in central Illinois (Urbana: 40°04'43.5"N 88°13'34.0"W; Seymour: 40°02'15.8"N 88°23'36.6"W; Athens: 39°56'41.3"N 89°43'19.8"W). The field locations were selected from our cooperators' willingness to allow us to conduct research at each location. Access to available land and planting equipment were key factors in selection of locations. We selected multiple locations to reduce the risk of adverse weather and/or soil conditions that would preclude establishing experiments according to our objective. Additionally, each location was selected to ensure adequate weed pressure, but individual weed species present at each site was not a criteria of location selection.

The soils at Urbana and Seymour are a Drummer silty clay loam (fine-silty, mixed, superactive, mesic Typic Endoaquolls) with 5.5% organic matter and 6.7 pH. Athens soils included an Ipava silt loam (fine, smectitic, mesic Aquic Argiudolls) and a Clarksdale silt loam (fine,

smectitic, mesic Udollic Endoaqualfs). The Ipava silt loam had a 5.8 pH with 4.3% organic matter, whereas the Clarksdale silt loam had a 6.5 pH with 2.5% organic matter.

### ***General Field Methods***

Experiments in 2021 were initiated at Urbana on April 5, and at Athens and Seymour on April 6. Trials were established following secondary tillage. Either Xtendflex (dicamba, glufosinate, and glyphosate-resistant) soybean (Asgrow® 33XF1, Bayer Crop Science, USA 800 N Lindbergh Blvd, Creve Coeur, MO 63141, USA), (GH3442XF® Syngenta, USA, 410 South Swing Road Greensboro, NC 27409, USA), or E3 (2,4-D, glufosinate, and glyphosate-resistant) soybean (XO3341E®, BASF, USA, 100 Park Avenue, Florham Park, NJ 07932, USA), (GH3442XF® Syngenta, USA, 410 South Swing Road Greensboro, NC 27409, USA) was planted in rows spaced 76 cm apart at a seeding rate of 345,947 seeds ha<sup>-1</sup> at all locations. Monthly precipitation totals for each location are presented in Table 1. Precipitation within 21 days after planting was 6, 3, and 3 cm at Athens, Seymour, and Urbana, respectively.

The experiment was arranged in a randomized complete block with four replications of plots measuring 3 m by 9 m. The treatment design was a five by four factorial of PRE and POST treatments. Treatment structure for each site included 1/2x or 1x PRE-only, POST-only, and 1/2x or 1x PRE followed by (fb) POST (Table 2). PRE treatments included a premix of S-metolachlor + metribuzin (Boundary; Syngenta, Greensboro, NC) applied at 0x, 1/2x, or 1x label recommended rates either at planting, or approximately two weeks after planting and prior to soybean emergence. The POST treatments were applied when weeds reached 10 cm in height and included glyphosate (1260 g ae ha<sup>-1</sup> Roundup PowerMax®, Bayer Crop Science, St. Louis, MO) + glufosinate (655 g ai ha<sup>-1</sup> Liberty, BASF, Research Triangle Park, NC) + liquid ammonium sulfate (Amsol; Winfield Solutions, St. Paul, MN) added at 3.4 kg ha<sup>-1</sup> alone or with pyroxasulfone (63 g ai ha<sup>-1</sup> or 126 g ai ha<sup>-1</sup>, Zidua, BASF, Research Triangle Park, NC). The rationale for including pyroxasulfone was to assess the benefit of extended residual weed control later into the growing season relative to glyphosate + glufosinate alone. Dates of PRE and POST applications are presented in Table 2. All treatments, including application rates and timings, are presented in Table 3.

The premix of S-metolachlor + metribuzin was chosen as the PRE treatment due to the general lack of soybean injury and broad-spectrum weed control. A mixture of glyphosate + glufosinate was selected for POST treatments since both herbicides are non-selective and with no soil-residual activity. Moreover, volatility concerns of glyphosate and glufosinate are negligible

compared to other POST herbicides in herbicide-resistant soybean (Duke and Powles 2008; Takano and Dayan 2020). Although glufosinate demonstrates minimal translocation, and efficacy is often environmentally dependent, resistant weed species are few (Heap 2024).

Herbicides were applied with a CO<sub>2</sub>-pressurized backpack sprayer equipped with AI 110025VS nozzles for PRE, and AIXR 110025 Teejet Air Induction XR nozzles (TeeJet Technologies, 200 W. North Avenue, Glendale Heights, IL 60139, USA) for POST applications. Nozzles were spaced 50 cm apart and calibrated to deliver 187 L/ha at 5.6 km h<sup>-1</sup> and 248 kPa.

### ***Data Collection***

Data collection included days until weed emergence in nontreated plots and all PRE treatments, days to crop emergence, and days to 10 cm tall weeds. Weed species were combined for analysis since the scope of this project was not to evaluate control of any individual species, but rather to evaluate the overall concept of weed control in an early planted soybean environment. Visual evaluation of weed control and soybean injury were made on a scale ranging from 0% (no control or injury) to 100% (complete control) compared with the nontreated beginning at the POST application timing and again 14 and 28 days after each POST application (DAPO). A late-season visual assessment was also made 49 days after the final POST application (DAFPO). Weed density (plants m<sup>-2</sup>) and biomass (g m<sup>-2</sup>) were recorded from two 0.25 m<sup>2</sup> quadrats per plot at the POST application timing and again at the 28 DAPO weed control assessments. Each plot's two biomass samples were combined prior to drying at 65°C, and dry biomass was recorded. Soybean grain yield was determined at maturity using an ALMACO SPC40 combine with a 76-cm row head (ALMACO, Nevada, IA) by harvesting the center two rows of each plot. Final yields were adjusted to 13% moisture.

### ***Statistical analysis***

Weed control (at POST, 14 DAPO, 28 DAPO, and 49 DAFPO), weed biomass (at POST and 28 DAPO), weed density (at POST and 28 DAPO), days to 10 cm tall weeds, and soybean yield were analyzed separately as linear mixed effect models using the LME4 package in R (Bates et al. 2014). PRE and POST treatment, as well as their interactions, were treated as fixed effects, while location and replication were treated as random effects in the models. Mean comparisons were made using Tukey's honestly significant difference test at  $\alpha = 0.05$  with degrees of freedom calculated according to the Kenward-Roger method. Response variables demonstrating a significant PRE by POST interaction in Table 4 were included in Table 5 to compare all combinations of PRE and POST treatments.

## Results and Discussion

### *Soybean Injury*

Soybean injury did not exceed 5% for any PRE treatment regardless of application rate or timing (data not presented). Soybean injury from all POST treatments was 10% or less at seven DAPO and declined over time.

### *Weed control*

Overall, the full rate of PRE extended the timing of the POST application by seven days compared to the half rate of PRE, and 14 days compared to no PRE (Table 4). PRE herbicides are a valuable tool for delaying weed emergence and limiting weed interference with soybean (Knezevic et al. 2019).

Weed species rated across all sites included: velvetleaf (*Abutilon theophrasti* Medik), Palmer amaranth (*Amaranthus palmeri* S. Watson), waterhemp, common lambsquarters (*Chenopodium album* L.), large crabgrass (*Digitaria sanguinalis* (L) Scop.), ivyleaf morningglory (*Ipomoea hederacea* Jacq.), fall panicum (*Panicum dichotomiflorum* Michx.), giant foxtail (*Setaria faberi* Herrm) and common cocklebur (*Xanthium strumarium* L.). Weed control at the initial POST application was influenced by rate of the PRE herbicides whether applied at planting or delayed. Weed control with the full rate of PRE was at least 95% regardless of application timing, while control with the ½ rate of PRE was 88 to 91% (Table 4). In comparison, Ellis and Griffin (2002) observed no difference in weed control when using a half or full rate of pendimethalin + imazaquin, pendimethalin, metolachlor, dimethenamid + imazaquin, sulfentrazone + chlorimuron, and metribuzin + chlorimuron.

POST glyphosate + glufosinate was selected to control all weeds that had emerged through the PRE herbicide, thereby allowing the evaluation of any potential benefit of adding a soil-residual herbicide (pyroxasulfone) with the POST for control of later-emerging weeds. By 14 DAPO, control of all emerged weeds was at least 93% for treatments including a PRE (Table 4). Weed control from PRE-only treatments ranged from 83–89% across application rates and timings 14 DAPO (Table 5). In contrast, weed control with any PRE treatment followed by POST with or without pyroxasulfone ranged from 93–98% 14 DAPO. Incomplete weed control (90–93%) was observed in POST-only treatments with and without pyroxasulfone 14 DAPO. This would have occurred for two reasons. Weed density in POST-only treatments would have made it difficult to achieve adequate coverage with glufosinate, which is crucial for it to control weeds (Knoche M

1994). Control of morningglory and waterhemp with glyphosate would have been insufficient alone. Secondly, species such as velvetleaf, cocklebur, and morningglory emerged within 14 days after the POST application regardless of the inclusion of pyroxasulfone. A POST application too early could allow later emerging weed seedlings to contribute to the soil seedbank. Waterhemp did as well, and this is expected as it emerges in multiple flushes throughout the growing season, especially after a rainfall event (Hartzler et al. 1999).

By 28 DAPO, there were no differences in weed control among treatments regardless of PRE rate or timing (Table 4). Weed control ranged from 96 to 98%. At 28 DAPO, there was no improvement in weed control by including pyroxasulfone with the POST treatment. In contrast, Grey et al. (2013) reported improved weed control by including pyroxasulfone with the POST application of glyphosate + fomesafen. Weed control from treatments not receiving a POST was less compared with treatments with a POST (Table 4).

At 49 DAPO, neither PRE rate nor timing resulted in a difference in weed control among treatments; weed control ranged from 92–95%. PRE fb POST treatments provided 92–97% weed control. Control with POST-only treatments was 94–97%, similar to PRE + POST treatments (Table 4). Weed control was less variable when herbicide treatments included both a PRE fb POST, although POST-only treatments provided similar levels of control.

### ***Weed density and biomass***

There were no differences among PRE rate or timing on weed density or weed biomass at the first POST. At 28 DAPO, weed densities in PRE-only treatments ranged from 21–41 weeds m<sup>-2</sup>, while POST-only treatments ranged from 14–26 weeds m<sup>-2</sup> (Table 5). Weed density was lower 28 DAPO for PRE fb POST treatments relative to POST-only treatments, yet no statistical differences were apparent. Including pyroxasulfone with the POST did result in lower weed densities but not significantly different compared to POST treatments without pyroxasulfone. Sarangi and Jhala (2019) did find a difference in Palmer Amaranth density when they collectively analyzed POST vs POST with residual 28 DAPO, however, velvetleaf density was not different. When broken down by each treatment to evaluate Palmer Amaranth density 28 DAPO, chloransulam-methyl + pyroxasulfone/fluthiacet-methyl was the only POST treatment with a residual herbicide to display differences, while all individual treatments had no effect on velvetleaf density.

Weed biomass was comparable at the first POST regardless of PRE rate or timing, showing no difference and ranging from 2 to 3 g m<sup>-2</sup>. At 28 DAPO, weed biomass of PRE-only treatments ranged from 27 to 51 g m<sup>-2</sup>. Weed biomass of POST-only treatments were similar as well and ranged from 5 to 11 g m<sup>-2</sup>. PRE fb POST treatments resulted in weed biomass of 1 to 6 g m<sup>-2</sup> 28 DAPO (Table 5).

### ***Soybean canopy closure/weed emergence***

Despite the variability in soybean canopy closure timing at each site, other than Urbana the only weeds noted to contribute to the weed seedbank were POST escapes in PRE fb POST treatments. Ivyleaf morningglory and common cocklebur were two weed species that emerged after the POST application and were not suppressed by the canopy in Urbana in 2021. Weed emergence was observed in treatments not receiving pyroxasulfone in the POST application at the other sites, but where common cocklebur was not present, these other weeds were suppressed by soybean canopy. Common cocklebur has shown the ability to tolerate reduced light levels under shaded conditions, which may explain why it was not suppressed by soybean (Regnier and Stoller 1989).

Early-planted soybean can achieve row closure sooner than later planted soybean. Later emerging weeds likely would be suppressed or have higher mortality rates in an early planted soybean environment (Arsenijevic et al. 2022). Velvetleaf emerging later in the season experienced higher mortality levels when under a soybean canopy (Lindquist et al. 1995). However, later emerging waterhemp has shown the ability to produce seed under shaded conditions in a standard soybean planting timing (Hartzler et al. 2004).

### ***Soybean yield***

Soybean yield for POST-only and PRE fb POST treatments was similar (Table 4). Soybean yield was greater for treatments receiving a POST compared to treatments without a POST. Soybean yield has been similar between reduced and full labeled rates of PRE herbicides (Muyonga et al. 1996). Soybean yield may be most affected by the timing of weed emergence, with earlier emerging weeds posing the greatest threat to yield loss (Kropff et al. 1992). PRE herbicides minimize the duration of weed competition with the crop when it is most vulnerable. External stresses during seed fill reduce soybean yield, which PRE-only treatments would allow for given greater weed interference during this reproductive period (Foroud et al. 1993).

## **Practical Implications**

Applying a full rate of S-metolachlor + metribuzin extended the timing of the glyphosate + glufosinate application compared to .5x rate or no PRE, although delaying S-metolachlor + metribuzin closer to soybean emergence offered no advantage in weed control or extending days to the POST application. Both rates of pyroxasulfone when included with glyphosate + glufosinate did not significantly reduce weed densities 28 DAPO. This may be explained by earlier soybean development in relation to weed emergence. A more developed soybean canopy would reduce the fluctuation of soil surface temperature and incident sunlight earlier in the season, reducing weed seedling emergence (Norsworthy and Oliveira 2007). PRE fb POST treatments provided the highest level of weed control and soybean yield. The PRE-only treatments did not yield as high as the POST-only treatments, which were similar to PRE fb POST treatments. Klingaman and Oliver (1994) reported increased competitiveness of soybean with entireleaf morningglory (*Ipomoea hederacea* var. *integriuscula*) and sicklepod (*Senna obtusifolia*) when planted in early May compared with early June. Planting soybean earlier should improve suppression of later-emerging weed species, yet this environment may be more conducive for earlier emerging weed species (Werle et al. 2014), and soil disturbance may promote summer annual species to shift to earlier emergence. Current and future research on early-planted soybean in comparison to conventional soybean planting timing includes injury potential from various soil-residual herbicides, herbicide carryover potential, and POST timing efficacy with and without a soil-residual herbicide.

## **Acknowledgments**

We thank the staff and undergraduate students in the University of Illinois Herbicide Evaluation program for their invaluable assistance in conducting this research.

## **Funding**

This research was funded by the Illinois Soybean Association.

## **Competing Interests**

No competing interests have been declared.

## References

- Arsenijevic N, DeWerff R, Conley S, Ruark M, Werle R (2022) Influence of integrated agronomic and weed management practices on soybean canopy development and yield. *Weed Technol* 36:73–78
- Bates D, Mächler M, Bolker B, Walker S (2014) Fitting linear mixed-effects models using lme4. R package version 1.1–7
- Corrigan KA, Harvey, RG (2000) Glyphosate with and without residual herbicides in no-till glyphosate-resistant soybean (*Glycine max*). *Weed Technol* 14:569–577
- Cowan P, Weaver SE, Swanton CJ (1998) Interference between pigweed (*Amaranthus spp.*), barnyardgrass (*Echinochloa crus-galli*), and soybean (*Glycine max*). *Weed Sci* 46:533–539
- Duke SO (2018) The history and current status of glyphosate. *Pest Manag Sci* 74:1027–1034
- Duke SO, Powles SB (2008) Glyphosate: a once-in-a-century herbicide. *Pest Manag Sci* 64:319–325
- Ellis JM, Griffin JL (2002) Benefits of soil-applied herbicides in glyphosate-resistant soybean (*Glycine max*). *Weed Technol* 16:541–547
- Foroud N, Mündel HH, Saindon G, Entz T (1993) Effect of level and timing of moisture stress on soybean plant development and yield components. *Irrigation Sci* 13:149–155
- Gonzini LC, Hart SE, Wax LM (1999) Herbicide combinations for weed management in glyphosate-resistant soybean (*Glycine max*). *Weed Technol* 13:354–360
- Grey TL, Cutts III GS, Newsome LJ, Newell III SH (2013) Comparison of pyroxasulfone to soil residual herbicides for glyphosate resistant Palmer amaranth control in glyphosate resistant soybean. *Crop Manag* 12:1–6
- Hager, A. G., Wax, L. M., Stoller, E. W., & Bollero, G. A. (2002) Common waterhemp (*Amaranthus rudis*) interference in soybean. *Weed Sci* 50:607–610
- Hamman B, Egli DB, Koning G (2002) Seed vigor, soilborne pathogens, preemergent growth, and soybean seedling emergence. *Crop Sci* 42:451–457
- Hartzler RG, Battles BA, Nordby D (2004) Effect of common waterhemp (*Amaranthus rudis*) emergence date on growth and fecundity in soybean. *Weed Sci* 52:242–245

- Hartzler RG, Buhler DD, Stoltenberg DE (1999) Emergence characteristics of four annual weed species. *Weed Sci* 47:578–584
- Heap I (2024) The International Herbicide-Resistant Weed Database. <https://www.weedscience.org/Pages/Species.aspx>. Accessed: November 6, 2024
- Klingaman TE, Oliver LR (1994) Influence of cotton (*Gossypium hirsutum*) and soybean (*Glycine max*) planting date on weed interference. *Weed Sci* 42:61–65
- Knezevic SZ, Pavlovic P, Osipitan OA, Barnes ER, Beiermann C, Oliveira MC, Lawrence N, Scott JE, Jhala A (2019) Critical time for weed removal in glyphosate-resistant soybean as influenced by preemergence herbicides. *Weed Technol* 33:393–399
- Knoche M (1994) Effect of droplet size and carrier volume on performance of foliage-applied herbicides. *Crop Prot* 13:163–178
- Koger CH, Price AJ, Faircloth JC, Wilcut JW, Nichols SP (2007) Effect of residual herbicides used in the last post-directed application on weed control and cotton yield in glyphosate- and glufosinate-resistant cotton. *Weed Technol* 21:378–383
- Kropff MJ, Weaver SE, Smits MA (1992) Use of ecophysiological models for crop-weed interference: relations amongst weed density, relative time of weed emergence, relative leaf area, and yield loss. *Weed Sci* 40:296–301
- Lindquist JL, Maxwell BD, Buhler DD, Gunsolus JL (1995) Velvetleaf (*Abutilon theophrasti*) recruitment, survival, seed production, and interference in soybean (*Glycine max*). *Weed Sci* 43:226–232
- Muyonga KC, Defelice MS, Sims BD (1996) Weed control with reduced rates of four soil applied soybean herbicides. *Weed Sci* 44:148–155
- Norsworthy JK, Oliveira MJ (2007) A model for predicting common cocklebur (*Xanthium strumarium*) emergence in soybean. *Weed Sci* 55:341–345
- Oplinger ES, Philbrook BD (1992) Soybean planting date, row width, and seeding rate response in three tillage systems. *J Prod Agric* 5:94–99
- Priess GL, Norsworthy JK, Roberts TL, Gbur EE (2020) Weed control and soybean injury from preplant vs. preemergence herbicide applications. *Weed Technol* 34:718–726
- Regnier EE, Stoller EW (1989) The effects of soybean (*Glycine max*) interference on the canopy architecture of common cocklebur (*Xanthium strumarium*), jimsonweed (*Datura stramonium*), and velvetleaf (*Abutilon theophrasti*). *Weed Sci* 37:187–195

- Sarangi D, Jhala AJ (2019) Palmer amaranth (*Amaranthus palmeri*) and velvetleaf (*Abutilon theophrasti*) control in no-tillage conventional (non-genetically engineered) soybean using overlapping residual herbicide programs. *Weed Technol* 33:95–105
- Shaner DL (2000) The impact of glyphosate-tolerant crops on the use of other herbicides and on resistance management. *Pest Manag Sci* 56:320–326
- Takano HK, Dayan FE (2020) Glufosinate-ammonium: a review of the current state of knowledge. *Pest Manag Sci* 76:3911–3925
- [USDA ESMIS] U.S. Department of Agriculture Economics, Statistics and Market Information System (2024) [usda.library.cornell.edu/concern/publications/8336h188j?locale=en&page=14#release-items](https://usda.library.cornell.edu/concern/publications/8336h188j?locale=en&page=14#release-items). Accessed: November 8, 2023
- [USDA NASS] U.S. Department of Agriculture–National Agriculture Statistics Service (2022) [https://www.nass.usda.gov/Quick\\_Stats/Ag\\_Overview/stateOverview.php?state=ILLINOIS](https://www.nass.usda.gov/Quick_Stats/Ag_Overview/stateOverview.php?state=ILLINOIS). Accessed: April 15, 2023
- [USDA NASS] U.S. Department of Agriculture–National Agriculture Statistics Service (2023) <https://www.nass.usda.gov/Newsroom/archive/2023/01-12-2023.php>. Accessed: April 20, 2023
- Van Acker RC, Swanton CJ, Weise SF (1993) The critical period of weed control in soybean [*Glycine max* (L.) Merr.]. *Weed Sci* 41:194–200
- Werle R, Sandell LD, Buhler DD, Hartzler RG, Lindquist JL (2014) Predicting emergence of 23 summer annual weed species. *Weed Sci* 62:267–279
- Wilcox JR, Frankenberger EM. (1987) Indeterminate and determinate soybean responses to planting date. *Agron J* 79:1074–1078

Table 1. Monthly total precipitation at Athens, Seymour, and Urbana, Illinois in 2021.

Month	Athens	Seymour	Urbana
	----- cm -----		
	-----		
April	7	5	5
May	14	9	8
June	12	17	17
July	12	10	9
August	13	6	7
Total	58	47	46

Table 2. Herbicide application dates in early planted soybean trials at Athens, Seymour, and Urbana, Illinois in 2021. Postemergence herbicide applications were made when weeds were 10 cm tall.

Application timing	Athens	Seymour	Urbana
PRE <sup>a</sup>	Apr 6	Apr 7	Apr 5
Delayed PRE	Apr 16	Apr 16	Apr 16
1st POST <sup>b</sup>	Jun 3	May 27	May 27
2nd POST <sup>c</sup>	Jun 10	Jun 11	Jun 4
3rd POST <sup>d</sup>	Jun 17	Jun 17	–

<sup>a</sup>PRE application the day of soybean planting

<sup>b</sup>Treatments at 1<sup>st</sup> POST that received herbicide application were POST-only across all sites, along with half rate of PRE (HPRE) at Urbana

<sup>c</sup>Treatments at 2<sup>nd</sup> POST that received herbicide application were HPRE at Athens and Seymour, and full rate of PRE (FPRE) at Urbana

<sup>d</sup>Treatments at 3<sup>rd</sup> POST that received herbicide application were FPRE at Athens and Seymour

Table 3. Herbicide treatments applied in early planted soybean trials at Athens, Seymour, and Urbana, Illinois in 2021.

Treatment	PRE	Rate <sup>a</sup> g ai ha <sup>-1</sup>	Timing <sup>b</sup>	POST	Rate g ai ha <sup>-1</sup>
1	Nontreated Control (No PRE or POST)	–	–	–	–
2	S- metolachlor + metribuzin	1,766 + 420	at planting	–	–
3	S- metolachlor + metribuzin	1,766 + 420	at planting	glyphosate + glufosinate	1260 + 655
4	S- metolachlor + metribuzin	1,766 + 420	at planting	glyphosate + glufosinate + pyroxasulfon e	1260 + 655 + 63
5	S- metolachlor + metribuzin	1,766 + 420	at planting	glyphosate + glufosinate + pyroxasulfon e	1260 + 655 + 126
6	S- metolachlor + metribuzin	883 + 210	at planting	–	–
7	S- metolachlor + metribuzin	883 + 210	at planting	glyphosate + glufosinate	1260 + 655
8	S- metolachlor + metribuzin	883 + 210	at planting	glyphosate + glufosinate + pyroxasulfon e	1260 + 655 + 63
9	S- metolachlor + metribuzin	883 + 210	at planting	glyphosate + glufosinate + pyroxasulfon e	1260 + 655 + 126
10	S- metolachlor + metribuzin	1,766 + 420	2WAP	–	–

<sup>a</sup>Rate for glyphosate expressed as g ae ha<sup>-1</sup>

<sup>b</sup>2WAP= two weeks after planting

Table 3. Continued

Treatment	PRE	Rate	Timing	POST	Rate
11	S-metolachlor + metribuzin	1,766 + 420	2WAP	glyphosate glufosinate	+ + 1260 + 655
12	S-metolachlor + metribuzin	1,766 + 420	2WAP	glyphosate glufosinate pyroxasulfone	+ + + 1260 + 655 + 63
13	S-metolachlor + metribuzin	1,766 + 420	2WAP	glyphosate glufosinate pyroxulfone	+ + + 1260 + 655 + 126
14	S-metolachlor + metribuzin	883 + 210	2WAP	–	–
15	S-metolachlor + metribuzin	883 + 210	2WAP	glyphosate glufosinate	+ + 1260 + 655
16	S-metolachlor + metribuzin	883 + 210	2WAP	glyphosate glufosinate pyroxasulfone	+ + + 1260 + 655 + 63
17	S-metolachlor + metribuzin	883 + 210	2WAP	glyphosate glufosinate pyroxasulfone	+ + + 1260 + 655 + 126
18	–	–	–	glyphosate glufosinate	+ + 1260 + 655
19	–	–	–	glyphosate glufosinate pyroxasulfone	+ + + 1260 + 655 + 63
20	–	–	–	glyphosate glufosinate pyroxasulfone	+ + + 1260 + 655 + 126

Table 4. Summary of main effects and interactions for weed response and soybean yield in early planted soybean trials across three locations in Illinois in 2021<sup>a</sup>.

Main effect	Days after planting until 10 cm tall weeds	Weed control at POST <sup>b</sup>	Weed density at POST	Weed biomass at POST	Weed control 14 DAPO <sup>c</sup>	Weed control 28 DAPO	Weed density 28 DAPO	Weed biomass 28 DAPO	Weed control 49 DAFPO	Soybean yield				
			plants m <sup>-2</sup>	g m <sup>-2</sup>			plants m <sup>-2</sup>	g m <sup>-2</sup>		kg ha <sup>-1</sup>				
PRE treatment	*	*	*	*	*	*	*	*	*	ns				
No PRE	54	c	0	d	68	a	12.4	a	67	95	36	43.6	95	4,828
1x S-metolachlor + metribuzin at planting	68	a	96	a	20	b	3.0	b	94	98	12	7.5	95	4,801
0.5x S-metolachlor + metribuzin at planting	61	b	91	bc	19	b	2.0	b	93	96	15	13.3	93	4,613
1x S-metolachlor + metribuzin 2WAP	68	a	95	ab	20	b	2.4	b	95	97	13	8.1	94	5,030
0.5x S-metolachlor + metribuzin 2WAP	61	b	88	c	20	b	1.9	b	94	96	21	16.1	92	4,983

Table 4. Continued

POST Timing	NA	NA	NA	NA	*	*	*	*	*	*			
No POST	.	.	.	.	69	90	b	39	56	84	b	4,257	b
Glyphosate + glufosinate	.	.	.	.	95	97	a	21	4.3	94	a	5,010	a
Glyphosate + glufosinate + 0.5x pyroxasulfone	.	.	.	.	95	97	a	10	3.9	97	a	5,010	a
Glyphosate + glufosinate + 1x pyroxasulfone	.	.	.	.	95	98	a	7	3	97	a	4,983	a
Interaction													
PRE*POST	NA	NA	NA	NA	*	ns	*	*	ns	ns			

<sup>a</sup>Values shown are means. Main effect means among PRE or POST treatment within a column with no common letter are significantly different according to Tukey's honest significance test at  $\alpha = 0.05$ . Significant at \* $P < 0.05$ . Not significant = ns.

<sup>b</sup>0 represents no control and 100 represents complete control.

<sup>c</sup>DAPO, days after POST; DAFPO, days after final POST; 2WAP, 2 weeks after planting

Table 5. Weed control, density, and biomass in response to PRE and POST treatment across three locations in Illinois in 2021<sup>a</sup>

PRE treatment	POST treatment							
	No POST		Glyphosate + glufosinate		Glyphosate + glufosinate + 0.5x pyroxasulfone		Glyphosate + glufosinate + 1x pyroxasulfone	
<b>Weed Control (%) 14 DAPO<sup>b</sup></b>								
No PRE	0	c	90	ab	90	ab	92	ab
1x S-metolachlor + metribuzin at planting	88	ab	96	a	96	a	96	a
0.5x S-metolachlor + metribuzin at planting	83	b	96	a	96	a	93	ab
1x S-metolachlor + metribuzin 2WAP	88	ab	96	a	97	a	97	a
0.5x S-metolachlor + metribuzin 2WAP	87	ab	95	a	96	a	96	a
<b>Weed Density (plants m<sup>-2</sup>) 28 DAPO<sup>c</sup></b>								
No PRE	79	a	26	bc	25	bc	14	bc
1x S-metolachlor + metribuzin at planting	24	bc	12	bc	7	bc	6	bc
0.5x S-metolachlor + metribuzin at planting	28	bc	22	bc	6	bc	6	bc
1x S-metolachlor + metribuzin 2WAP	21	bc	23	bc	3	c	4	c
0.5x S-metolachlor + metribuzin 2WAP	41	b	25	bc	8	bc	7	bc
<b>Weed Biomass (g m<sup>-2</sup>) 28 DAPO</b>								
No PRE	148.							
1x S-metolachlor + metribuzin at planting	6	a	10.2	cd	10.6	bcd	4.5	cd
0.5x S-metolachlor + metribuzin at planting	26.7	bcd	0.8	d	1.6	d	0.9	d
1x S-metolachlor + metribuzin 2WAP	42.9	bc	4.6	cd	3.3	cd	2.3	d
0.5x S-metolachlor + metribuzin 2WAP	29.1	b	1.5	d	0.5	d	1.1	d
50.7	a	4.2	cd	3.5	cd	5.8	cd	

<sup>a</sup>Values shown are means. Means among PRE or POST treatment with no common letter are significantly different according to Tukey's honest significance test at  $\alpha = 0.05$ . Significant at \* $P < 0.05$ . Comparisons for each response variable can be made across PRE and POST treatments.

<sup>b</sup>Percent of non-treated, where 0 represents no control and 100 represents complete control.

<sup>c</sup>DAPO, Days after POST