

In conclusion, I would agree with Professor Casey that defensive medicine and kneejerk practice is no substitute for sound clinical judgement. Based on the scientific facts which are now available to the medical profession, I do question the objective basis of what she considers to be sound clinical judgement.

Brian E Leonard, PhD, DSc, MRJA
*Professor of Pharmacology,
Pharmacology Department,
University College Galway,
Co Galway,
Ireland.*

References

1. Leonard BE. Editorial: Litigation and the toxicity of psychotropic drugs. *Ir J Psychol Med* 1996; 13: 88-9
2. Casey P. Commentary: Litigation and the toxicity of psychotropic drugs. *Ir J Psychol Med* 1996; 13: 89-90
3. Carlston A, Allebeck P, Brandt L. Are suicide rates in Sweden associated with changes in the prescribing of medicines? *Acta Psychiatr Scand* 1996; 94: 94-100.
4. Farmer RDT. Suicide and poisons. *Human Psychopharmac* 1994; 9(1): S1-S21.
5. Henry JA, Alexander CA, Sener EK. Relative mortality from overdose of antidepressives. *BMJ* 1995; 310: 221-4.
6. Daly RJ. Medico-legal aspects of prescribing. *Human Psychopharmac* 1994; 9(1): S31-S36.
7. de Gier JJ. Drugs other than alcohol and driving in the European Union. Study coordinated with the support of the European Blood Safety Federation and the Directorate General for Transport of the Commission for the European Communities. EC document 1995; IHP: 95-94.
8. Ray WA, Fought RL, Decker MD. Psychoactive drugs and the risk of injurious motor vehicle crashes in elderly drivers. *Am J Epidemiol* 1992; 16: 873-83.
9. Montgomery SA, Henry JA, McDonald G, *et al*. Selective serotonin re-uptake inhibitors: meta analysis of discontinuation rates. *Int Clin Psychopharmac* 1994; 9: 47-53.

Dysmnestic MCQs

Sir - An experienced neurologist reported great difficulty in finding the correct answers to neurological MCQ questions when these were requested by the first author whilst he was preparing for the College exams. Experienced psychiatrists often have the same experience when attempting to answer psychiatric MCQ questions for their trainees. The basic problem was illustrated when the second author sought guidance with this question¹ - Q(28). "Following a head injury, the following is true: - Anterograde amnesia may persist for longer than post-traumatic amnesia." The answer, "true".

The correct answer is that it is controversial and misleading. The question was derived from Kopelman² - "post-traumatic amnesia (PTA) refers to memory loss for events following a lesion. Anterograde amnesia (AA) refers to an impairment in learning new material. AA, may persist long after the termination of PTA." These definitions are uncomfortably alike for clinical purposes, as Kopelman agrees when he writes: "AA is commonly used synonymously with PTA."²

Varying definitions abound, eg. one source defines PTA in terms of memory and AA in terms of conscious level. The problem would be solved if we had only the following terms;

post-traumatic disturbance of consciousness (PTDC) and PTA, each meaning what they said. Unless terminology reflects what is being observed it is dangerously misleading. How many trainees can remember the difference between hypnagogic and hypnopompic hallucinations before they see the go (- ing to sleep) in hypnagogic?

Brian O'Shea, MB, BCh, FRCPsych,
Consultant Psychiatrist
Rita Condren, MB, BCh,
*Senior House Officer,
Newcastle Hospital,
Greystones,
Co Wicklow,
Ireland.*

References

1. Green B. *The MRCPsych Study Manual*. Dordrecht, The Netherlands: Kluwer Academic Publications, 1993.
2. Kopelman MD. Amnesia: organic and psychogenic. *Br J Psychiatry* 1987; 150: 428-42.
3. Gelder M, Gath D, Xayou R, Cowen P. *Oxford Textbook of Psychiatry*, 3rd ed. Oxford: Oxford University Press, 1996.

Psychiatric inpatient suicide

Sir - Coakley *et al* in their thoughtful study of inpatient suicides¹ recommend two types of investigation following an inpatient suicide; a formal inquiry and a confidential informal review of the event. I have worked in a system where such a practice has been in place for many years and I support their recommendations in this regard.

When a suicide occurs in either the inpatient or outpatient setting a procedure known as a "Psychological Autopsy" is activated. Administrative and clinical staff who have been involved in any way with the patient, as well as care givers from the community are invited to review the deceased patient's contact with our services, the treatment strategies that were used, and make recommendations to prevent a similar tragedy. A record which is kept confidential within the department is kept of the recommendations. This procedure serves as a healing ritual for staff and helps protect staff morale. It helps to deal with the feelings of loss and grief that many staff experience when the tragedy of a suicide occurs.

Dr Patrick G Coll, MB, FRCP (C)
*Clinical Assistant Professor,
Department of Psychiatry,
University of Calgary,
Calgary General Hospital,
841 Centre Ave East,
Calgary, Alberta T2E 0A1,
Canada.*

Reference

1. Coakley GM, Carey TG, Owens JM. A study of psychiatric inpatient suicides. *Ir J Psychol Med* 1996; 13(3): 102-4