

Vitamin D deficiency in minority populations

Glen B Taksler^{1,*}, David M Cutler^{2,3}, Edward Giovannucci⁴ and Nancy L Keating^{5,6}

¹Medicine Institute, Cleveland Clinic, 9500 Euclid Avenue, G1-40F, Cleveland, OH 44195, USA; ²Department of Economics and Kennedy School of Government, Harvard University, Cambridge, MA, USA; ³National Bureau of Economic Research, Cambridge, MA, USA; ⁴Departments of Nutrition and Epidemiology, Harvard School of Public Health, Boston, MA, USA; ⁵Department of Health Care Policy, Harvard Medical School, Boston, MA, USA; ⁶Department of Medicine, Brigham and Women's Hospital, Boston, MA, USA

Submitted 25 November 2013: Final revision received 26 February 2014: Accepted 3 March 2014: First published online 15 April 2014

Abstract

Objective: Black and Hispanic individuals synthesize less vitamin D per unit of sun exposure than white individuals. The relationship between UV radiation and vitamin D insufficiency in minorities has not been well explored.

Design: Prospective cohort study.

Setting: Using the National Health and Nutrition Examination Survey, we obtained serum vitamin D levels for non-Hispanic Whites, Hispanics and non-Hispanic Blacks aged ≥ 18 years from 2000–2006. We linked these data with the average monthly solar UV index by census tract and data on sun exposure, vitamin D supplementation, health and demographics. We used multivariable regression analyses to assess vitamin D deficiency (< 15 ng/ml) and insufficiency (< 20 ng/ml) in January (when the UV index was lowest) by race/ethnicity and geography.

Subjects: Adults (n 14 319) aged ≥ 18 years.

Results: A 1-point increase in the UV index was associated with a 0.51 ng/ml increase in vitamin D (95% CI 0.35, 0.67 ng/ml; $P < 0.001$). Non-Hispanic Black race and Hispanic ethnicity were associated with a 7.47 and 3.41 ng/ml decrease in vitamin D, respectively (both $P < 0.001$). In January, an estimated 65.4% of non-Hispanic Blacks were deficient in vitamin D, compared with 28.9% of Hispanics and 14.0% of non-Hispanic Whites. An estimated 84.2% of non-Hispanic Blacks were insufficient in vitamin D *v.* 56.3% of Hispanics and 34.8% of non-Hispanic Whites. More non-Hispanic Blacks were estimated to be deficient in vitamin D in January in the highest UV index quartile than were non-Hispanic Whites in the lowest UV index quartile (60.2% *v.* 25.7%).

Conclusions: Wintertime vitamin D insufficiency is pervasive among minority populations, and not uncommon among non-Hispanic Whites.

Keywords
Vitamins
Vitamin D
Disparities
Race
Survival

The Institute of Medicine estimates that 54% of non-Hispanic Blacks have very low levels of vitamin D (25-hydroxycholecalciferol (25(OH)D₃) < 16 ng/ml), *v.* 27% of Mexican Americans and 11% of non-Hispanic Whites^(1,2). A growing literature^(1–3) suggests that vitamin D may be associated with reduced mortality (overall⁽⁴⁾ and hospital-associated⁽⁵⁾), incidence of cancer^(6–13), CVD^(14,15), diabetes⁽¹⁶⁾, fractures^(17,18) and ageing-related functional limitations⁽¹⁹⁾, although some studies find an unclear relationship^(20,21) or a harmful association with disease⁽²²⁾. Randomized controlled trials are underway to examine the potential role of vitamin D in the prevention of multiple diseases, including cancer, CVD and diabetes^(20,23).

There are known racial and seasonal variations in vitamin D. Black individuals synthesize less vitamin D per unit of sun exposure than white individuals, because darker complexion acts as a natural sunscreen^(24,25).

The majority of vitamin D is estimated to derive from sunshine for most people⁽²⁶⁾, with the liver converting cholecalciferol into 25(OH)D₃, the form of vitamin D typically measured in blood serum. 25(OH)D₃ has a reported half-life of 15 d⁽²⁷⁾, and therefore decreases from summer (when solar radiation is highest) to winter⁽²⁸⁾. Observational studies suggest higher cancer mortality rates for patients diagnosed in winter^(29,30), at northern latitudes^(31,32) and in areas with less UV radiation from sunshine^(33,34), although genetics may contribute⁽³⁵⁾. Additionally, previous research suggests that racial differences in the synthesis of vitamin D from sunshine are largest in January⁽³⁶⁾. However, studies have not adequately considered the association between the UV index and racial differences in vitamin D deficiency or insufficiency.

We assessed the extent of vitamin D insufficiency by race/ethnicity and geography (as categorized by the

*Corresponding author: Email taksleg@ccf.org

UV index) in January, when low UV radiation limited the body's ability to synthesize vitamin D from sunshine.

Experimental methods

Data and participants

We obtained data on serum vitamin D and race/ethnicity from the National Health and Nutrition Examination Survey (NHANES)⁽³⁷⁾ for 14 319 individuals aged ≥ 18 years from 2000 to 2006. Individuals had known serum 25-hydroxyvitamin D (25(OH)D), race (non-Hispanic White, Hispanic, non-Hispanic Black), age, gender, marital status, household income, smoking status and BMI. We also obtained restricted data on each participant's census tract of residence, month and year of NHANES examination and testing, and serum 25(OH)D levels for 2000 (vitamin D data were publicly available for later years).

For UV irradiance, we used satellite data from the National Aeronautics and Space Administration in increments of 1° latitude and 1.25° longitude, throughout the USA, collected from 1978 to 2005^(38,39). These data covered the UV-B and UV-A spectrum (280–400 nm). We obtained the average noontime UV index ('UV index') for each calendar month, for the centroid of each NHANES respondent's census tract^(40,41). The UV index measures UV radiation reaching the Earth's surface; in the continental USA, it ranges from 0–5 in January to 4–12 in July⁽⁴⁰⁾. The index is a function of latitude, time, solar irradiation, elevation, ozone and cloud cover^(40,41). Although we used the long-term average UV index, the correlation of the long-term average and the 2000–2005 average was $\geq 99.0\%$ for each calendar month.

Control variables

Using NHANES data, we obtained each participant's age, gender, marital status, annual household income, hair colour, skin reaction to half an hour of sun exposure without sunscreen or protective clothing, how often an individual typically stayed in the shade on a sunny day when outside for > 1 h, number of sunburns in the past year, how often an individual wore a hat that covered the face, ears and neck, how often an individual wore a long-sleeved shirt, how often an individual wore sunscreen, the sun protection factor (SPF) most often used, average daily vitamin D supplementation during the past 30 d, BMI, smoking status, and month and year of examination. Table 1 provides categorizations of each variable.

Analyses

To better understand why vitamin D might vary across races and ethnicities, we performed three analyses. First, we estimated the number of minutes outdoors at noon in each calendar month for fair-skinned white, Hispanic and dark-skinned black individuals to synthesize 15 μg (600 IU) of vitamin D, the RDA for adults aged ≤ 70 years⁽¹⁾.

This calculation was defined as the minimum erythral dose (MED) of UV radiation required for individuals of each race to synthesize 15 μg (600 IU) of vitamin D, divided by the MED supplied per minute of noontime outdoor exposure. For the numerator, we followed dermatology literature to expect that fair-skinned white, Hispanic and dark-skinned black individuals (Fitzpatrick skin types I, IV and VI, respectively, where I is the lightest and VI is the darkest skin tone⁽⁴²⁾) would require 0.25, 0.56 and 1.25 MED of UV radiation, respectively, to synthesize 15 μg (600 IU) of vitamin D, with 25.5% of body surface area – the head, neck, arms and hands – uncovered (Appendix 1)⁽⁴³⁾. Again following dermatology literature⁽⁴³⁾, we assumed linearity of this relationship, so that half the MED exposure or uncovered body surface area would yield half the vitamin D absorption. For the denominator, we used the definition of the UV index (1 unit = 0.0072 MED/min)⁽⁴⁴⁾. We also assumed a relationship between average maximum daily temperature and body parts exposed to sunlight ($\geq 23.9^\circ\text{C}$ ($\geq 75^\circ\text{F}$): face, neck, hands, arms exposed; 15.6–23.8°C (60–74°F): face, neck, hands, lower arms exposed; –1.1 to 15.5°C (30–59°F): face, neck, hands exposed; –9.4 to –1.2°C (15–29°F): face, neck exposed; $\leq -9.5^\circ\text{C}$ ($\leq 14^\circ\text{F}$): face exposed)^(36,45,46). We then estimated the difference in minutes required to synthesize 15 μg (600 IU) of vitamin D, between each minority population (Hispanics, dark-skinned blacks) and fair-skinned whites. To better illustrate our methods, consider Atlanta, Georgia in January. Atlanta had an average UV index of 2.54⁽³⁸⁾ and maximum temperature of 11.1°C (52°F)⁽⁴⁵⁾. We estimated that fair-skinned whites would synthesize 15 μg (600 IU) vitamin D in 31 min (0.25 MED required to synthesize 15 μg of vitamin D \div (0.0072 MED/UV index unit \times 2.54 UV index units) \times (25.5% exposed body surface area in baseline formula \div 11.5% of exposed body surface area at 11.1°C))^(36,38,40,41,43). Similarly, we estimated that Hispanics and dark-skinned blacks would synthesize 15 μg (600 IU) vitamin D in 69 and 153 min, respectively^(36,38,49,41,43). Therefore, estimated racial/ethnic differences in the number of minutes to synthesize 15 μg (600 IU) of vitamin D were 41 min for Hispanics and 126 min for dark-skinned blacks, each as compared with fair-skinned whites.

Second, we used multiple regression models to assess the relationship between serum vitamin D and the UV index. The dependent variable was vitamin D (ng/ml) at the time of NHANES examination and the main independent variables of interest were each participant's race/ethnicity and the average UV index 1 month prior to the exam. We selected 1 month prior to the exam because vitamin D has an estimated half-life of 15 d⁽²⁷⁾, and the former was more highly correlated with vitamin D than the average UV index in the month of the exam or 2–3 months prior to the exam. Control variables were as described above. Observations were weighted using NHANES-provided weights for its medical examination sample, and robust standard errors were used. In sensitivity

Table 1 Summary statistics of the study population; adults aged ≥ 18 years ($n = 14\,319$), National Health and Nutrition Examination Survey (NHANES), 2000–2006

	Non-Hispanic White ($n = 7554$)	Non-Hispanic Black ($n = 2991$)	Hispanic ($n = 3774$)	<i>P</i>
Demographics				
Age (years)	47.3	43.0	39.9	< 0.001
Female	51.4	56.0	50.5	
Married	62.9	36.3	57.2	
Annual household income				
< \$US 25 000	19.2	33.3	34.6	< 0.001
\$US 25 000–34 999	10.6	13.6	15.6	
\geq \$US 35 000	65.8	46.1	42.8	
Serum vitamin D				
Vitamin D (ng/ml)	25.9	14.9	20.7	
Category				
Deficient (<15 ng/ml)	8.3	53.4	21.7	< 0.001
Insufficient (<20 ng/ml)	23.3	77.3	46.4	
Alternative insufficient (<30 ng/ml)	70.2	97.0	88.3	
UV index in census tract of residence				
UV index month of exam (level)	5.8	5.8	5.3	
UV index 1 month before exam (level)	5.9	5.6	5.5	
UV index 1 month before exam (categorized)				
< 3.0	6.5	7.3	14.3	< 0.001
3.0–3.9	9.0	12.8	17.4	
4.0–4.9	13.6	18.0	15.6	
5.0–5.9	16.3	18.9	16.0	
6.0–6.9	27.9	19.0	13.1	
7.0–7.9	20.3	21.4	14.7	
≥ 8.0	6.5	2.8	8.9	
Skin type				
Hair colour				
Black	2.9	41.3	30.8	< 0.001
Red	2.2	0.7	0.3	
Blonde	11.0	0.2	0.8	
Light brown	12.1	2.0	7.6	
Medium brown	15.2	3.1	5.6	
Dark brown	14.9	14.2	22.5	
Other	0.2	0.3	0.1	
Missing	12.8	8.2	5.5	
Skin reaction to 1/2 h of sun exposure without sunscreen or protective clothing				
No skin reaction	23.8	58.0	40.2	< 0.001
Severe sunburn with blisters	2.7	0.4	1.2	
Severe sunburn for a few days with peeling	10.7	0.9	6.4	
Mildly burned with some tanning	30.6	3.4	17.4	
Turn darker without sunburn	11.9	23.2	25.8	
Other	0.4	0.6	1.0	
Missing	19.9	13.6	8.1	
Sun behaviours				
How often stay in the shade on a sunny day, when outside for > 1 h				
Always	1.8	6.6	5.7	< 0.001
Most of the time	8.5	14.1	11.5	
Sometimes	18.5	16.8	16.3	
Rarely	9.9	5.4	7.2	
Never	4.5	6.2	7.5	
Don't go in the sun	0.4	0.6	0.5	
Missing†	14.1	9.4	5.3	
Not asked†	42.4	40.9	46.0	
Number of sunburns in past year				
0	2.8	42.2	24.7	< 0.001
1	12.9	3.4	9.1	
2	12.6	3.0	7.1	
≥ 3	15.3	1.2	7.9	
Missing†	14.1	9.4	5.4	
Not asked†	42.4	40.9	46.0	
How often wear a hat that covers face, ears and neck				
Always	3.2	5.0	4.4	< 0.001
Most of the time	3.7	4.6	4.2	
Sometimes	6.9	8.9	6.5	
Rarely	6.4	6.4	4.6	
Never	23.0	24.2	28.5	
Missing†	14.4	9.9	5.8	
Not asked†	42.4	41.0	46.0	

Table 1 Continued

	Non-Hispanic White (<i>n</i> 7554)	Non-Hispanic Black (<i>n</i> 2991)	Hispanic (<i>n</i> 3774)	<i>P</i>
How often wear a long-sleeved shirt				
Always	0.9	1.2	3.8	< 0.001
Most of the time	1.6	2.5	3.3	
Sometimes	6.1	10.2	8.5	
Rarely	8.7	7.9	6.1	
Never	26.1	27.4	26.5	
Missing†	14.4	9.9	5.8	
Not asked†	42.4	40.9	46.0	
How often wear sunscreen				
Always	5.4	1.4	4.3	< 0.001
Most of the time	8.0	1.2	3.9	
Sometimes	11.1	4.0	7.7	
Rarely	7.1	3.4	5.1	
Never	11.6	39.2	27.3	
Missing†	14.4	9.9	5.8	
Not asked†	42.4	40.9	46.0	
Sunscreen SPF used most often				
None or < 15	28.3	49.6	34.5	< 0.001
15–29	10.9	3.3	6.8	
≥30	18.4	6.3	12.7	
Not asked†	42.4	40.9	46.0	
Other personal behaviours or characteristics				
Vitamin D supplementation during past 30 d (avg. IU/d)	1.8	1.0	0.8	< 0.001
BMI (kg/m ²)	28.1	30.1	28.4	< 0.001
Smoker				
Every day	22.4	22.7	13.9	< 0.001
Some days	3.2	3.5	7.8	
Not at all	27.7	15.5	19.2	
Month/year of exam				
Month				
January	2.9	3.7	9.5	
February	5.4	5.7	13.3	
March	5.3	10.4	12.0	
April	9.3	12.9	5.6	
May	10.3	12.6	4.9	
June	13.0	6.4	3.6	
July	8.1	6.0	2.9	
August	12.5	9.7	4.1	
September	10.9	8.1	7.5	
October	10.0	8.6	14.5	
November	7.7	8.9	12.5	
December	4.6	7.0	9.7	
Year				
2000	15.2	14.9	17.3	
2001–02	27.2	25.9	28.7	
2003–04	28.8	29.0	27.3	
2005–06	28.8	30.1	26.7	

SPF, sun protection factor.

Data are presented as percentages, unless otherwise specified.

†Question not asked in 2000 and 2001–02. Percentage missing is shown for 2003–04 and 2005–06.

analyses, we considered categorizations of the UV index in groups (<3.0, 3.0–3.9, 4.0–4.9, 5.0–5.9, 6.0–6.9, 7.0–7.9, ≥8.0). We also considered an interaction term between the UV index and overweight/obese BMI (25.0–25.9 kg/m², ≥30.0 kg/m²), because studies suggest that vitamin D may be stored in adipose tissue⁽⁴⁷⁾, with levels inversely related to obesity^(47,48).

Third, we estimated the population-level distribution of vitamin D levels by race and ethnicity in January, the month with the lowest average UV index⁽⁴⁵⁾. To do so, we adjusted vitamin D levels for each NHANES participant to January by performing the regression of vitamin D

v. race/ethnicity and eleven monthly dummies (February–December, relative to January). We then subtracted the coefficient for the month of each participant's NHANES laboratory examination from his/her actual vitamin D level, to estimate January-adjusted vitamin D. For each race/ethnicity, we obtained the 2nd, 5th–95th (in 5 percentage point increments) and 98th percentiles of adjusted vitamin D levels, and bootstrapped to obtain intermediate points in the distribution. (NHANES Restricted Data Center privacy regulations prohibited us from recording the entire distribution.) We performed a similar analysis stratified by race/ethnicity and quartile of the January UV index.

We then examined the proportion of each group estimated to be deficient (<15 ng/ml)⁽¹⁾ or insufficient (<20 ng/ml according to the Institute of Medicine-recommended thresholds⁽¹⁾, or <30 ng/ml according to alternative criteria for insufficiency suggested by some researchers⁽⁴⁹⁾) in vitamin D in January.

Results

Table 1 presents summary statistics. Non-Hispanic Whites were older, wealthier and had more sunburns than Hispanics and non-Hispanic Blacks. Non-Hispanic Whites had the highest average vitamin D levels (25.87 ng/ml), followed by Hispanics (20.72 ng/ml) and non-Hispanic Blacks (14.93 ng/ml).

Figure 1 shows racial/ethnic differences in the estimated number of minutes required to synthesize 15 μ g (600 IU) of vitamin D from sunshine in January, using the relationship between temperature and sun exposure described in the Methods. Hispanics required 23, 78 and 110 min more than fair-skinned whites in the counties containing Los Angeles, New York City and Chicago, respectively, based on UV index and expected body area exposed to sunlight. Dark-skinned blacks required ≥ 120 min more than fair-skinned whites in the vast majority of the USA. For comparison, in July, the racial difference was typically <15 min between Hispanics and fair-skinned whites, and <30 min between dark-skinned blacks and fair-skinned whites (not shown).

Table 2 shows regression results for key variables of interest. (See Appendix 2 for complete results.) A 1-point increase in the UV index 1 month prior to the examination was associated with a 0.51 ng/ml increase in 25(OH)D levels (95% CI 0.35, 0.67 ng/ml, $P < 0.001$; model 1). Non-Hispanic Black race and Hispanic ethnicity were associated with 25(OH)D levels lower by 7.47 ng/ml (95% CI -8.63, -6.31 ng/ml) and 3.41 ng/ml (95% CI -3.89, -2.93 ng/ml), respectively (both $P < 0.001$; model 1). This association was most evident when the UV index was < 5.0 (model 2).

Obesity, defined as a BMI ≥ 30.0 kg/m², was associated with a 2.51 ng/ml decrease in vitamin D (95% CI -3.86, -1.17 ng/ml, $P < 0.001$; model 3). Additionally, obese individuals had a 0.28 ng/ml attenuated association of vitamin D with the UV index (95% CI -0.50, -0.05 ng/ml, $P = 0.015$). For example, controlling for the interaction of BMI group with vitamin D (model 3), a 1-point increase in the UV index was associated with an average 0.66 ng/ml increase in vitamin D. For an obese individual, this association was attenuated to a 0.38 ng/ml increase in vitamin D (0.66 ng/ml average minus 0.28 ng/ml interaction term).

Figure 2 shows the estimated population-level distribution of serum vitamin D in January, by race/ethnicity (across all geographic locations). Mean January-adjusted vitamin D levels were 23.75 (SD 8.30) ng/ml, 19.29 (SD 6.94) ng/ml and

13.21 (SD 6.49) ng/ml for non-Hispanic Whites, Hispanics and non-Hispanic Blacks, respectively. Vertical lines in Fig. 2 indicate thresholds for vitamin D deficiency (<15 ng/ml)⁽¹⁾ and insufficiency (<20 ng/ml or <30 ng/ml, depending on the definition used)^(1,49). In January, an estimated 65.4% of non-Hispanic Blacks were deficient in vitamin D, compared with 28.9% of Hispanics and 14.0% of non-Hispanic Whites. An estimated 84.2% of non-Hispanic Blacks were insufficient (<20 ng/ml) in vitamin D *v.* 56.3% of Hispanics and 34.8% of non-Hispanic Whites. Using alternative criteria for insufficiency (<30 ng/ml), all non-Hispanic Blacks were insufficient in vitamin D, compared with 91.9% of Hispanics and 78.2% of non-Hispanic Whites.

Figure 3 repeats the analysis by race/ethnicity and geographic location (as categorized by approximate quartile of the January UV index in each participant's census tract of residence). Mean January-adjusted vitamin D levels ranged from 19.79 to 24.38 ng/ml for non-Hispanic Whites, from 16.74 to 19.90 ng/ml for Hispanics and from 10.62 to 14.27 ng/ml for non-Hispanic Blacks. For each race/ethnicity, mean January-adjusted vitamin D increased as the UV index rose. In the lowest quartile (January UV index < 3.44), an estimated 75.1% of non-Hispanic Blacks and 38.1% of Hispanics were deficient in vitamin D, compared with 25.7% of non-Hispanic Whites. In the highest quartile (January UV index > 6.00), an estimated 60.2% of non-Hispanic Blacks and 25.6% of Hispanics were deficient in vitamin D, compared with 12.8% of non-Hispanic Whites.

Discussion

Previous literature suggests that it is difficult for adults residing in Boston and north to produce enough vitamin D in winter^(43,47,50), resulting in widespread wintertime vitamin D deficiency at northern latitudes⁽⁵¹⁾. Our results suggest that wintertime vitamin D insufficiency in minority populations is pervasive across the USA. Using nationally representative NHANES data from 2000–2006, we estimated that 84.2–100% of non-Hispanic Blacks and 56.3–91.9% of Hispanics were insufficient in vitamin D in January (depending on a 20 *v.* 30 ng/ml threshold). Even in non-Hispanic Whites, vitamin D insufficiency during winter was not uncommon, with an estimated 34.8–78.2% of individuals affected.

Moreover, the association of race with wintertime vitamin D appeared at least as strong as that of geography. The estimated disparity in January vitamin D levels between non-Hispanic Whites and non-Hispanic Blacks across the USA (23.75 - 13.21 = 10.54 ng/ml, Fig. 2) exceeded that between non-Hispanic Whites residing in the lowest *v.* highest quartile of the January UV index (24.38 - 19.79 = 4.59 ng/ml, Fig. 3). Similarly, more non-Hispanic Blacks were estimated to be deficient in January




Fig. 1 (colour online) Racial/ethnic differences in estimated number of minutes required to synthesize 15 µg (600 IU) of vitamin D from sunlight, in January. Results shown for noontime, assuming no sunscreen and a relationship between average maximum daily temperature and sun exposure ($\geq 23.9^{\circ}\text{C}$ ($\geq 75^{\circ}\text{F}$): face, neck, hands, arms exposed; $15.6\text{--}23.8^{\circ}\text{C}$ ($60\text{--}74^{\circ}\text{F}$): face, neck, hands, lower arms exposed; -1.1 to 15.5°C ($30\text{--}59^{\circ}\text{F}$): face, neck, hands exposed; -9.4 to -1.2°C ($15\text{--}29^{\circ}\text{F}$): face, neck exposed; $\leq -9.5^{\circ}\text{C}$ ($\leq 14^{\circ}\text{F}$): face exposed) for: (a) black (darkest skin) minus white (fairest skin); (b) Hispanic minus white (fairest skin)

Table 2 Regression results. Dependent variable: serum 25-hydroxyvitamin D (ng/ml)

Regression model	Model 1 UV index		Model 2 UV index – groups		Model 3 Interaction with obesity	
	Mean	SE	Mean	SE	Mean	SE
Race/ethnicity						
Non-Hispanic White	Ref.	–	Ref.	–	Ref.	–
Non-Hispanic Black	–7.47***	0.59	–7.96***	0.24	–8.01***	0.24
Hispanic	–3.41***	0.25	–3.37***	0.24	–3.32***	0.25
UV index						
UV index 1 month before exam	0.51***	0.08			0.66***	0.11
Groups of UV index 1 month before exam						
UV index < 3			–3.15***	0.43		
3 ≤ UV index < 4			–2.50***	0.36		
4 ≤ UV index < 5			–2.23***	0.36		
5 ≤ UV index < 6			–0.88**	0.33		
6 ≤ UV index < 7			Ref.	–		
7 ≤ UV index < 8			–0.02	0.32		
UV index ≥ 8			0.39	0.38		
BMI						
Continuous (kg/m ²)	–0.28***	0.01	–0.28***	0.01		
25.0–30.0 kg/m ²					–0.27	0.72
≥30.0 kg/m ²					–2.51***	0.69
Interaction of UV index with BMI						
BMI 25.0–30.0 kg/m ² × UV index					–0.21	0.12
BMI ≥30.0 kg/m ² × UV index					–0.28*	0.11
<i>R</i> ²	0.29		0.29		0.28	
<i>F</i>	84.89		80.88		81.12	
<i>n</i>	14 319		14 319		14 319	

Ref., reference category.

Robust standard errors were used.

Coefficients for demographics, sun behaviours, other personal characteristics and month/year of exam are shown in Appendix 2.

P* < 0.05, *P* < 0.01, ****P* < 0.001.




Fig. 2 (colour online) Estimates of serum 25-hydroxyvitamin D, adjusted to January, by race/ethnicity; adults aged ≥ 18 years (*n* 14 319), National Health and Nutrition Examination Survey (NHANES), 2000–2006. Vertical lines represent thresholds for vitamin D deficiency (<15 ng/ml) and insufficiency (<20 ng/ml or <30 ng/ml, depending on the source)

vitamin D in the highest quartile of the UV index than were non-Hispanic Whites in the lowest quartile of the UV index (60.2% *v.* 25.7%, Fig. 3). The association of Hispanic ethnicity with vitamin D appeared similarly strong to geography; 25.7% of non-Hispanic Whites residing in counties in the lowest quartile of the January UV index and 25.6% of Hispanics residing in counties in the highest quartile of the January UV index were deficient in vitamin D (Fig. 3).

Alternatively viewed, 95% confidence intervals for our regression models suggested that a 1-point increase in the UV index was associated with a 0.35–0.67 ng/ml increase in 25(OH)D (coefficient = 0.51; Table 2, model 1). Since the January UV index ranges from 0 to 5 in the USA⁽⁴⁰⁾, geography contributed up to 3.35 ng/ml of wintertime vitamin D (0.67 × 5). This magnitude was about the same as the decrease in January vitamin D associated with Hispanic ethnicity (3.41 ng/ml) but well below that for non-Hispanic Black race (7.47 ng/ml; Table 2, model 1).

These results suggest a potentially more pervasive vitamin D insufficiency among non-Hispanic Blacks in winter than previously known. For example, a 2011 Institute of Medicine report using 2003–2006 NHANES data suggested that 53.6%, 27.2% and 10.6% of non-Hispanic Blacks, Mexican Americans and non-Hispanic Whites had vitamin D levels < 16 ng/ml (40 nmol/l)^(1,2), similar to the percentages we found for vitamin D deficiency in Fig. 2. Differences may be in part due to methodology; the Institute of Medicine adjusted vitamin D levels to northern latitudes (across seasons)⁽¹⁾, while we adjusted to January (across latitudes). Our analysis further adds the distribution of vitamin D levels by race/ethnicity and geography. Using the Institute of Medicine’s threshold of ≤ 20 ng/ml for vitamin D insufficiency, we found that 84.2% of non-Hispanic Blacks, 56.3% of Hispanics




Fig. 3 (colour online) Estimates of serum 25-hydroxyvitamin D, adjusted to January, by January UV index in census tract of residence; adults aged ≥ 18 years (n 14 319), National Health and Nutrition Examination Survey (NHANES), 2000–2006. Results shown for: (a) January UV index < 3.44 (quartile 1); (b) January UV index 3.44–4.16 (quartile 2); (c) January UV index 4.17–6.00 (quartile 3); and (d) January UV index > 6.00 (quartile 4). Vertical lines represent thresholds for vitamin D deficiency (< 15 ng/ml) and insufficiency (< 20 ng/ml or < 30 ng/ml, depending on the source)

and 34.8% of non-Hispanic Whites were insufficient in January; results were more extreme (including 78.2% of non-Hispanic Whites) at thresholds of ≤ 30 ng/ml.

Previous research also suggests that it may be infeasible to synthesize adequate quantities of vitamin D from sunshine in the northern USA^(52–54). Our results extend this finding to wide portions of the USA for Hispanics and non-Hispanic Blacks. In particular, it may be much more time-consuming, and therefore impractical, for minorities to synthesize the RDA of vitamin D from sunshine in winter, compared with non-Hispanic Whites (Fig. 1).

Our results support the need for further evidence to evaluate the benefits of prioritized vitamin D testing and supplementation of minority populations in winter. The US Preventive Services Task Force found incomplete evidence to recommend for or against vitamin D supplementation in postmenopausal women to prevent fractures⁽⁵⁵⁾, but did not consider the general population or subgroups based on race/ethnicity. Similarly, the Canadian province of British Columbia, which borders Washington state, suggested that most adults would benefit from a supplement of 20–25 μg (800–1000 IU) daily, but did not consider potential differences based on race/ethnicity⁽⁵⁶⁾. Some evidence from community-based studies suggests that adults could tolerate far higher doses without toxicity⁽⁵⁷⁾, although the exact threshold is unknown.

These potential benefits would need to be weighed against lower fracture risk in blacks⁽⁵⁸⁾ and recent research suggesting that black individuals tend to have lower levels of vitamin D-binding protein⁽⁵⁹⁾, potentially signifying fewer health risks from lower total vitamin D than the general population⁽⁵⁹⁾. Yet, given the possibility that vitamin D may be associated with disease^(1–23), it may be reasonable to investigate a potential association between racial disparities in vitamin D and racial disparities in mortality.

Limitations

NHANES conducts examinations at northern latitudes during summer months and southern latitudes during winter months⁽⁴³⁾. However, because the decrease in vitamin D associated with a UV index < 3 was larger than the increase associated with a UV index ≥ 8 (Table 2), we expect that deficiencies would be even more prevalent were wintertime data available at all latitudes. Second, we did not have information on dietary intake of vitamin D (outside of supplementation), physical activity and outdoor occupation, which may be positively associated with vitamin D^(48,60). Third, although our findings suggest that individuals with more sun exposure had higher vitamin D, we did not consider the risk of skin cancer.

Conclusions

Wintertime vitamin D insufficiency is pervasive among minority populations, potentially contributing to racial disparities in disease incidence and mortality. Additional research is needed to assess the benefits of preferential routine vitamin D supplementation in minorities.

Acknowledgements

Acknowledgements: This study was completed while G.B.T. was with the Departments of Population Health and Medicine, New York University School of Medicine, New York, NY, USA. The work was performed at the National Center for Health Statistics, Hyattsville, MD, USA and the National Bureau of Economic Research, Cambridge, MA, USA. **Financial support:** This study was funded by the A. David Mazzone Awards Program of the Dana-Farber Cancer Institute (grant number 2011_DR_002). The A. David Mazzone Awards Program had no role in the design, analysis or writing of this article. **Conflict of interest:** None. **Authorship:** conception and design: G.B.T., D.M.C. and N.L.K.; analysis and interpretation of the data: G.B.T., D.M.C., E.G. and N.L.K.; drafting of the article: G.B.T.; critical revision of the article for important intellectual content: G.B.T., D.M.C., E.G. and N.L.K.; final approval of the article: G.B.T., D.M.C., E.G. and N.L.K.; provision of study materials or patients: not applicable; statistical expertise: G.B.T.; administrative, technical or logistic support: G.B.T.; collection and assembly of data: G.B.T. **Ethics of human subject participation:** This study was considered exempt by the Institutional Review Board of Harvard Medical School and did not meet criteria for human subjects review for the Institutional Review Board of New York University School of Medicine.

References

- Institute of Medicine (2011) *Dietary Reference Intakes for Calcium and Vitamin D*. Washington, DC: The National Academies Press.
- Looker AC, Dawson-Hughes B, Calvo MS *et al.* (2002) Serum 25-hydroxyvitamin D status of adolescents and adults in two seasonal subpopulations from NHANES III. *Bone* **30**, 771–777.
- Hosseini-Nezhad A & Holick MF (2013) Vitamin D for health: a global perspective. *Mayo Clin Proc* **88**, 720–755.
- Zittermann A, Iodice S, Pilz S *et al.* (2012) Vitamin D deficiency and mortality risk in the general population: a meta-analysis of prospective cohort studies. *Am J Clin Nutr* **95**, 91–100.
- Lange N, Litonjua AA, Gibbons FK *et al.* (2013) Pre-hospital vitamin D concentration, mortality, and bloodstream infection in a hospitalized patient population. *Am J Med* **126**, 640.e19–640.e27.
- Toner CD, Davis CD & Milner JA (2010) The vitamin D and cancer conundrum: aiming at a moving target. *J Am Diet Assoc* **110**, 1492–1500.
- Davis CD (2008) Vitamin D and cancer: current dilemmas and future research needs. *Am J Clin Nutr* **88**, issue 2, 565S–569S.
- Moan J, Porojnicu AC, Dahlback A *et al.* (2008) Addressing the health benefits and risks, involving vitamin D or skin cancer, of increased sun exposure. *Proc Natl Acad Sci USA* **105**, 668–673.
- Freedman DM, Looker AC, Chang S *et al.* (2007) Prospective study of serum vitamin D and cancer mortality in the United States. *J Natl Cancer Inst* **99**, 1594–1602.
- Lappe JM, Travers-Gustafson D, Davies KM *et al.* (2007) Vitamin D and calcium supplementation reduces cancer risk: results of a randomized trial. *Am J Clin Nutr* **85**, 1586–1591.
- Garland CF, Garland FC, Gorham ED *et al.* (2006) The role of vitamin D in cancer prevention. *Am J Public Health* **96**, 252–261.
- Moukayed M & Grant WB (2013) Molecular link between vitamin D and cancer prevention. *Nutrients* **5**, 3993–4021.
- Robsahm TE, Schwartz GG & Tretli S (2013) The inverse relationship between 25-hydroxyvitamin D and cancer survival: discussion of causation. *Cancers (Basel)* **5**, 1439–1455.
- Tomson J, Emberson J, Hill M *et al.* (2013) Vitamin D and risk of death from vascular and non-vascular causes in the Whitehall study and meta-analyses of 12,000 deaths. *Eur Heart J* **34**, 1365–1374.
- Wang L, Song Y, Manson JE *et al.* (2012) Circulating 25-hydroxy-vitamin D and risk of cardiovascular disease: a meta-analysis of prospective studies. *Circ Cardiovasc Qual Outcomes* **5**, 819–829.
- Song Y, Wang L, Pittas AG *et al.* (2013) Blood 25-hydroxy vitamin D levels and incident type 2 diabetes: a meta-analysis of prospective studies. *Diabetes Care* **36**, 1422–1428.
- Bischoff-Ferrari HA, Willett WC, Wong JB *et al.* (2009) Prevention of nonvertebral fractures with oral vitamin D and dose dependency: a meta-analysis of randomized controlled trials. *Arch Intern Med* **169**, 551–561.
- Bischoff-Ferrari HA, Dawson-Hughes B, Staehelin HB *et al.* (2009) Fall prevention with supplemental and active forms of vitamin D: a meta-analysis of randomised controlled trials. *BMJ* **339**, b3692.
- Sohl E, van Schoor NM, de Jongh RT *et al.* (2013) Vitamin D status is associated with functional limitations and functional decline in older individuals. *J Clin Endocrinol Metab* **98**, E1483–E1490.
- Kupferschmidt K (2012) Uncertain verdict as vitamin D goes on trial. *Science* **337**, 1476–1478.
- Murdoch DR, Slow S, Chambers ST *et al.* (2012) Effect of vitamin D₃ supplementation on upper respiratory tract infections in healthy adults: the VIDARIS randomized controlled trial. *JAMA* **308**, 1333–1339.
- Marketwire (2008) Novacea announces preliminary findings from data analysis of Ascent-2 phase 3 trial. <http://www.marketwire.com/press-release/novacea-announces-preliminary-findings-from-data-analysis-of-ascent-2-phase-3-trial-nasdaq-novc-864465.htm> (accessed November 2012).
- Lappe JM & Heaney RP (2012) Why randomized controlled trials of calcium and vitamin D sometimes fail. *Dermato-endocrinol* **4**, 95–100.
- Clemens TL, Adams JS, Hendeson SL *et al.* (1982) Increased skin pigment reduces the capacity of skin to synthesise vitamin D₃. *Lancet* **1**, 74–76.
- Ginde AA, Liu MC & Camargo CA Jr (2009) Demographic differences and trends of vitamin D insufficiency in the US population, 1988–2004. *Arch Intern Med* **169**, 626–632.
- Holick MF (2003) Vitamin D: a millennium perspective. *J Cell Biochem* **88**, 296–307.
- Jones G (2008) Pharmacokinetics of vitamin D toxicity. *Am J Clin Nutr* **88**, issue 2, 582S–586S.
- Hypönen E & Power C (2007) Hypovitaminosis D in British adults at age 45 y: nationwide cohort study of dietary and lifestyle predictors. *Am J Clin Nutr* **85**, 860–868.

29. Robsahm TE, Tretli S, Dahlback A *et al.* (2004) Vitamin D₃ from sunlight may improve the prognosis of breast-, colon- and prostate cancer (Norway). *Cancer Causes Control* **15**, 149–158.
30. Lagunova Z, Porojnicu AC, Dahlback A *et al.* (2007) Prostate cancer survival is dependent on season of diagnosis. *Prostate* **67**, 1362–1370.
31. Hanchette CL & Schwartz GG (1992) Geographic patterns of prostate cancer mortality. Evidence for a protective effect of ultraviolet radiation. *Cancer* **70**, 2861–2869.
32. Schwartz GG & Hanchette CL (2006) UV, latitude, and spatial trends in prostate cancer mortality: all sunlight is not the same (United States). *Cancer Causes Control* **17**, 1091–1101.
33. Grant WB (2006) Lower vitamin-D production from solar ultraviolet-B irradiance may explain some differences in cancer survival rates. *J Natl Med Assoc* **98**, 357–364.
34. Grant WB & Garland CF (2006) The association of solar ultraviolet B (UVB) with reducing risk of cancer: multifactorial ecologic analysis of geographic variation in age-adjusted cancer mortality rates. *Anticancer Res* **26**, 2687–2699.
35. Grant WB (2011) The roles of ultraviolet-B irradiance, vitamin D, apolipoprotein E ε₄, and diet in the risk of prostate cancer. *Cancer Causes Control* **22**, 157–158.
36. Taksler GB, Cutler DM, Giovannucci E *et al.* (2013) Ultraviolet index and racial differences in prostate cancer incidence and mortality. *Cancer* **119**, 3195–3203.
37. Centers for Disease Control and Prevention, National Center for Health Statistics (1999)–2000, 2001–2002, 2003–2004, 2005–2006) *National Health and Nutrition Examination Survey Data*. Hyattsville, MD: US Department of Health and Human Services, CDC; available at <http://www.cdc.gov/nchs/nhanes.htm>
38. National Aeronautics and Space Administration (2010) Erythral UV exposure, Total Ozone Mapping Spectrometer. http://toms.gsfc.nasa.gov/ery_uv/euv_v8.html (accessed August 2010).
39. Boscoe FP & Schymura MJ (2006) Solar ultraviolet-B exposure and cancer incidence and mortality in the United States, 1993–2002. *BMC Cancer* **6**, 264.
40. Environmental Protection Agency (1994) *Experimental UV Index*. EPA 430-F-94-019. Washington, DC: EPA.
41. Herman JR & Celarier EA (1999) Erythral exposure data product. <http://macuv.gsfc.nasa.gov/doc/erynotes.pdf> (accessed August 2010).
42. Fitzpatrick TB (1975) Soleil et peau (Sun and skin). *Journal de Médecine Esthétique* **2**, 33–34.
43. Terushkin V, Bender A, Psaty EL *et al.* (2010) Estimated equivalency of vitamin D production from natural sun exposure versus oral vitamin D supplementation across seasons at two US latitudes. *J Am Acad Dermatol* **62**, 929.e1–929.e9.
44. Kiedron P, Stierle S & Lantz K (2007) Instantaneous UV index and daily UV dose calculations. <http://www.esrl.noaa.gov/gmd/grad/neubrew/docs/UVindex.pdf> (accessed February 2014).
45. National Climatic Data Center (2011) GHCN-Daily v2.92. ftp://ftp.ncdc.noaa.gov/pub/data/ghcn/daily/ghcnd_all.tar.gz (accessed April 2012).
46. Lund CC & Broder NC (1944) Estimation of areas of burns. *Surg Gynecol Obstet* **79**, 352–358.
47. Holick MF (2004) Vitamin D: importance in the prevention of cancers, type 1 diabetes, heart disease, and osteoporosis. *Am J Clin Nutr* **79**, 362–371.
48. McCullough ML, Weinstein SJ, Freedman DM *et al.* (2010) Correlates of circulating 25-hydroxyvitamin D: Cohort Consortium Vitamin D Pooling Project of Rarer Cancers. *Am J Epidemiol* **172**, 21–35.
49. Holick MF (2007) Vitamin D deficiency. *N Engl J Med* **357**, 266–281.
50. Webb AR, Kline L & Holick MF (1988) Influence of season and latitude on the cutaneous synthesis of vitamin D₃: exposure to winter sunlight in Boston and Edmonton will not promote vitamin D₃ synthesis in human skin. *J Clin Endocrinol Metab* **67**, 373–378.
51. Merewood A, Mehta SD, Grossman X *et al.* (2010) Widespread vitamin D deficiency in urban Massachusetts newborns and their mothers. *Pediatrics* **125**, 640–647.
52. Godar DE, Pope SJ, Grant WB *et al.* (2011) Solar UV doses of adult Americans and vitamin D(3) production. *Dermato-endocrinology* **3**, 243–250.
53. Vitamin D Council (2013) How do I get the vitamin D my body needs? <http://www.vitamindcouncil.org/about-vitamin-d/how-do-i-get-the-vitamin-d-my-body-needs/> (accessed July 2013).
54. Webb AR & Engelsen O (2006) Calculated ultraviolet exposure levels for a healthy vitamin D status. *Photochem Photobiol* **82**, 1697–1703.
55. Moyer VA; US Preventive Services Task Force (2013) Vitamin D and calcium supplementation to prevent fractures in adults: US Preventive Services Task Force recommendation statement. *Ann Intern Med* **158**, 691–696.
56. Anon. (2011) Vitamin D: a patient guide for adults. http://www.bcguidelines.ca/pdf/vitamind_patient_guide.pdf (accessed July 2013).
57. Garland CF, French CB, Baggerly LL *et al.* (2011) Vitamin D supplement doses and serum 25-hydroxyvitamin D in the range associated with cancer prevention. *Anticancer Res* **31**, 607–611.
58. Jacobsen SJ, Goldberg J, Miles TP *et al.* (1990) Hip fracture incidence among the old and very old: a population-based study of 745,435 cases. *Am J Public Health* **80**, 871–873.
59. Powe CE, Evans MK & Wenger J (2013) Vitamin D-binding protein and vitamin D status of black Americans and white Americans. *N Engl J Med* **369**, 1991–2000.
60. Freedman DM, Dosemeci M & McGlynn K (2002) Sunlight and mortality from breast, ovarian, colon, prostate, and non-melanoma skin cancer: a composite death certificate based case-control study. *Occup Environ Med* **59**, 257–262.

Appendix 1

Further details on the outdoor exposure required to synthesize vitamin D

As described in the main text, we estimated the number of minutes outdoors at noon in each calendar month for fair-skinned white, Hispanic and dark-skinned black individuals to synthesize 15 µg (600 IU) of vitamin D, the (RDA) for adults ≤70 years⁽¹⁾. This calculation was defined as the minimum erythral dose (MED) of UV radiation required for individuals of each race to synthesize 15 µg (600 IU) of vitamin D, divided by the MED supplied per minute of noontime outdoor exposure. In this Appendix, we describe our calculations in more detail.

For the numerator, we state in the main text that we followed dermatology literature to expect that fair-skinned white, Hispanic and dark-skinned black individuals would require 0.25, 0.56 and 1.25 MED of UV radiation, respectively, to synthesize 15 µg (600 IU) of vitamin D, with 25.5% of body surface area – the head, neck, arms and hands – uncovered. These numbers derive from an estimated 0.42, 0.94 and 2.08 MED (87.6, 197.0 and 437.8 J/m²) of UV radiation, respectively, to synthesize 25 µg

(1000 IU) of vitamin D⁽⁴³⁾. Since the dermatology literature suggests linearity of this relationship⁽⁴³⁾, we multiplied by 60% (15 µg/25 µg, 600 IU/1000 IU).

For the denominator, we used the definition of the UV index (1 unit = 0.0072 MED/min)⁽⁴⁴⁾. Each unit of the UV index is defined as 25 mW/m², or 0.43 MED/h⁽⁴⁴⁾.

We also assumed a relationship between average maximum daily temperature and body parts exposed to

sunlight (≥23.9°C (≥75°F): face, neck, hands, arms exposed; 15.6–23.8°C (60–74°F): face, neck, hands, lower arms exposed; –1.1 to 15.5°C (30–59°F): face, neck, hands exposed; –9.4 to –1.2°C (15–29°F): face, neck exposed; ≤–9.5°C (≤14°F): face exposed). Following previous literature, we assumed that the face, neck, hands, upper arms and lower arms encompassed 3.5%, 2.0%, 6.0%, 8.0% and 6.0% of body surface area, respectively⁽⁴⁶⁾.

Appendix 2

Complete regression results for Table 2. Dependent variable: serum 25-hydroxyvitamin D (ng/ml)

Regression model	Model 1†	Model 2	Model 3	Model 4†	Model 5†
Description	UV index	UV index – quartiles	UV index – median	UV index groups	Interaction with obesity
Race/ethnicity (omitted: Non-Hispanic White)					
Non-Hispanic Black	–7.47***	–7.49***	–7.87***	–7.96***	–8.01***
Hispanic	–3.41***	–3.38***	–3.39***	–3.37***	–3.32***
UV index					
UV index 1 month before exam (continuous)					
UV index 1 month before exam	0.51***				0.66***
Non-Hispanic Black × UV index	–0.09				
BMI 25.0–30.0 kg/m ² × UV index					–0.21
BMI ≥ 30.0 kg/m ² × UV index					–0.28*
Quartiles of UV index 1 month before exam (omitted: Quartile 1)					
Quartile 2		0.41			
Quartile 3		1.45***			
Quartile 4		1.27***			
Non-Hispanic Black × quartile 2		–0.82			
Non-Hispanic Black × quartile 3		–0.62			
Non-Hispanic Black × quartile 4		–0.62			
Half of UV index 1 month before exam (omitted: UV index < median)					
UV index ≥ median			1.15***		
Non-Hispanic Black × UV index ≥ median			–0.24		
Groups of UV index 1 month before exam (omitted: 6–7)†					
UV index < 3				–3.15***	
3 ≤ UV index < 4				–2.50***	
4 ≤ UV index < 5				–2.23***	
5 ≤ UV index < 6				–0.88**	
7 ≤ UV index < 8				–0.02	
UV index ≥ 8				0.39	
Demographics					
Age	–0.05***	–0.05***	–0.05***	–0.05***	–0.05***
Female	–0.26	–0.27	–0.27	–0.25	–0.30
Married	0.81***	0.83***	0.83***	0.81***	0.87***
Annual household income (omitted: < \$US 25 000)					
\$US 25 000–34 999	0.16	0.15	0.14	0.17	0.20
≥ \$US 35 000	0.44*	0.42*	0.43*	0.44*	0.49*

Continued

Regression model	Model 1†	Model 2	Model 3	Model 4†	Model 5†
Description	UV index	UV index – quartiles	UV index – median	UV index groups	Interaction with obesity
Skin type					
Hair colour (omitted: Black)					
Red	1.78*	1.80*	1.80*	1.72*	1.84*
Blonde	2.25***	2.23***	2.25***	2.24***	2.30***
Light brown	2.10***	2.11***	2.12***	2.09***	2.12***
Medium brown	1.69***	1.66***	1.68***	1.67***	1.70***
Dark brown	1.11***	1.08***	1.11***	1.06***	1.17***
Other	2.65	2.79*	2.82*	2.68	2.70
Missing	2.07***	2.09***	2.11***	2.03***	2.08***
Skin reaction to 1/2 h of sun exposure without sunscreen or protective clothing (omitted: none)					
Severe sunburn with blisters	-0.18	-0.18	-0.19	-0.18	-0.19
Severe sunburn for a few days with peeling	-0.21	-0.19	-0.18	-0.19	-0.24
Mildly burned with some tanning	0.91***	0.92***	0.92***	0.94***	0.94***
Turn darker without sunburn	0.51*	0.54*	0.54*	0.51*	0.51*
Other	-1.33	-1.32	-1.33	-1.27	-1.21
Missing	-0.59	-0.63	-0.62	-0.60	-0.49
Sun behaviours					
How often stay in the shade on a sunny day, when outside for > 1 h (omitted: sometimes)					
Always	-1.95***	-1.93***	-1.94***	-2.01***	-2.17***
Most of the time	-1.69***	-1.68***	-1.68***	-1.72***	-1.77***
Rarely	2.03***	2.01***	2.00***	2.05***	2.09***
Never	1.46**	1.42**	1.42**	1.48**	1.53**
Don't go in the sun	5.74	5.52	5.69	5.53	5.80
Missing	6.92***	6.85***	6.97***	6.81***	6.53***
Number of sunburns in past year (omitted: zero)					
1	1.16***	1.17***	1.17***	1.14***	1.26***
2	0.87***	0.87***	0.87***	0.86***	0.93***
≥ 3	0.61***	0.61***	0.62***	0.61***	0.66***
Missing	5.13	5.08	5.10	4.91	5.50
How often wear a hat that covers face, ears and neck (omitted: never)					
Always	0.50	0.50	0.51	0.52	0.32
Most of the time	-0.01	0.03	0.04	0.00	-0.03
Sometimes	0.23	0.25	0.26	0.27	0.24
Rarely	-0.61	-0.58	-0.58	-0.57	-0.61
How often wear a long-sleeved shirt (omitted: never)					
Always	-1.85*	-1.88*	-1.88*	-1.79*	-1.60*
Most of the time	-2.09***	-2.14***	-2.18***	-2.05***	-1.95***
Sometimes	-0.49	-0.53	-0.53	-0.43	-0.48
Rarely	-0.93*	-0.93*	-0.94*	-0.91*	-0.82*
How often wear sunscreen (omitted: never)					
Always	2.17	2.16	2.13	2.25	2.22
Most of the time	1.88	1.87	1.85	1.95	1.87
Sometimes	1.36	1.38	1.34	1.41	1.34
Rarely	0.13	0.12	0.09	0.22	0.11

Continued

Regression model	Model 1†	Model 2	Model 3	Model 4†	Model 5†
Description	UV index	UV index – quartiles	UV index – median	UV index groups	Interaction with obesity
Sunscreen SPF used most often (omitted: < 15)					
15–29	–0.45	–0.38	–0.36	–0.49	–0.53
≥30	–0.72	–0.67	–0.66	–0.75	–0.80
Missing	0.18	0.24	0.23	0.21	0.11
Missing data for hat, long-sleeved shirt, sunscreen					
Missing	–10.54**	–10.39**	–10.54**	–10.14**	–10.49*
Other personal behaviours or characteristics					
Vitamin D supplementation during past 30 d (avg. IU/d)	0.28***	0.28***	0.28***	0.28***	0.27***
Smoker					
Every day	–1.26***	–1.23***	–1.23***	–1.28**	–1.15***
Some days	0.25	0.28	0.27	0.22	0.23
Missing	–0.33	–0.30	–0.31	–0.33	–0.34
BMI					
BMI (kg/m ²)	–0.28***	–0.28***	–0.28***	–0.28***	
25.0–30.0 kg/m ²					–0.27
≥30.0 kg/m ²					–2.51***
Month/year of exam					
Month (omitted: January)					
February	–0.39	–0.32	–0.39	–0.42	–0.34
March	–1.11*	–0.97	–0.90	–1.25*	–1.16*
April	0.03	0.02	0.07	–0.39	0.15
May	–0.58	0.68	0.68	–1.25*	–0.50
June	0.44	2.00***	2.04***	–0.52	0.52
July	1.98**	3.16***	3.20***	1.33*	2.12***
August	2.19***	3.20***	3.24***	1.52**	2.25***
September	0.55	2.11***	2.22***	–0.27	0.64
October	0.44	1.96***	1.92***	–0.36	0.55
November	0.96	1.11*	1.16*	0.43	1.08*
December	1.23*	1.30*	1.38**	0.94	1.29*
Year (omitted: 2000)					
2001–02	–1.63***	–1.63***	–1.67***	–1.43***	–1.64***
2003–04	–1.82	–1.96	–1.94	–1.83	–1.80
2005–06	–0.86	–1.04	–1.05	–0.79	–0.87
Intercept	30.60***	31.98***	32.16***	34.96***	23.49***
<i>R</i> ²	0.29	0.29	0.28	0.29	0.28
<i>F</i>	84.89	80.04	84.73	80.88	81.12
<i>n</i>	14 319	14 319	14 319	14 319	14 319

SPF, sun protection factor.
 Robust standard errors were used.
 P* < 0.05, *P* < 0.01, ****P* < 0.001.
 †Model presented in main article.