

Research Article

Cite this article: Ciuca L *et al.* (2025) Uncovering human *Dirofilaria repens* infections: new cases in Southern Italy. *Parasitology*, 1–10. <https://doi.org/10.1017/S0031182025000290>

Received: 2 December 2024
Revised: 10 February 2025
Accepted: 28 February 2025

Keywords:

Dirofilaria repens-gravid females;
epidemiology; human dirofilariosis;
Southern Italy; zoonotic infection

Corresponding author: Lavinia Ciuca;
Email: lavinia.ciuca@unina.it

Uncovering human *Dirofilaria repens* infections: new cases in Southern Italy

Lavinia Ciuca¹ , Simona Gabrielli², Patrizia Forgiione³, Evaristo Di Napoli¹, Orlando Paciello¹, Maria Vittoria Panariello⁴, Marianna Ascierto⁴, Luciana Petrucci⁴, Maria Ortensia Montella¹, Maria Paola Maurelli¹  and Laura Rinaldi¹ 

¹Department of Veterinary Medicine and Animal Production, Center for Monitoring of Parasitosis (CREMOPAR), University of Naples Federico, Naples, Italy; ²Department of Public Health and Infectious Diseases, University of Rome 'Sapienza', Rome, Italy; ³Local Health Authority Naples 1 Center, Pellegrini Hospital, Naples, Italy and ⁴Azienda of Colli of Naples, UOC Microbiology and Virology, Naples, Italy

Abstract

Dirofilaria repens is the primary etiological agent of human dirofilariosis in the Old World, with Italy reporting the highest number of cases in Europe. This study describes two new cases of *D. repens* infection in humans, in southern Italy, where canine dirofilariosis is endemic. The first case involved a 33-year-old man from Caserta (Campania, Southern Italy) who presented with a subcutaneous mass on the upper eyelid. Surgical excision revealed an immature female *D. repens* worm measuring 14 cm, lacking microfilariae in both the uterus and peripheral blood. The second case was a 67-year-old man from Pozzuoli (Metropolitan City of Naples, Southern Italy) with an oval-shaped nodule in the left frontal scalp region. A live gravid female *D. repens* worm measuring 15–16 cm was extracted, also without microfilariae in the peripheral blood and no male worm detected. PCR sequencing confirmed a 100% match with *D. repens*. Both patients tested positive for *D. repens* antibodies by IgG ELISA. These cases underscore the continuous spread of human dirofilariosis in southern Italy and highlight diagnostic challenges due to variable clinical presentations. The discovery of a gravid female without microfilaremia suggests complexities in the parasite's life cycle in humans, challenging the notion of humans as strict dead-end hosts. Given the rising prevalence in both humans and dogs, a comprehensive epidemiological study is recommended. Inclusion of dirofilariosis in the national surveillance system for notifiable diseases would improve case identification and tracking, aiding in better monitoring and control of this zoonotic infection.

Introduction

Dirofilaria immitis and *Dirofilaria repens* (Spirurida, Onchocercidae) are zoonotic vector-borne filarial parasites, transmitted by Culicidae mosquitoes, with dogs as the main reservoirs of infection (Capelli *et al.*, 2018; Genchi and Kramer, 2020; Perles *et al.*, 2024), in which the nematodes may cause heartworm disease or subcutaneous nodules, respectively (Simón *et al.*, 2022). Data reported in recent studies shows an increase in the prevalence of both *D. immitis* and *D. repens* in Europe and in southeastern regions of Asia and Africa (McCall *et al.*, 2008; Capelli *et al.*, 2018; Genchi and Kramer, 2020). In Italy, the changing distribution patterns of both *D. immitis* and *D. repens* show an increase in their prevalence in dogs from areas considered non-endemic, such as regions in southern Italy (Genchi *et al.*, 2009, 2019; Panarese *et al.*, 2022; Ciucă *et al.*, 2023; Napoli *et al.*, 2023). Human infections are mainly due to *D. repens*, which remains a persistent public health problem (Szénási *et al.*, 2008; Montoya-Alonso *et al.*, 2010; Ciucă *et al.*, 2018; Simón *et al.*, 2022; Perles *et al.*, 2024). Following transmission of the infecting larvae (L3) by mosquitoes, humans are usually accidental hosts for *D. immitis* and *D. repens*, although cases with fertile adults and/or microfilariaemia due to *D. repens* have been reported (Grandi *et al.*, 2008; Poppert *et al.*, 2009; Sergiev *et al.*, 2009). Zoonotic infection results in the formation of nodular structures as a consequence of the circumscribed inflammatory response mounted around dead or dying larval stages. In most patients, *D. immitis* localizes in the lungs, causing pulmonary dirofilariosis, while *D. repens* localizes in the subcutaneous tissue – causing the formation of subcutaneous nodules – or under the conjunctiva (ocular dirofilariosis). Over the years, several cases of pulmonary nodules caused by *D. repens* have also been reported in the Old World (Pampiglione *et al.*, 2009; Benzaquen *et al.*, 2015; Gabrielli *et al.*, 2021; Momčilović *et al.*, 2022). Pulmonary dirofilariosis can be confused with pulmonary neoplasms (benign, carcinoma, metastases), tuberculosis or fungal infections (Awe *et al.*, 1975; Ro *et al.*, 1989; Oliva *et al.*, 2019) and represents a diagnostic challenge (Simón *et al.*, 2022). Between 1977 and 2016,

© The Author(s), 2025. Published by Cambridge University Press. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted re-use, distribution and reproduction, provided the original article is properly cited.



Figure 1. Case 1. (A) Patient with a serpiginous lump on the upper eyelid of the left eye at first presentation. (B, C) Patient with swelling of the upper eyelid of the left eye in the form of a lump after one week of treatment. (D) Patient after surgery. (E) Worm recovered after excision of the nodule from upper eyelid of the left eye.

more than 3500 human cases due to *D. repens* and 25 due to *D. immitis* were reported in Europe (Genchi and Kramer, 2017; Capelli *et al.*, 2018). Since humans are not natural hosts, microfilariae are usually absent from peripheral blood. However, the parasite occasionally evades the host's immune system to reach sexual maturity, resulting in microfilaremic infections that have so far been described in 23 human cases (Simón *et al.*, 2022; Tasić-Otasevic *et al.*, 2023). Italy is traditionally endemic for human dirofilariosis and is one of the countries with the highest number of human cases identified so far (Muro *et al.*, 1999; Avellis *et al.*, 2011; Capelli *et al.*, 2018). In fact, the number of human cases of dirofilariosis caused by *D. repens* reported in the literature since Addario's first observation in 1885 amounted to approximately 410 in 1995, 181 of which had occurred in Italy (Addario, 1885; Pampiglione and Rivasi, 2000). In 2001, there was a further description of 60 cases between 1990 and 1999 (Pampiglione *et al.*, 2001). More recently, eight further cases of *D. repens* infection were described by Gabrielli *et al.* (2021). The present study aims to describe two new human cases of *D. repens* infection occurring in a region of southern Italy which has recently become endemic for canine dirofilariosis (Ciucă *et al.*, 2023).

Materials and methods

Case histories and sampling of adult nematodes

First case (case 1)

In October 2023, a 33-year-old Italian-born man living in Caserta (Campania, southern Italy) was referred to the emergency hospital (Ospedale Dei Pellegrini) of Naples, with complaints of pruritus and swelling of the left upper eyelid (Figure 1A). In his description, he mentioned that he had been taking long walks along the river at Castel Volturno six months prior to his admission to the hospital. The man reported that he had been living with two dogs, aged 3 and 6 years old, for a period of 5 years. At the initial ophthalmological examination, the patient exhibited an oedema in a serpentine shape on the upper eyelid of the left eye (Figure 1A). No evidence of swelling or similar rashes was observed in other parts of the body. The ophthalmological examination revealed that

the visual acuity in both eyes was within the normal range. Given the possibility of a parasitic infection, the patient was treated with a dose of 12 mg of ivermectin. The following day, the patient was transferred to another hospital in Naples. Subsequently, he underwent the following additional diagnostic procedures: a chest X-ray, a CT scan, an ECG, blood tests, copromicroscopy examination and an IgG test for SARS-CoV-2. With the exception of the detection of *Blastocystis* spp. in the stool sample, all results were within the normal range. The patient was also examined by a dermatological specialist who prescribed a course of albendazole 400 mg twice a day, for 7 days. After treatment, the patient returned with a swelling of the upper eyelid in the form of a lump (Figure 1B,C). The snake-like appearance was no longer present. The patient was then referred for surgical excision of the nodule that had formed. A live and motile worm of approximately 14 cm in length was found during the surgical excision of the nodule on the left upper eyelid (Figure 1D,E; supplementary file 1).

Second case (case 2)

In September 2024, a 67-year-old Italian man from Pozzuoli, a coastal town in the Metropolitan City of Naples (southern Italy) was referred to a private dermatology clinic in Naples for an oval-shaped subcutaneous mass located in the left frontal region (Figure 2A). An ultrasound of the skin showed an oval-shaped nodule measuring 7.7×3.9 mm, hypoechoic with well-defined, avascular margins located in the scalp area (Figure 2B). Based on the ultrasound, a cyst or fibroma of the scalp was suspected, and surgical removal was recommended. During the surgical removal, a live worm measuring 10 cm in length was found (Figure 2C,D). The patient had no history of travel abroad in the past 3 years and owned an 18-month-old dog.

The 2 adult nematodes (the first along with the excised nodule from Case 1 and the second from Case 2), were sent to the Laboratory of Parasitology of the University of Naples Federico II, for histological and molecular analysis. Based on the final surgical results, there was a suspicion of infection with *Dirofilaria* spp. At this point, blood and serum samples were obtained from both patients to verify the presence of microfilariae in the blood and of specific antibodies against *Dirofilaria* spp. in the serum. Similarly




Figure 2. Case 2. (A) Patient with a subcutaneous mass located in the left frontal region of the scalp; (B) an ultrasound of the skin revealed an oval-shaped, hypoechoic nodule measuring 7.7×3.9 mm with well-defined, avascular margins in the scalp area, suggestive of the presence of a nematode; (C) patient after surgery; (D) a 10 cm-long nematode surgically removed from the nodule in the scalp area.

blood and serum samples were collected from all dogs owned by both patients (2 from Case 1 and 1 from Case 2) and tested for the presence of microfilariae, *D. immitis* antigens and antibodies against *D. repens* somatic antigens.

The immunological tests were performed at the Department of Public Health and Infectious Diseases, University of Rome 'Sapienza', Rome, Italy.

Laboratory investigation of the human cases: histology, molecular analysis, modified Knott's test and immunological tests

The worms and nodule samples (Case 1 and Case 2) were excised, then preserved in neutral buffered formalin to maintain tissue integrity. After fixation, they were embedded in paraffin, allowing for thin slicing at 3–5 μ m thickness for detailed microscopic analysis. The sections were then stained with haematoxylin and eosin, a standard method for distinguishing cellular and tissue structures due to haematoxylin staining nuclei blue and eosin staining the cytoplasm and extracellular matrix pink. Digital images were captured using a Panoramic SCAN II (3DHISTECH) digital slide scanner, enabling detailed examination and production of representative images for further analysis.

For molecular identification, genomic DNA was extracted from 25 mg of tissue from each nematode, using the DNeasy® Blood and Tissue kit (Qiagen, Germany), following the manufacturer's instructions. Molecular analyses were performed in accordance with the multiplex PCR protocol described by Rishniw *et al.* (2006) (5.8 + ITS2 region) which allows for the simultaneous detection of *D. immitis* and *D. repens*.

A modified Knott's test was used for the detection of circulating microfilariae of *D. repens* (Genchi *et al.*, 2021) as outlined below. One mL of EDTA blood (3×1 mL for each tube) was mixed with 9 mL of distilled water and centrifuged for 3–5 min at 170 g. The supernatant was then removed from the tube and the contents were stained with 1–2 drops of 1% methylene blue. A drop was placed on a microscope slide covered with a cover slip and observed under an optical microscope at $100\times$.

The patient serum samples were analysed using an home-made ELISA test to detect the specific IgG antibody response. This involved the use of adult *D. immitis* and *D. repens* somatic antigens as described in Cabrera *et al.* (2018) and Mendoza-Roldán *et al.* (2021). The antigens were isolated from worms obtained by necropsy of naturally infected dogs, washed, and then they were macerated and sonicated (three cycles of 70 kHz, 30 s) in sterile saline solution (Prieto *et al.*, 1997). The homogenate was centrifuged at 16 000g for 30 min. The supernatant was dialyzed against 0.01M PBS, pH 7.2. The protein concentration was measured (Bradford, 1976) and an ELISA microplate was coated with antigens at a final concentration of 0.8 μ g/ μ L. The serum sample was tested in solid-phase ELISA at dilution 1:80 and 1:150 to detect anti-*D. repens* and anti-*D. immitis* IgG antibodies, respectively. Goat anti-human-IgG-antihuman IgG (H + L) conjugated to horseradish peroxidase (Sig-ma- Aldrich, MO, USA) was used as a secondary antibody at 1:40.000 dilution in both cases. The Optical density (OD) was determined at a wavelength of 492 nm (Easy-Reader, Bio Rad, CA, USA). The cut-off points (OD = 1.12 for DiSA and OD = 1.03 for DrSA) were established by calculating the mean value + 3 standard deviations (3SD) of 30 serum samples obtained from clinically healthy humans (negative controls).

Laboratory investigations on the patients' dogs

The dogs were screened for *Dirofilaria* spp. microfilariae using the Knott test (only 1 mL of EDTA blood) (Genchi *et al.*, 2021). Additionally, the Petcheck Heartworm Canine (IDEXX) test was used according to the manufacturer's instructions, to detect *D. immitis* antigen. All dogs were also screened for antibodies against *D. repens* (Mendoza-Roldán *et al.*, 2021).

Results

Histology – case 1 and case 2

In the first case, the nematode displayed a whitish, cylindrical body measuring 14 cm in length. Histological characteristics were consistent with an immature female nematode




Figure 3. Histological analysis case 1. (A,B) Longitudinal section of the parasite showing a thick, intensely eosinophilic cuticle covered with several layers of chitin raised in ridges (red arrows) (E.E. 40×); (C) longitudinal section of the parasite: caudal end. (E.E. 20×). (D) Longitudinal section of the parasite, coelomic cavity: a monolayer of cubic or cylindrical cells is observed (compatible with the intestine, red arrows). No microfilariae are observed. (E.E. 40×); (E,F) nodule. Cutis sections: the stratum corneum epithelium (blue arrow) is diffusely hyperplastic. The dermis is extensively expanded due to the presence of a chronic inflammatory infiltrate, mainly characterised by a high number of activated macrophages, epithelioid cells and eosinophilic granulocytes and a lower number of small mature lymphocytes and plasma cells (yellow arrows). There are multifocal areas of repair characterised by the presence of blood vessels (green arrows), fibroblasts associated with fibrillar matrix (fibroplasia) (E.E. 20×).

(Wong and Brummer, 1978; Brindicci *et al.*, 2019). Additionally, a thick, intensely eosinophilic cuticle was observed, covered with multiple chitinous layers forming ridges. This ridged appearance of the cuticle was characteristic of *D. repens* (Figure 3A,B). The nematode's coelomic cavity contained a monolayer of cuboidal or cylindrical cells, consistent with intestinal tissue. No microfilariae were observed in the uterus (Figure 3C,D). Macroscopically, the nodule measured 1 × 1 cm, was spheroidal in shape, firm in consistency, and smooth on the surface. Histologically, the outer skin epithelium of the nodule was diffusely hyperplastic. The dermis was significantly expanded due to chronic inflammatory infiltrate, mainly comprising numerous activated macrophages, epithelioid cells, and eosinophilic granulocytes, with a smaller number of mature lymphocytes and plasma cells (Figure 3E,F).

For the second case, the nematode displayed a whitish, cylindrical body and with ridged appearance of the cuticle measuring 15–16 cm in length. Histological characteristics were consistent with a fertile female nematode (Ciucă *et al.*, 2020; Simón *et al.*, 2022), showing in the transverse section the uterus with microfilariae (Figure 4A,B).

Knott test and multiplex PCR (case 1 and case 2)

The modified Knott's test revealed the absence of microfilariae in the blood of both patients.

The multiplex PCR method amplified a 484-base pair fragment specific to *D. repens* in both patients. The sequences of the PCR products obtained are available in the GenBank database under the following accession numbers: AY693808 (Case 1) and MW242631.1 (Case 2).

The first patient (Case 1) showed a positive response to *D. repens* somatic antigens (OD = 1.24) and a negative response to *D. immitis* (OD = 0.64) somatic antigens as determined by in-house IgG.

Similarly, the second patient (Case 2) was seropositive for *D. repens* (OD = 1.86) while no specific IgG were detected against *D. immitis* antigens (OD = 0.63).

Test results for *Dirofilaria* spp in dogs

All dogs tested negative for *Dirofilaria* infection in both the modified Knott's test and the antigen test. In addition, the ELISA IgG assay for antibodies against *D. repens* also yielded negative results for the three dogs tested.

Discussion

The present study unveiled two new cases of human *Dirofilaria repens* infection in southern Italy and involved clinical, serological, haematological and molecular analysis in the two patients and their dogs, adopting an integrated medical and veterinary approach.




Figure 4. Histological analysis case 2. (A,B) Longitudinal section of the parasite: intestine (black arrow) and uterus containing microfilariae (blue arrow) (E.E. 20×). Transverse section of the parasite: a thick, intensely eosinophilic cuticle is observed, covered by multiple layers of chitin raised into crests (red arrows), intestine (black arrow), and uterus (blue arrow) with microfilariae (green arrow) (E.E. 20×).

Dirofilaria repens is the main etiological agent of human dirofilariasis in the Old World (Genchi and Kramer, 2020). The rise in dirofilariasis cases across Europe has been linked to factors such as climate change, an increase in mosquito species capable of transmitting *Dirofilaria* spp., and the intensification of international dog trade and movement (Cancrini *et al.*, 2007; Genchi *et al.*, 2009; Capelli *et al.*, 2013, 2018; Salamatin *et al.*, 2013; Simón *et al.*, 2017; Perles *et al.*, 2024). In Ukraine, for instance, dirofilariasis was included in the national surveillance system for notifiable diseases in 1975, and the country has thousands of cases reported in the archives of the Ministry of Health, even if not all cases are documented in international scientific journals (Salamatin *et al.*, 2013; ESDA, 2017; Simón *et al.*, 2017, 2022).

Human dirofilariasis cases have been reported mainly from canine dirofilariasis endemic areas in the Mediterranean region (Capelli *et al.*, 2018; Tahir *et al.*, 2019), with Italy being the country with the highest number of reported cases to date (Pampiglione *et al.*, 2001; Capelli *et al.*, 2018; Gabrielli *et al.*, 2021; Simon *et al.*, 2022, 2024). Considering that Italy is an endemic country for canine and human dirofilariasis (Capelli *et al.*, 2018; Genchi *et al.*, 2019, 2023; Brianti *et al.*, 2022; Napoli *et al.*, 2023) not only is an increased number of cases of *D. immitis* and *D. repens* in both dogs and humans expected, but also a high frequency of aberrant migrations in humans, especially due to *D. repens*, as already shown in dogs (Pierantozzi *et al.*, 2017; Napoli *et al.*, 2024). As an example, *D. repens* infections of the male genitalia have been documented in humans (Pampiglione and Rivasi, 2000; Simón *et al.*, 2022) including a case of testicular infection with *D. repens* recently reported in a child from northeastern Italy (Ugolini *et al.*, 2022). Moreover, it should also be noted that two cases of *D. immitis* infection in humans, have already been reported in central Italy, one with localization in the right temporal bulbar subconjunctival space and the other with a 2 cm nodule in the right upper lung (Avellis *et al.*, 2011; Palicelli *et al.*, 2022).

The two human cases of *D. repens* infection reported in this study provide further insight into the epidemiological and clinical features of human dirofilariasis in an endemic region of southern Italy, where canine dirofilariasis is prevalent (Ferrara *et al.*, 2022a; Ciucă *et al.*, 2023; Napoli *et al.*, 2023). Indeed, human dirofilariasis cases in southern Italy have been progressively increasing. The earliest reported cases date back to 1885, with 11 cases documented (Pampiglione *et al.*, 2001). After a long hiatus, a new case emerged in an hospital in Naples in 2012: a 17-year-old male who presented with a pulmonary nodule that was initially suspected to be a lung tumour; histopathological examination confirmed the presence of *D. repens*, highlighting the diagnostic challenges posed by the atypical pulmonary location mimic malignancies (Gabrielli *et al.*, 2012). In 2018, Ferrari *et al.* reported a rare case of pulmonary dirofilariasis caused by *D. repens* in a 62-year-old man from Palermo,

who presented with pulmonary nodules that were initially suspected to be malignant based on imaging studies. Subsequently, in 2019, a case was reported in Bari, involving an 82-year-old woman with a subcutaneous *D. repens* lesion on her right thigh (Brindicci *et al.*, 2019). Moreover, in 2022, it was reported a case of cutaneous dirofilariasis caused by *D. repens* in an Italian girl from Naples (southern Italy) who spends her summer holidays in the Tuscan countryside (northern Italy) (Ferrara *et al.*, 2022b).

Our study describes two additional autochthonous cases of subcutaneous *D. repens* infections in southern Italy: a 33-year-old man from Caserta (Case 1), and a 67-year-old man from Pozzuoli, a coastal town in the Metropolitan City of Naples (Case 2). These cases evidence the continuous spreading of human dirofilariasis in southern Italy and the importance of accurate diagnostic approaches. Moreover this findings, alongside our cases of subcutaneous localization, reflects the diverse manifestations of *D. repens*, demonstrating its potential for internal organ involvement, albeit rarely, as also documented in literature (Ro *et al.*, 1989; Ferrari *et al.*, 2018; Oliva *et al.*, 2019). Together, these cases highlight the need for awareness of atypical presentations of dirofilariasis in endemic regions, where superficial and deep tissue nodules may both occur.

Genetic analysis has shown considerable variability among isolates from both humans and dogs in Italy, indicating potential intraspecific diversity that may influence transmission dynamics and clinical manifestations (Genchi and Kramer, 2017). This genetic diversity could also explain the variety in human presentations, from superficial nodules to rare internal cases. Such variability underscores the importance of molecular diagnostics in distinguishing between different strains, as it may have implications for tracking infection sources and understanding regional differences in transmission and pathogenicity (Gabrielli *et al.*, 2024). However, a major limitation of the present study is that the genetic variability of the strains in both cases was not characterized.

The distribution of patients by age and sex is not uniform. Most reported cases of *D. repens* infection occur in individuals aged 20–69, with the highest incidence observed in those aged 50–59 years and a greater prevalence in women than in men (Salamatin *et al.*, 2013; Gabrielli *et al.*, 2021; Simón *et al.*, 2022). In the current study, one patient (Case 2) falls within the age range with the highest incidence, while the other patient (Case 1) is within the broader 20–69 age range. Moreover, both patients were male, aligning with findings from other studies (Pampiglione *et al.*, 1994; PupiĆ-Bakrač *et al.*, 2021) and contrasting with the cases reviewed by Simón *et al.* (2022). Several studies have indicated that there is no statistically significant correlation between human subcutaneous dirofilariasis and either age or gender (Pampiglione *et al.*, 2001; Fontes-Sousa *et al.*, 2019; PupiĆ-Bakrač *et al.*, 2021). Nevertheless, geographic variations in age distributions have been

noted. For example, However, while cases in children are rare in Italy (Bertozzi et al., 2015; Pansini et al., 2022; Ugolini et al., 2022), in Sri Lanka it is common to find infected children under 5 years of age – possibly due to differences in clothing, customs, and vector preferences (Senanayake et al., 2013; Balendran et al., 2022). Additionally, a case in a 100-year-old individual is also unusual in Italy, perhaps reflecting the reduced effectiveness of insecticides in the Piedmont area (Pampiglione et al., 2001). Ultimately, the risk factors for this disease extend beyond age and gender and include the endemicity of the region (evidenced by infections in dogs and cats), areas with increased vector exposure, and lifestyle factors, such as herein presented, where both patients frequently took long walks in areas known to have canine cases. Other factors, such as individual immunological status, may also contribute, as suggested by other studies (Pupić-Bakrač et al., 2021; Simón et al., 2022).

Almost all human cases of *D. repens* reported in Italy have been asymptomatic or have presented with transient localized symptoms such as swelling, erythema, rash and pruritus (Rivasi et al., 2006; Ermakova et al., 2017; Gabrielli et al., 2021; Pupić-Bakrač et al., 2021). These symptoms are usually associated with subcutaneous or submucosal nodules in various parts of the body and can even involve the eye through the conjunctiva (Pampiglione et al., 1995, 2001; Pampiglione and Rivasi, 2000; Eccher et al., 2008; Palicelli et al., 2014; Fontanelli Sulekova et al., 2016). Only rarely the worm can migrate to deeper tissues such as muscles and lungs (Ermakova et al., 2017; Gabrielli et al., 2021; Pupić-Bakrač et al., 2021; Simón et al., 2022; Napolitano et al., 2023). The two cases in our study align with typical human presentations with subcutaneous nodules without blood microfilariae (Pampiglione and Rivasi, 2000; Pampiglione et al., 2001; Gabrielli et al., 2021). In Case 1, the patient presented with a subcutaneous mass in the upper eyelid, while Case 2 presented with an oval-shaped nodule in the left frontal region of the scalp. Furthermore, the nematode's presence in the subcutaneous tissue of both cases 'resembles *D. repens*', as evidenced by similar cases from Italy (Brindicci et al., 2019; Gabrielli et al., 2021; Campana et al., 2022). Both cases in the present study showed infection sites similar to those commonly reported in a Ukrainian study, which analysed clinical data from 755 cases and found that 64.6% (488) of parasitic lesions were located in the head, including 297 cases around the eyes (Salamatin et al., 2013). Additionally, the eyelid is the most commonly reported anatomical site for nodules caused by *D. repens*, as highlighted in the recent case review by Simón et al. (2022). This prevalence is likely because parasitic lesions in other areas are more difficult for patients to detect.

Humans are considered accidental hosts for *Dirofilaria* spp., as nematodes rarely reach sexual maturity or produce microfilariae in human blood (Simón et al., 2012; Ermakova et al., 2017). However, some studies have documented *D. repens* infections with microfilariae in blood or local tissues (Kłudkowska et al., 2018; Potters et al., 2018; Pupić-Bakrač et al., 2021; Tasić-Otasevic et al., 2023).

In our study, the first patient presented with an immature adult female *D. repens* worm measuring 14 cm in length, lacking microfilariae in both the uterus and the blood, but exhibiting the presence of antibodies in the serum sample. Similar characteristics – including sexually immature nematodes, absence of microfilariae, and a positive IgG immune response – were reported in some of the cases described in central Italy (Gabrielli et al., 2021). The second case was initially suspected to be a cyst or fibroma based on ultrasound findings but was later identified as a live 16–17 cm *D. repens* worm upon surgical excision, highlighting the challenge of distinguishing dirofilariasis from benign tumours. A similar case of subcutaneous

D. repens infection in the scalp region of a patient, was also reported in Slovenia in 2017 (Kotnik et al., 2022).

Furthermore, in both cases of the present study, the worms were encapsulated within nodules. This finding is consistent with the global cases reviewed by Simón et al. (2022), who reported that 96.68% (408) of *D. repens* worms were encapsulated within nodules, while only 3.32% (14) were free.

An intriguing aspect of our study was the discovery of a gravid female *D. repens* within the nodule of the second case, although no microfilariae were present in the blood and no male worm was detected. While most nematodes isolated from humans are immature, several studies have documented mature *D. repens* females carrying microfilariae in their uterus without releasing them into the bloodstream (Supriaga et al., 2004; Poppert et al., 2009; Makaveev et al., 2022; Simón et al., 2022). Furthermore, Ermakova et al. (2017) reported sexually mature *D. repens* in 10.4% of nodules, suggesting that humans may not be a 'dead-end' host for this helminth. Even though the detection of *D. repens* microfilariae in the bloodstream suggests the probable presence of both male and female adult worms within the human host, the conclusive identification of both sexes in human cases has yet to be confirmed. Notably, the present case is the first molecularly confirmed instance in Italy since 1992 of a gravid female *D. repens* without the presence of microfilariae in either the peripheral blood or local tissues. In the case documented by Pampiglione et al. (1992), a 53-year-old woman from the Campania region (Salerno province) had a parasite located in the left submammary region. The detection of microfilariae in the uterus suggested that she was also harbouring a mature male worm, although none was identified. This finding implied the potential for microfilariae release into the bloodstream, yet microfilaremia was not observed. Conversely, Fontanelli Sulekova et al. (2016) reported a case in Central Italy where microfilariae were detected in a fine-needle aspirate of a subcutaneous nodule, despite the absence of microfilaremia in peripheral blood samples. The infection did not become fully patent, indicating that while microfilariae can be present in local tissues, they may not always enter systemic circulation or lead to a fully developed infection.

These cases highlight the variability in human *D. repens* infections and suggest that the parasite's life cycle in humans may be more complex than previously thought. The presence of gravid females and tissue microfilariae without corresponding microfilaremia challenges the notion of humans as strict dead-end hosts and underscores the need for further research to understand the transmission dynamics and host–parasite interactions of *D. repens* in human infections.

A recent study reviewing *D. repens* cases in humans with local microfilaremia or blood microfilaremia in Europe from 1992 to 2021 identified 20 patients (Pupić-Bakrač et al., 2021). Only four of these patients had a history of chronic immune disorders, while the others were either immunocompetent or had no relevant medical data available. In our study, the two patients had no history of chronic immune disorders or immunodeficiency and exhibited haematological and biochemical parameters within normal ranges. In contrast, cases of *D. repens* with microfilariae in the bloodstream have been associated with the development of eosinophilia (Simón et al., 2012; Pupić-Bakrač et al., 2021).

The diagnosis of human dirofilariasis can be challenging and difficult to achieve. Most cases present with 'silent signs', absence of microfilariae, or are misinterpreted as malignant tumours (Capelli et al., 2018; Ferrari et al., 2018; Miterpáková et al., 2022; Simón et al., 2022). Diagnosis is primarily based on histopathological

examination of excised nodules (Miterpáková *et al.*, 2022; Simón *et al.*, 2022). In most cases, similar to our cases, the nematodes are identified by the typical features of striated cuticle for *D. repens* and smooth for *D. immitis* (with ridges and striae only on the ventral surface of the male caudal extremity) (Wong and Brummer, 1978; Simón *et al.*, 2012). However, due to the presence of immature nematodes in the histology of nodules, it is sometimes difficult to distinguish them from fully developed adults at the species level. Thus, molecular analysis is a better tool to diagnose *Dirofilaria* spp. in humans, but also to understand the hidden diversity of *Dirofilaria* spp. (Rossi *et al.*, 2015; Palicelli *et al.*, 2022).

Current serological tests for human dirofilariasis have limitations in sensitivity and specificity. Commercial ELISA kits, such as those from Bordier Affinity Products SA, detect antibodies against multiple filarial genera, including *Dirofilaria*, while home-made IgG ELISA tests target *D. immitis* and *D. repens*. These limitations are exemplified by the low OD values observed in our cases, possibly due to the immature worm in the first case. Despite this, serological tests can serve as preliminary epidemiological screening tools to assess human exposure risk, especially in regions endemic for canine dirofilariasis. Human immunological responses to *Dirofilaria* spp. correlate with high dirofilariasis prevalence in canine populations (Ciucă *et al.*, 2018; Genchi and Kramer, 2020; Mendoza-Roldán *et al.*, 2021; Brianti *et al.*, 2022; Perles *et al.*, 2024).

A persistent challenge in the diagnosis of human dirofilariasis, despite its worldwide endemic status, is the difficulty many physicians face in recognizing clinical cases. This lack of awareness often leads to delays in diagnosis as physicians do not initially consider dirofilariasis. Such diagnostic delays are noted in numerous reports not only in Italy but worldwide, where patients struggle with the infection for a long time and often require guidance and support from specialists in parasitology to obtain a diagnosis (Pampiglione and Rivasi, 2000; Ilyasov *et al.*, 2013; Ahmed *et al.*, 2022; Miterpáková *et al.*, 2022; Simón *et al.*, 2022). For example, in Case 1 from our study, the patient was treated with ivermectin and albendazole for one week after the initial evaluation and subsequently referred to another hospital for further assessment – without excising the nodule as a primary intervention or recognizing dirofilariasis as a possible cause. Similarly, in Case 2, the nodule was initially suspected to be cystic in nature. Consequently, the classical presentation of *D. repens* infection went unrecognized at first, even though Italy is an endemic area for dirofilariasis in both dogs and humans. In southern Italy in particular, the prevalence of both *D. repens* and *D. immitis* is increasing, with prevalence in shelter dogs ranging from 10% to 75% (Ferrara *et al.*, 2022a; Ciucă *et al.*, 2023; Napoli *et al.*, 2023).

However, it should be noted that the dogs owned by both patients in this study tested negative for *D. repens* microfilariae. This finding indicates that the infections in humans were not directly linked to their own pets, despite close contact with potential reservoir hosts highlights the complexity of the transmission dynamics and the importance of considering the wider dog population in endemic areas as well as environmental exposure to the mosquito vectors such as *Aedes albopictus* and *Culex pipiens* (Capelli *et al.*, 2018; Ferrara *et al.*, 2022a).

In contrast to our study, a case of *D. repens* in subcutaneous tissue was described in Croatia in which the patients' own dogs were microfilaraemic (Pupić-Bakrač *et al.*, 2021). This highlights the variability of zoonotic transmission dynamics and underlines the importance of comprehensive diagnostic approaches in both humans and their pets.

In conclusion, Italy is facing increasing prevalence rates of dirofilariasis in dogs (Capelli *et al.*, 2013; Genchi and Kramer, 2020; Brianti *et al.*, 2022; Ciucă *et al.*, 2023; Napoli *et al.*, 2023) and a steady increase incidence in humans, with the highest number of reported cases in Europe (Pampiglione *et al.*, 2001; Rivasi *et al.*, 2006; Gabrielli *et al.*, 2021; Mendoza-Roldán *et al.*, 2021; Pupić-Bakrač *et al.*, 2021; Palicelli *et al.*, 2022; Perles *et al.*, 2024). Therefore, a comprehensive and harmonized epidemiological study of human exposure to *Dirofilaria* spp. in Italy would be advisable to demonstrate the need for its inclusion in national disease surveillance and to better monitor the increasing number of zoonotic cases.

Supplementary material. The supplementary material for this article can be found at <https://doi.org/10.1017/S0031182025000290>.

Acknowledgements. We thank the medical doctors for providing us with details regarding the clinical aspects of the cases presented in this study.

Author contributions. LC and LR conceived and designed the study. LC, PF, MVP, MA, and LP collected clinical data and patient samples. SG, MOM, MPM, and LC performed molecular and immunological analyses. EN and OP conducted histological examinations and pathological assessments. LC analysed the data and drafted the manuscript. SG, OP, MPM, PF, MOM, and LP contributed to data interpretation and critically revised the manuscript. LR and SG supervised the study and provided critical revisions. All authors read and approved the final manuscript.

Financial support. This study was supported by EU funding within the NextGeneration EU-MUR PNRR Extended Partnership initiative on Emerging Infectious Diseases (Project no. PE00000007, INF-ACT).

Competing interests. The authors declare there are no conflicts of interest.

Ethical standards. This study was conducted in accordance with the ethical standards of the institutional and national research committees and with the 1964 Helsinki Declaration and its later amendments. Ethical approval for this study was obtained from the Ethics Committee of the University of Naples Federico II under approval number 0010200. All procedures performed involving human participants were in accordance with the ethical standards of the institutional and national research committees. Oral informed consent was obtained from all individual participants included in the study. Participants were provided with detailed information about the study's objectives, procedures, potential risks and benefits and assured of their right to withdraw at any time without any consequences.

References

- Addario C (1885) Su di un nematode dell'occhio umano. *Annali Di Oftalmologia* 14, 135–148.
- Ahmed N, Tonelli L, Labagnara G, Doglioni C and Pedica F (2022) Subcutaneous dirofilariasis in Italy: A diagnostic hypothesis to remind when the anamnesis is misleading. *Pathologica* 114, 174–177. doi: 10.32074/1591-951X-297
- Avellis FO, Kramer LH, Mora P, Bartolino A, Benedetti P and Rivasi F (2011) A case of human conjunctival dirofilariasis by *Dirofilaria immitis* in Italy. *Vector-Borne and Zoonotic Diseases* 11, 451–452. doi: 10.1089/vbz.2010.0067
- Awe RJ, Mattox KL, Alvarez BA, Stork WJ, Estrada R and Greenberg SD (1975) Solitary and bilateral pulmonary nodules due to *Dirofilaria immitis*. *American Review of Respiratory Disease* 112, 445–449. doi: 10.1164/arrd.1975.112.3.445
- Balendran T, Yatawara L and Wickramasinghe S (2022) Human Dirofilariasis caused by *Dirofilaria repens* in Sri Lanka from 1962 to 2020. *Acta Parasitologica* 67, 628–639. doi: 10.1007/s11686-022-00543-3
- Benzaquen M, Brajon D, Delord M, Yin N, Bittar F, Toga I, Berbis P and Parola P (2015) Cutaneous and pulmonary dirofilariasis due to *Dirofilaria repens*. *British Journal of Dermatology* 173, 788–791. doi: 10.1111/bjd.13859

- Bertozzi M, Rinaldi VE, Prestipino M, Giovenali P and Appignani A (2015) Dirofilariasis Mimicking an Acute Scrotum. *Pediatric Emergency Care* **31**, 715–716. doi: 10.1097/PEC.0000000000000346
- Bradford MM (1976) A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Analytical Biochemistry* **7**, 248–254. doi: 10.1016/0003-2697(76)90527-3
- Brianti E, Panarese R, Napoli E, De Benedetto G, Gaglio G, Bezerra-Santos MA, Mendoza-Roldan JA and Otranto D (2022) *Dirofilaria immitis* infection in the Pelagie archipelago: The southernmost hyperendemic focus in Europe. *Transboundary and Emerging Diseases* **69**, 1274–1280. doi: 10.1111/tbed.14089
- Brindicci G, Santoro CR, Signorile F, Leone A, Di Ciaula G, Monno L and Angarano G (2019) Subcutaneous human dirofilariosis by *D. repens* in South Italy: A case report. *New Microbiologica* **42**, 234–236.
- Cabrera ED, Carretón E, Morchón R, Falcón-Cordón Y, Falcón-Cordón S, Simón F and Montoya-Alonso JA (2018) The Canary Islands as a model of risk of pulmonary dirofilariasis in a hyperendemic area. *Parasitology Research* **117**, 933–936. doi: 10.1007/s00436-018-5774-1
- Campana I, Fania L, Samela T, Campana L, Zecchi V, Ricci F and Abeni D (2022) A case of human *Dirofilaria repens* in Rome, Italy: A clinical and radiological challenge. *Dermatology Reports* **14**, 9354. doi: 10.4081/dr.2022.9354
- Cancrini G, Scaramozzino P, Gabrielli S, Di Paolo M, Toma L and Romi R (2007) *Aedes albopictus* and *Culex pipiens* implicated as natural vectors of *Dirofilaria repens* in central Italy. *Journal of Medical Entomology* **44**, 1064–1066. doi: 10.1603/0022-2585(2007)44[1064:aaacpi]2.0.co;2
- Capelli G, Frangipane Di Regalbano A, Simonato G, Cassini R, Cazzin S, Cancrini G, Otranto D and Pietrobello M (2013) Risk of canine and human exposure to *Dirofilaria immitis* infected mosquitoes in endemic areas of Italy. *Parasites and Vectors* **6**, 60. doi: 10.1186/1756-3305-6-60
- Capelli G, Genchi C, Baneth G, Bourdeau P, Brianti E, Cardoso L, Danesi P, Fuehrer HP, Giannelli A, Ionică AM, Maia C, Modrý D, Montarsi F, Krücken J, Papadopoulos E, Petric D, Pfeffer M, Savić S, Otranto D, Poppert S and Silaghi C (2018) Recent advances on *Dirofilaria repens* in dogs and humans in Europe. *Parasites and Vectors* **11**, 663. doi: 10.1186/s13071-018-3205-x
- Ciucă L, Caruso V, Illiano S, Bosco A, Maurelli MP and Rinaldi L (2023) Corrigendum: Emerging risk of *Dirofilaria* spp. infection in shelter dogs in southern Italy. *Frontiers in Veterinary Science* **10**, 1290005. doi: 10.3389/fvets.2023.1290005
- Ciucă L, Roman C, Prisco F, Miron L, Acatrinei D, Paciello O, Maurelli MP, Vismarra A, Cringoli G and Rinaldi L (2020) First report of *Dirofilaria repens* infection in a microfilaremic cat from Romania. *Veterinary Parasitology: Regional Studies and Reports* **22**, 100497. doi: 10.1016/j.vprsr.2020.100497
- Ciucă L, Simón F, Rinaldi L, Kramer L, Genchi M, Cringoli G, Acatrinei D, Miron L and Morchón R (2018) Seroepidemiological survey of human exposure to *Dirofilaria* spp. in Romania and Moldova. *Acta Tropica* **187**, 169–174. doi: 10.1016/j.actatropica.2018.07.012
- Eccher A, Dalfior D, Gobbo S, Martignoni G, Brunelli M, Decaminada W, Bonetti F, Rivasi F, Barbareschi M and Menestrina F (2008) Periorbital subcutaneous tumor-like lesion due to *Dirofilaria repens*. *International Journal of Surgical Pathology* **16**, 101–103. doi: 10.1177/1066896907307040.z
- Ermakova L, Nagorny S, Pshenichnaya N, Ambalov Y and Boltachiev K (2017) Clinical and laboratory features of human dirofilariasis in Russia. *IDCases* **19**, 112–115. doi: 10.1016/j.idcr.2017.07.006
- European Society of Dirofilariasis and Angiostrongylosis (ESDA) (2017) Guidelines for clinical management of human *Dirofilaria* infections. Available at: <https://www.esda.vet/guidelines-tutorials/> (Accessed 1 March 2021).
- Ferrara D, Barbieri A, D'Auria D, Castelli L, Gholami Shanagolabad P, Zeccolini M and Esposito F (2022b) The worm “repens” in the city: A case of subcutaneous dirofilariasis in a 12-year-old female. *Journal of Radiology Review* **9**, 29–31. doi: 10.23736/S2723-9284.21.00160-6
- Ferrara M, Maglione R, Ciccarelli D, Mundis SJ, Di Loria A, Pisaroglo de Carvalho M and Santoro D (2022a) Prevalence of *Dirofilaria repens* in dogs living in deltaic coastal plain of the Volturno River (Italy): A geographical risk model of infection. *Journal of Helminthology* **96**, e12. doi: 10.1017/S0022149X22000062
- Ferrari PA, Grisolia A, Reale S, Liotta R, Mularoni A and Bertani A (2018) A rare case of human pulmonary dirofilariasis with nodules mimicking malignancy: Approach to diagnosis and treatment. *Journal of Cardiothoracic Surgery* **13**, 65. doi: 10.1186/s13019-018-0750-5
- Fontanelli Sulekova L, Gabrielli S, De Angelis M, Milardi GL, Magnani C, Di Marco B, Taliani G and Cancrini G (2016) *Dirofilaria repens* microfilariae from a human node fine-needle aspirate: A case report. *BMC Infectious Diseases* **16**, 248. doi: 10.1186/s12879-016-1582-3
- Fontes-Sousa AP, Silvestre-Ferreira AC, Carretón E, Esteves-Guimarães J, Maia-Rocha C, Oliveira P, Lobo L, Morchón R, Araújo F, Simón F and Montoya-Alonso JA (2019) Exposure of humans to the zoonotic nematode *Dirofilaria immitis* in Northern Portugal. *Epidemiology & Infection* **30**, 147–282. doi: 10.1017/S0950268819001687
- Gabrielli S, Brustenga L, Morganti G, Ciucă L, Barlozzari G, Rigamonti G, Orlandi M, Sforna M and Veronesi F (2024) Genetic variability of *Dirofilaria repens* isolates from humans and dogs in Italy. *Veterinary Parasitology* **326**, 110096. doi: 10.1016/j.vetpar.2023.11009
- Gabrielli S, Mangano V, Furzi F, Oliva A, Vita S, Poscia R, Fazii P, Di Paolo J, Marocco R, Mastroianni CM, Bruschi F and Mattiucci S (2021) Molecular identification of new cases of human dirofilariosis (*Dirofilaria repens*) in Italy. *Pathogens* **10**, 251. doi: 10.3390/pathogens10020251
- Gabrielli S, Petrucci L, Coppola MG and Cancrini G (2012) A Case of Human Pulmonary Dirofilariasis Attributed to *D. Repens* in Italy. Parma, Italy: Third European Dirofilaria Days.
- Genchi C and Kramer LH (2017) Subcutaneous dirofilariosis (*Dirofilaria repens*): An infection spreading throughout the Old World. *Parasites and Vectors* **10**(Suppl. 2), 517. doi: 10.1186/s13071-017-2434-8
- Genchi C and Kramer LH (2020) The prevalence of *Dirofilaria immitis* and *D. repens* in the Old World. *Veterinary Parasitology* **280**, 108995. doi: 10.1016/j.vetpar.2019.108995
- Genchi C, Rinaldi L, Mortarino M, Genchi M and Cringoli G (2009) Climate and *Dirofilaria* infection in Europe. *Veterinary Parasitology* **163**, 286–292. doi: 10.1016/j.vetpar.2009.03.026
- Genchi M, Ciucă L, Vismarra A, Ciccone E, Cringoli G, Kramer L and Rinaldi L (2021) Evaluation of alternative reagents on the performance of the modified Knott's test. *Veterinary Parasitology* **298**, 109555. doi: 10.1016/j.vetpar.2021.109555
- Genchi M, Kramer L, Venco L, Ciucă L and Vismarra A (2023) How Italian practitioners manage dirofilariosis in dogs and cats? Data of a second national survey. *Veterinary Parasitology: Regional Studies and Reports* **46**, 100934. doi: 10.1016/j.vprsr.2023.100934
- Genchi M, Rinaldi L, Venco L, Cringoli G, Vismarra A and Kramer L (2019) *Dirofilaria immitis* and *D. repens* in dog and cat: A questionnaire study in Italy. *Veterinary Parasitology* **267**, 26–31. doi: 10.1016/j.vetpar.2019.01.014
- Grandi G, Morchón R, Kramer L, Kartashev V and Simón F (2008) *Wolbachia* in *Dirofilaria repens*, an agent causing human subcutaneous dirofilariasis. *Journal of Parasitology* **94**, 1421–1423. doi: 10.1645/GE-1575.1
- Ilyasov B, Kartashev V, Bastrikov N, Morchón R, González-Miguel J and Simón F (2013) Delayed diagnosis of dirofilariasis and complex ocular surgery, Russia. *Emerging Infectious Diseases* **19**, 326–328. doi: 10.3201/eid1902.121388
- Kłudkowska M, Pielok Ł, Frąckowiak K, Masny A, Gołab E and Paul M (2018) *Dirofilaria repens* infection as a cause of intensive peripheral microfilaremia in a Polish patient: Process description and cases review. *Acta Parasitologica* **63**, 657–663. doi: 10.1515/ap-2018-0077
- Kotnik T, Rataj AV and Šoba B (2022) *Dirofilaria repens* in dogs and humans in Slovenia. *Journal of Veterinary Research* **66**, 117–123. doi: 10.2478/jvetres-2022-0008
- Makaveev I, Galey A, Marinova P, Martinov M, Harizanov R, Rainova I and Karcheva M (2022) First case of ocular dirofilariosis in Bulgaria caused by gravid female *Dirofilaria repens*. *Annals of Parasitology* **68**, 191–193. doi: 10.17420/ap6801.424
- McCall JW, Genchi C, Kramer LH, Guerrero J and Venco L (2008) Heartworm disease in animals and humans. *Advances in Parasitology* **66**, 193–285. doi: 10.1016/S0065-308X(08)00204-2

- Mendoza-Roldán JA, Gabrielli S, Cascio A, Manoj RRS, Bezerra-Santos MA, Benelli G, Brianti E, Latrofa MS and Otranto D (2021) Zoonotic *Dirofilaria immitis* and *Dirofilaria repens* infection in humans and an integrative approach to the diagnosis. *Acta Tropica* **223**, 106083. doi: 10.1016/j.actatropica.2021.106083
- Miterpáková M, Antolová D, Rampalová J, Undesser M, Krajčovič T and Vichová B (2022) *Dirofilaria immitis* pulmonary dirofilariasis, Slovakia. *Emerging Infectious Diseases* **28**, 482–485. doi: 10.3201/eid2802.211963
- Momčilović S, Gabrielli S, Đenić N, Živković N, Stevanović G, Krstić M, Randelović M and Tasić-Otašević S (2022) New cases of human dirofilariasis on the Balkan Peninsula—“Masked intruders” uncovered by a surgeon. *Parasitology International* **86**, 102482. doi: 10.1016/j.parint.2021.102482
- Montoya-Alonso JA, Mellado I, Carretón E, Cabrera-Pedrero ED, Morchón R and Simón F (2010) Canine dirofilariasis caused by *Dirofilaria immitis* is a risk factor for the human population on the island of Gran Canaria, Canary Islands, Spain. *Parasitology Research* **107**, 1265–1269. doi: 10.1007/s00436-010-1987-7
- Muro A, Genchi C, Cordero M and Simón F (1999) Human dirofilariasis in the European Union. *Parasitology Today* **15**, 386–389. doi: 10.1016/s0169-4758(99)01496-9
- Napoli E, De Benedetto G, Ciucă L, Bosco A, Lia RP, Veneziano V, Bezerra Santos MA, Otranto D, Rinaldi L and Brianti E (2023) New distribution patterns of *Dirofilaria immitis* in Italy. *Frontiers in Veterinary Science* **10**, 1162403. doi: 10.3389/fvets.2023.1162403
- Napoli E, De Benedetto G, Di Giorgio S, Sfacteria A, Marino G, Mazzara D, Giambrone G, Gaglio G and Brianti E (2024) How frequently *Dirofilaria repens* localizes in dog testes? *Veterinary Parasitology: Regional Studies and Reports* **55**, 101112. doi: 10.1016/j.vprsr.2024.101112
- Napolitano AG, Pallotto C, Pourmolkara D, Vinci D and Coviello E (2023) *Dirofilaria* infestation of pectoralis major muscle in Italy, a case report. *Clinica Terapeutica* **174**, 322–325. doi: 10.7417/CT.2023.2544
- Oliva A, Gabrielli S, Pernazza A, Pagini A, Daralioi T, Mantovani S, Mattiucci S, D'Amati G and Mastroianni CM (2019) *Dirofilaria repens* infection mimicking lung melanoma metastasis. *Open Forum Infectious Diseases* **6**, ofz049. doi: 10.1093/ofid/ofz049
- Palicelli A, Deambrogio C, Arnulfo A, Rivasi F, Paganotti A and Boldorini R (2014) *Dirofilaria repens* mimicking an ovarian mass: Histologic and molecular diagnosis. *APMIS* **122**, 1045–1046. doi: 10.1111/apm.12259
- Palicelli A, Veggiani C, Rivasi F, Gustinelli A and Boldorini R (2022) Human pulmonary dirofilariasis due to *Dirofilaria immitis*: The first Italian case confirmed by polymerase chain reaction analysis, with a systematic literature review. *Life* **12**, 1584. doi: 10.3390/life12101584
- Pampiglione S, Canestri Trotti G and Rivasi F (1995) Human dirofilariasis due to *Dirofilaria (Nochtiella) repens*: A review of world literature. *Parassitologia* **37**, 149–193.
- Pampiglione S, Del Maschio O, Pagan V and Rivasi F (1994) Pulmonary dirofilariasis in man: A new Italian case. Review of the European literature. *Parasite* **1**, 379–385. doi: 10.1051/parasite/1994014379
- Pampiglione S and Rivasi F (2000) Human dirofilariasis due to *Dirofilaria (Nochtiella) repens*: An update of world literature from 1995 to 2000. *Parassitologia* **42**, 231–254.
- Pampiglione S, Rivasi F, Angeli G, Boldorini R, Incensati RM, Pastormerlo M, Pavesi M and Ramponi A (2001) Dirofilariasis due to *Dirofilaria repens* in Italy, an emergent zoonosis: Report of 60 new cases. *Histopathology* **38**, 344–354. doi: 10.1046/j.1365-2559.2001.01099.x
- Pampiglione S, Rivasi F and Gustinelli A (2009) Dirofilarial human cases in the Old World, attributed to *Dirofilaria immitis*: A critical analysis. *Histopathology* **54**, 192–204. doi: 10.1111/j.1365-2559.2008.03197.x
- Pampiglione S, Schmid C and Montaperto C (1992) Human dirofilariasis: Discovery of a gravid female of *Dirofilaria repens* in a subcutaneous nodule. *Pathologica* **84**, 77–81.
- Panarese R, Iatta R, Beugnet F and Otranto D (2022) Incidence of *Dirofilaria immitis* and *Leishmania infantum* infections in sheltered dogs from Southern Italy. *Transboundary and Emerging Diseases* **69**, 891–894. doi: 10.1111/tbed.14025
- Pansini A, Magenes VC, Casini F, Guida G, De Sanctis M, Canonica CPM, Rossi RS, Zuccotti G and Giacomet V (2022) Testicular Dirofilariasis in an Italian 11-Year-old Child. *The Pediatric Infectious Disease Journal* **12**, e539–e540. doi: 10.1097/INF.00000000000003706
- Perles L, Dantas-Torres F, Krücken J, Morchón R, Walochnik J and Otranto D (2024) Zoonotic dirofilariases: One, no one, or more than one parasite. *Trends in Parasitology* **40**, 257–270. doi: 10.1016/j.pt.2023.12.007
- Pierantozzi M, Di Giulio G, Traversa D, Aste G and Di Cesare A (2017) Aberrant peritoneal localization of *Dirofilaria repens* in a dog. *Veterinary Parasitology: Regional Studies and Reports* **10**, 62–64. doi: 10.1016/j.vprsr.2017.08.002
- Poppert S, Hodapp M, Krueger A, Hegasy G, Niesen WD, Kern WV and Tannich E (2009) *Dirofilaria repens* infection and concomitant meningoencephalitis. *Emerging Infectious Diseases* **15**, 1844–1846.
- Potters I, Vanfraechem G and Bottieau E (2018) *Dirofilaria repens* nematode infection with microfilaremia in traveler returning to Belgium from Senegal. *Emerging Infectious Diseases* **24**, 1761–1763. doi: 10.3201/eid2409.180462
- Prieto C, Venco L, Simon F and Genchi C (1997) Feline heartworm (*Dirofilaria immitis*) infection: Detection of specific IgG for the diagnosis of occult infections. *Veterinary Parasitology* **70**, 209–217. doi: 10.1016/s0304-4017(97)00008-3
- Pupić-Bakrač A, Pupić-Bakrač J, Beck A, Jurković D, Polkinghorne A and Beck R (2021) *Dirofilaria repens* microfilaremia in humans: Case description and literature review. *One Health* **13**, 100306. doi: 10.1016/j.onehlt.2021.100306
- Rishniw M, Barr SC, Simpson KW, Frongillo MF, Franz M and Alpizar JL (2006) Discrimination between six species of canine microfilariiae by single polymerase chain reaction. *Veterinary Parasitology* **135**, 303–314. doi: 10.1016/j.vetpar.2005.10.013
- Rivasi F, Boldorini R, Criante P, Leutner M and Pampiglione S (2006) Detection of *Dirofilaria (Nochtiella) repens* DNA by polymerase chain reaction in embedded paraffin tissues from two human pulmonary locations. *APMIS* **114**, 567–574. doi: 10.1111/j.1600-0463.2006.apm_423.x
- Ro JY, Tsakalakis PJ, White VA, Luna MA, Chang-Tung EG, Green L, Cribb L and Ayala AG (1989) Pulmonary dirofilariasis: The great imitator of primary or metastatic lung tumor. A clinicopathologic analysis of seven cases and a review of the literature. *Human Pathology* **20**, 69–76. doi: 10.1016/0046-8177(89)90205-0
- Rossi A, Peix Á, Pavlikovskaya T, Sagach O, Nikolaenko S, Chizh N, Kartashev V, Simón F and Siles-Lucas M (2015) Genetic diversity of *Dirofilaria* spp. isolated from subcutaneous and ocular lesions of human patients in Ukraine. *Acta Tropica* **142**, 1–4. doi: 10.1016/j.actatropica.2014.10.021
- Salamatina RV, Pavlikovska TM, Sagach OS, Nikolayenko SM, Korniyushin VV, Kharchenko VO, Masny A, Cielecka D, Konieczna-Salamatina J, Conn DB and Golab E (2013) Human dirofilariasis due to *Dirofilaria repens* in Ukraine, an emergent zoonosis: Epidemiological report of 1465 cases. *Acta Parasitologica* **58**, 592–598. doi: 10.2478/s11686-013-0187-x
- Senanayake PM, Infaq MLM, Adikaram SGS and Udagama PV (2013) Ocular and subcutaneous dirofilariasis in a Sri Lankan infant: An environmental hazard caused by dogs and mosquitoes. *Paediatrics and International Child Health* **33**, 111–112. doi: 10.1179/2046905512Y.0000000024
- Sergiev VP, Supriaga VG, Morozov EN and Zhukova LA (2009) Human dirofilariasis: Diagnosis and the pattern of pathogen-host relations. *Medical Parasitology (Mosk)* **3**, 3–6.
- Simón F, Diosdado A, Siles-Lucas M, Kartashev V and González-Miguel J (2022) Human dirofilariasis in the 21st century: A scoping review of clinical cases reported in the literature. *Transboundary and Emerging Diseases* **69**, 2424–2439. doi: 10.1111/tbed.14210
- Simón F, González-Miguel J, Diosdado A, Gómez PJ, Morchón R and Kartashev V (2017) The complexity of zoonotic Filariasis epizootic and its consequences: A multidisciplinary view. *BioMed Research International*, 6436130. doi: 10.1155/2017/6436130
- Simón F, Siles-Lucas M, Morchón R, González-Miguel J, Mellado I, Carretón E and Montoya-Alonso JA (2012) Human and animal dirofilariasis: The emergence of a zoonotic mosaic. *Clinical Microbiology Reviews* **25**, 507–544. doi: 10.1128/CMR.00012-12

- Supriaga VG, Tsybina TN, Denisova TN, Morozov EN, Romanenko NA and Starkova TV** (2004) The first case of diagnosis of dirofilariasis from the microfilariae detected in the human subcutaneous tumor punctate. *Medical Parasitology (Mosk)* **6**, 6–8.
- Szénási Z, Kovács AH, Pampiglione S, Fioravanti ML, Kucsera I, Tónczos B and Tiszlavicz L** (2008) Human dirofilariasis in Hungary: An emerging zoonosis in central Europe. *Wiener Klinische Wochenschrift* **120**, 96–102. doi: 10.1007/s00508-008-0928-2
- Tahir D, Davoust B and Parola P** (2019) Vector-borne nematode diseases in pets and humans in the Mediterranean Basin: An update. *Veterinary World* **12**, 1630–1643. doi: 10.14202/vetworld.2019.1630-1643
- Tasić-Otasevic S, Golubović M, Trichei S, Zdravkovic D, Jordan R and Gabrielli S** (2023) Microfilaremic *Dirofilaria repens* Infection in Patient from Serbia. *Emerging Infectious Diseases* **12**, 2548–2550. doi: 10.3201/eid2912.230796
- Ugolini S, Lima M, Maffi M, Pierangeli F, Vastano M, Gargano T, Varani S, Gustinelli A, Caffara M and Fioravanti ML** (2022) *Dirofilaria repens* testicular infection in child, Italy. *Emerging Infectious Diseases* **28**, 2569–2572. doi: 10.3201/eid2812.220424
- Wong MM and Brummer ME** (1978) Cuticular morphology of five species of *Dirofilaria*: A scanning electron microscope study. *Journal of Parasitology* **64**, 108–114. doi: 10.2307/3279618