


COVID-19 Vulnerability Mapping of Asian Countries

Showmitra Kumar Sarkar , Md. Manjur Morshed and Tanmoy Chakraborty

Department of Urban and Regional Planning, Khulna University of Engineering & Technology (KUET), Bangladesh

Brief Report

Cite this article: Sarkar SK, Morshed MM, Chakraborty T. COVID-19 vulnerability mapping of Asian countries. *Disaster Med Public Health Prep.* 17(e241), 1–8. doi: <https://doi.org/10.1017/dmp.2022.139>.

Keywords:

Coronavirus; pandemic; factors; principal component analysis; Asian countries

Corresponding author:

Showmitra Kumar Sarkar,
E-mail: mail4dhrubo@gmail.com

Abstract

Objective: The objective of this study is to map vulnerability of Asian countries to the COVID-19 pandemic.

Method: According to the Intergovernmental Panel on Climate Change (IPCC) 2007 framework for natural hazards, vulnerability is a function of exposure, sensitivity, and adaptive capacity. From an extensive literature review, we identified 16 socioeconomic, meteorological, environmental, and health factors that influence coronavirus disease 2019 (COVID-19) cases and deaths. The underlying factors of vulnerability were identified using principal component analysis.

Results: Our findings indicate that the percentage of the urban population, obesity rate, air connectivity, and the population aged 65 and over, diabetes prevalence, and PM_{2.5} levels all contributed significantly to COVID-19 sensitivity. Subsequently, governance effectiveness, human development index (HDI), vaccination rate, and life expectancy at birth, and gross domestic product (GDP) all had a positive effect on adaptive capacity. The estimated vulnerability was corroborated by a Pearson correlation of 0.615 between death per million population and vulnerability.

Conclusion: This study demonstrates the application of universal indicators for assessing pandemic vulnerability for informed policy interventions such as the COVAX vaccine roll-out priority. Despite data limitations and a lack of spatiotemporal analysis, this study's methodological framework allows for ample data incorporation and replication.

Coronavirus disease 2019 (COVID-19) is an unprecedented event in modern history. The pandemic has impacted the majority of the world to some extent, resulting in a slew of social and economic costs. Pandemics are predicted to become more common in the coming years.^{1,2} As a result, minimizing pandemic vulnerability while increasing pandemic adaptive capacity is critical.^{3,4}

Numerous sensitivity factors have been linked to the spread of the pandemic. Demographic factors such as population density, age distribution, sex, and urbanization are correlated with COVID-19 cases and deaths.^{5–8} Second, meteorological and environmental factors such as average temperature, outdoor environment, particulate matter, environmental zones and wind speed are associated with COVID-19 cases and deaths.^{6,9–11} Third, underlying health factors such as diabetes, obesity and nutritional status have been identified as key contributors of COVID-19 related morbidity and mortality.^{12–14} Fourth, studies recognize ethnicity, socio-economic inequality, poverty and disposable income as significant factors of COVID-19 cases and deaths.^{10,13,15} Fifth, travel patterns and international connectivity are related to virus spread and infections.^{16–18} Finally, studies point to the (in)capacity of health systems to cope with pandemic crisis—essentially an indicator of governance effectiveness and adaptive capacity.^{15,19,20}

Vulnerability quantification is a necessary condition for proactive response. Measuring Asian vulnerability to COVID-19 is important given it continues to report the highest numbers of new infections in the world. India, the second largest Asian country in terms of population concentration, is leading in the number of new infections and deaths.²¹ This has far-reaching implications for neighboring South Asian countries and beyond. Asia is primarily composed of developing economies and the pandemic has slowed the region's continuous economic growth, affecting millions of people. The majority of available literature on COVID-19 vulnerability is either country- or global-specific. For example, the Social Vulnerability Index (SVI) in the United States considers socioeconomic status, household composition and disability, minority status and language, and housing type and transportation for measuring vulnerable groups during public health emergencies.²² Similarly, vulnerability to COVID-19 pandemic in India reflects on 5 socio-economic domains: Socioeconomic, demographic, housing and hygiene, epidemiological, and health system.²³ To date, there has been no study that measures Asian countries' vulnerability to COVID-19 pandemic using a natural disaster, which is an external stressor, framework.




Figure 1. Countries included in the study.

The objective of this research is to measure vulnerability of Asian countries to the COVID-19 pandemic. The Intergovernmental Panel on Climate Change (IPCC) 2007 framework for quantifying vulnerability to natural hazards can be applied to the COVID-19 pandemic. Vulnerability is comprised of 3 components: exposure, sensitivity, and adaptive capacity. We used principal component analysis (PCA) to identify the major factors affecting sensitivity and adaptive capacity to estimate vulnerability. Exposure refers to an external stress, such as COVID-19, whereas sensitivity and adaptive capacity are measures of susceptibility and coping capacity to the external stress, respectively.²⁴ Our research does not consider risk of COVID-19 transmission as a component of vulnerability (see Bjørnstad et al.²⁵). Subsequently, this research is based on the reported COVID-19 cases and deaths, which is different from the Susceptible-Exposed-Infectious-Removed (SEIRS) model that considers factors such as loss of immunity, birth, and death in COVID-19 transmission dynamics.²⁶

Methods

Study Area

Forty-three of 46 Asian countries were selected for the study (Figure 1). Due to the scarcity of data, this study excluded China and its territorial geographic areas (Hong Kong and Taiwan), North Korea, and Palestine.

Analytical Method

We have used vulnerability to natural disasters as an analytical framework for this study, where vulnerability is defined as the function of exposure, sensitivity, and adaptive capacity (Equation 1).^{27–30} According to the IPCC's 2007 report, vulnerability is the result of the interaction of exposure, sensitivity, and adaptive capacity. According to the IPCC's fourth assessment report (AR4), vulnerability to climate change is defined as "the degree to which a system is susceptible to and incapable of coping with adverse effects of climate change, including climate variability and extremes. Vulnerability is a function of the type, magnitude, and rate of climate change and variation to which a system is exposed, as well as its sensitivity and adaptive capacity".³¹ While IPCC does not define exposure, it is understood as an external stressor that contributes to vulnerability.³² Vulnerability is

thus viewed as the first-order impact caused by the exposure due to the system's sensitivity, which is moderated by its adaptive capacity.²⁴

$$\text{Vulnerability} = \text{Exposure} + \text{Sensitivity} - \text{Adaptive Capacity} \quad (1)$$

COVID-19 exposure, sensitivity, and adaptive capacity are determined by several independent variables (Table 1). Exposure is quantified by the number of cases per million population. Sensitivity and adaptive capacity variables are classified in such a way that their values increase with increasing sensitivity or adaptive capacity. For instance, a high prevalence of diabetes increases morbidity and mortality when associated with COVID-19.^{12,33} In this study, COVID-19-related deaths per million population are used as a proxy for a country's pandemic vulnerability.

The definition of vulnerable people can change dynamically in a COVID-19 scenario. Globally, the economic cost of the pandemic is estimated to be staggering at US \$10 trillion between 2020 and 2021.³⁴ Vulnerable people include not only the elderly and those with chronic illnesses and comorbidities, but also those who may struggle financially, mentally, or physically as a result of the onset of COVID.³⁵ However, this study's vulnerability estimation is limited to exposure, 8 sensitivity, and 7 adaptive capacity variables.

Multicollinearity refers to a linear correlation between a large number or all of the independent variables in a dataset, which makes estimating the relationship between each independent and dependent variable difficult.³⁶ Multicollinearity occurs when 1 independent variable shifts in response to a change in another independent variable. As a result, a high degree of multicollinearity among the independent variables may influence the results and their interpretation. Principal component analysis (PCA) is a widely used multivariate analysis technique that breaks down multiple correlated variables into several uncorrelated components.³⁷ We used PCA to eliminate multicollinearity in data.

Varimax rotation was used to calculate the factor loadings of sensitivity and adaptive capacity, which have an inverse relationship with the factor score. The Kaiser-Meyer-Olkin (KMO) and Bartlett's Test of Sphericity were used to determine the sampling adequacy for principal component analysis.³⁸ A statistically significant principal component analysis requires a minimum KMO value of 0.50. Significant factors are those that satisfy the Kaiser Eigenvalue criterion (>1) and account for a greater than one-third

Table 1. Exposure, sensitivity, adaptive capacity, and vulnerability factors

	Factors	Min	Max	Mean	SD	Source
Exposure	Cases per 1 million population	232.00	152606.00	37674.49	40416.89	(as of 04 August 2021) ²¹
Sensitivity	Population density (people per sq. km of land area)	2.04	7953.00	448.95	1245.14	³⁹
	Population ages 65 and above (% of total population)	1.16	28.00	6.82	4.94	
	Urban population (% of total population)	18.59	100.00	60.00	24.85	
	PM _{2.5} air pollution, mean annual exposure (micrograms per cubic meter)	5.90	99.73	38.09	23.78	
	Diabetes prevalence (% of population ages 20 to 79)	3.90	19.90	9.33	3.86	
	Obesity Rates	2.10	37.90	16.85	11.31	⁴⁰
	Death per 100,000 Cardio-vascular	30.29	361.17	178.06	89.05	⁴¹
	Air Connectivity Index	2.02	6.05	3.71	.86	⁴²
Adaptive capacity	Hospital beds (per 1,000 people)	0.30	63.00	4.78	9.44	³⁹
	Doctors (per 1,000 people)	0.01	6.13	1.79	1.34	
	Life expectancy at birth, total (years)	58.00	83.04	73.48	6.81	
	GDP per capita (current US\$)	503.32	181402.83	19328.19	32966.61	
	Governance Effective Index	-2.28	2.22	-.02	.95	⁴³
	Human Development Index	0.47	0.94	0.74	0.12	⁴⁴
	People vaccinated per 100	0.04	71.20	22.61	23.03	(as of 04 August 2021) ⁴⁵
Vulnerability	Deaths per 1 million population	0.90	1556.00	367.42	391.27	(as of 04 August 2021) ²¹

of the variance in the data. The normalized exposure, sensitivity, and adaptive capacity variables were used to calculate vulnerability to COVID-19. Equation 2 is used to normalize the data.

$$\acute{x} = \frac{x - \text{Min}(x)}{\text{Max}(x) - \text{Min}(x)} \quad (2)$$

Where,

\acute{x} = normalized data of a country;

x = data of a country;

Min(x) = the minimum value among the countries;

Max(x) = the maximum value among the countries.

To normalize India's exposure, for example, the number of cases per million population was calculated for each of the study countries. Similarly, the maximum number of cases per million (Bahrain) and minimum number of cases per million (Yemen) were noted. Following that, the normalized exposure of India was calculated using the min-max normalization technique (Equation 2). The primary advantage of min-max normalization is that it produces a distribution of data between 0 and 1, with 0 and 1 representing the lowest and highest values, respectively.

The vulnerability to COVID-19 of the respective countries is determined using Equation 1. On the basis of Jenks' natural breaks classification method, the normalized COVID-19 vulnerability is reclassified into 3 classes. Appendix Table 1 shows the normalized values of exposure, sensitivity, adaptive capacity, and vulnerability to COVID-19 (see also Figure 2). Finally, we validated the calculated vulnerability by comparing it to the actual vulnerability (death per million population from COVID-19) by Pearson correlation.

ArcGIS 10.5 was used to create maps of selected countries' exposure, sensitivity, adaptive capacity, and vulnerability. IBM-SPSS 25 was used to conduct the statistical analysis.

Data

COVID-19 exposure (cases/million as of 04 August 2021), sensitivity, adaptive capacity, and vulnerability (deaths/million) data/

variables were gathered from a variety of online sources. A total of 16 factors were identified following a thorough literature review. Three subgroups of variables had been defined: 1 for exposure, 8 for sensitivity, and 7 for adaptive capacity (Table 1).

Results

Principal Component Analysis

The Bartlett's Test of Sphericity yielded a statistically significant result (P -value 0.05), indicating that the data set was suitable for factor analysis. KMO tests for sensitivity (0.519) and adaptive capacity (0.576) indicated that the variables had a high degree of informational overlap. As a result, factor analysis was a viable method for eliminating multicollinearity between independent variables. Based on the Kaiser Eigenvalue criterion and scree plots, 3 components for sensitivity and 2 components for adaptive capacity were identified.

For sensitivity, 3 components independently explained 40.91 percent, 25.67 percent, and 13.49 percent (cumulatively 80.07 percent) of the data variance of the input variables of COVID-19 sensitivity (Table 2). Three factors in component 1—percentage of urban population (0.922), obesity rates (0.883), and air connectivity (0.675)—all contributed positively and significantly to the COVID-19 sensitivity, as determined by varimax rotation (Table 3). Component 2 isolates the population aged 65 and over as the only significant positive factor (0.589). Component 3 contains 2 significant positive variables: Diabetes prevalence (0.695) and PM_{2.5} (0.880).

Two adaptive capacity components accounted for 42.97% and 24.48% (cumulatively 67.46%) of the variance in the input variables, respectively. Three factors in component 1—governance effectiveness index (0.851), HDI (0.872) and vaccination rate (0.850)—all contributed positively and significantly to adaptive capacity against COVID-19. Component 2 of the adaptive capacity has 2 significant factors: Life expectancy at birth (0.934) and GDP (0.664). Sensitivity and adaptive capacity were calculated for each country using the principal component analysis components.




Figure 2. (a) Exposure, (b) sensitivity, (c) adaptive capacity, and (d) vulnerability to COVID-19.

Table 2. Result of component extraction of factor analysis

	Component	% of Variance	Cumulative %	Factor Score	Description (factor score)
Sensitivity	1	40.91	40.91	-1.877 to 1.663	(-) low sensitive
	2	25.67	66.58	-1.941 to 2.397	(+) highly sensitive
	3	13.49	80.07	-1.720 to 2.188	
Adaptive capacity	1	42.97	42.97	-2.013 to 2.125	(-) low adaptability
	2	24.48	67.45	-2.806 to 2.080	(+) highly adaptive

Table 3. Factors loading of the factors

Factors	Components*			
	1	2	3	
Sensitivity	Population density (people per sq. km of land area)	-	0.417	-
	Population ages 65 and above (% of total population)	-	0.589	-0.541
	Urban population (% of total population)	0.922	0.257	-
	Obesity rates	0.883	-0.348	0.237
	Diabetes prevalence (% of population ages 20 to 79)	0.409	-	0.695
	Death per 100,000 Cardio-vascular	-	-0.960	-
	PM _{2.5} air pollution, mean annual exposure (micrograms per cubic meter)	-	-0.145	0.880
	Air Connectivity Index	0.675	0.456	-
Adaptive capacity	Hospital beds (per 1000 people)	-0.135	-	-
	Doctors (per 1000 people)	0.431	-0.568	-
	Life expectancy at birth, total (years)	0.149	0.934	-
	Gross domestic products (GDP) per capita (current US\$)	0.138	0.664	-
	Governance Effective Index	0.851	-	-
	Human Development Index (HDI)	0.872	-	-
	People vaccinated per 100	0.850	-	-

*Bold values indicate high factor loadings.

In summary, high urbanization and obesity, along with air pollution, have the highest potential for increasing sensitivity to COVID-19. On the contrary, 4 variables—life expectancy at birth, governance effectiveness, HDI and vaccination rate—positively contribute to adaptive capacity against COVID-19 pandemic.

Exposure, Sensitivity, and Adaptive Capacity

The maps in Figure 2 depict the selected Asian countries' exposure (a), sensitivity (b), adaptive capacity (c), and vulnerability (d). Bahrain ranks highest in terms of exposure (normalized range 0.38-1.0), followed by the Maldives, Georgia, Jordan, Israel, and Kuwait (Figure 2a). Clearly, countries with small landmasses are more exposed than countries with larger landmasses. Oman, Mongolia, Iran, Iraq, Malaysia, and Azerbaijan, to name a few, have a moderate level of exposure (normalized range, 0.10-0.37). Saudi Arabia, Sri Lanka, the Philippines, Indonesia, and Singapore, among other countries, have a low level of exposure (0-0.09).

The most sensitive countries to COVID-19 are Qatar, Bahrain, Saudi Arabia, Turkey, and the United Arab Emirates (Figure 2b). On the other hand, Singapore, Qatar, the United Arab Emirates, and Bahrain, among others, had the highest adaptive capacity (Figure 2c). According to Equation 1, 14 of the 43 countries, including Bahrain, the Maldives, Georgia, Jordan, Kuwait, and Israel, had the highest vulnerability to COVID-19 (normalized range, 0.55-1.00) (Figure 2d). Similarly, 14 countries had a medium

(0.54-0.24) vulnerability to COVID-19, while the remaining countries had a low (0.23-0.00) vulnerability.

Finally, a Pearson correlation coefficient of 0.615 (at the 0.01 level of significance) between the calculated vulnerability and death per million population from COVID-19 for the selected Asian countries indicated an acceptable level of vulnerability estimation.

Discussion

Numerous studies have been conducted on the contextual and country-specific factors that contribute to COVID-19 vulnerability. International comparisons of vulnerability to the COVID-19 pandemic, on the other hand, are difficult due to a lack of data and contextual differences between countries. PCA is frequently used to study natural disaster behavior,⁴⁶ and it has the potential to identify common COVID-19 factors globally.⁴⁷⁻⁴⁹

We considered the COVID-19 pandemic as a hazard in this study. Therefore, we have used the IPCC's 2007 hazard framework to assess Asian countries' vulnerability to COVID-19, incorporating 16 variables that quantify each country's vulnerability. Using principal component analysis, a total of 5 components (3 for sensitivity and 2 for adaptive capacity) were found to be suitable for measuring vulnerability. Six variables—the percentage of the urban population, the population aged 65 and over, obesity rate, air connectivity, diabetes prevalence, and PM_{2.5}—all contributed

positively and significantly to the COVID-19 sensitivity. These findings corroborated with those of several other studies that have linked COVID-19 sensitivity to demographic factors such as urbanization and the population aged 65 and over,⁵⁻⁷ underlying health condition,^{12,14} air connectivity,¹⁶ and meteorological factors such as PM_{2.5}.⁹

Five factors explained COVID-19 adaptive capacity: The governance effectiveness index, the human development index (HDI), the vaccination rate, the life expectancy at birth, and the gross domestic product (GDP). Similar to the findings of Babu *et al.*,¹⁹ this research indicates that governance effectiveness is a predictor of an effective response to a pandemic. Bangladesh, for example, ranks low in adaptive capacity due to a poor track record managing the pandemic and residents' disregard for health guidelines.^{50,51} We used GDP and HDI as indicators of countries' socioeconomic characteristics, and a high value for these 2 variables increase adaptive capacity to COVID-19. Additionally, we demonstrated that vaccination is critical for building adaptive capacity against the COVID-19 pandemic.

The analytical framework used in conjunction with the findings of this study has implications for national and international policy responses to COVID-19. Separating exposure, sensitivity, and adaptive capacity enables a more precise assessment of a country's preparedness for a pandemic. India, for instance, has a high sensitivity but a low vulnerability due to its high adaptive capacity. Additionally, identifying universal risk factors for vulnerability enables socioeconomic and political policy interventions such as increased vaccination in high-density and urban areas, as well as for older adults. More importantly, the vulnerability of each country can serve as a benchmark for prioritizing vaccine roll-out through the COVID-19 Vaccines Global Access (COVAX) program in the event of future pandemics.

Finally, we used Pearson correlation to validate the estimated vulnerability to COVID-19 death. Our vulnerability, as estimated, accounts for 61.5% of COVID-19 deaths, which is quite acceptable. Apart from the study's limitations, discussed in the following section, vulnerability appears to be a dynamic concept. As a result, using death from COVID as a proxy for vulnerability is debatable. Incorporating variables relating to financial, mental, and physical consequences can undoubtedly help improve the vulnerability estimation model.

This study has several limitations. To begin, China (including Hong Kong and Taiwan), North Korea, and Palestine were not considered for measuring Asian Vulnerability. Second, the total number of confirmed cases and deaths was based on the official report that lacks data confidence. For instance, India demonstrated a high level of exposure but a low level of vulnerability. However, India's death toll is estimated to be 4 times that of official reports.⁵² Additionally, governance, vaccinations, and even population enumeration were reported with varying degrees of accuracy across these countries. Third, the study established vulnerability to COVID-related deaths. This is problematic, however, because the vulnerable population included those who suffered financially, psychologically, or physically as a result of COVID-19's onset. Fourth, despite the fact that COVID-19 is a respiratory illness, we do not have an explanation for death from cardio-vascular diseases with a high yet negative factor loading for sensitivity (see Table 3). Additionally, this research ignores several well-documented COVID-19 sensitivity factors, such as population density.⁵ Fifth, we disregarded the severity of COVID-19 infection and made no distinction between asymptomatic, mild, and severe cases. Finally, this study excluded

spatiotemporal analysis and relied exclusively on cross-sectional data rather than panel data.

Conclusions

COVID-19 is a global concern at the moment, and pandemics are expected to occur more frequently in the future. Assessing vulnerability to pandemics is a prerequisite for adaptive future planning in the event of a pandemic. We assessed Asian countries' vulnerability to COVID-19 using the IPCC's 2007 natural disaster framework and a set of universal indicators. COVID-19 vulnerability was determined by 1 component for exposure, 3 components for sensitivity, and 2 components for adaptive capacity. When death per million population was used as a proxy for vulnerability, our estimate was 61.5% accurate.

The findings of this study have policy implications for pandemic preparedness, including vaccine roll-out and response measures. Additionally, the analytical framework used in this study is applicable to other infectious and transmissible diseases, such as tuberculosis and dengue fever. Apart from data limitations, the study makes no attempt to account for spatiotemporal variation in COVID-19 vulnerability. However, the methodology used in this study leaves ample room for data manipulation, addition, and replication to improve the estimation of COVID-19 vulnerability.

Supplementary material. To view supplementary material for this article, please visit <https://doi.org/10.1017/dmp.2022.139>

Acknowledgments. We are indebted to the Editors of this journal and anonymous reviewers for their insightful comments and suggestions on earlier drafts of this manuscript.

Conflicts of interest. The authors declare that they are unaware of any competing financial interests or personal relationships that might appear to have influenced the work reported in this study.

Ethical standards. The authors declare that no ethical approval is required. Our institute does not require ethical approval for this type of study.

References

1. Marani M, Katul GG, Pan WK, *et al.* Intensity and frequency of extreme novel epidemics. *Proc Natl Acad Sci U S A.* 2021;118(35):e2105482118. doi: [10.1073/pnas.2105482118](https://doi.org/10.1073/pnas.2105482118)
2. Gavi. 5 reasons why pandemics like COVID-19 are becoming more likely. Published 2020. Accessed May 21, 2022. <https://www.gavi.org/vaccineswork/5-reasons-why-pandemics-like-covid-19-are-becoming-more-likely>
3. Connolly C, Ali SH, Keil R. On the relationships between COVID-19 and extended urbanization. *Dialogues Hum Geogr.* 2020;10(2):213-216. doi: [10.1177/2043820620934209](https://doi.org/10.1177/2043820620934209)
4. Sharifi A, Khavarian-Garmsir AR. The COVID-19 pandemic: impacts on cities and major lessons for urban planning, design, and management. *Sci Total Environ.* 2020;749:1-3. doi: [10.1016/j.scitotenv.2020.142391](https://doi.org/10.1016/j.scitotenv.2020.142391)
5. Velasco JM, Tseng WC, Chang CL. Factors affecting the cases and deaths of covid-19 victims. *Int J Environ Res Public Health.* 2021;18(2):1-10. doi: [10.3390/ijerph18020674](https://doi.org/10.3390/ijerph18020674)
6. Sharif N, Dey SK. Impact of population density and weather on COVID-19 pandemic and SARS-CoV-2 mutation frequency in Bangladesh. *Epidemiol Infect.* 2021;149:e16. doi: [10.1017/S0950268821000029](https://doi.org/10.1017/S0950268821000029)
7. Sarkar SK, Ekram KMM, Das PC. Spatial modeling of COVID-19 transmission in Bangladesh. *Spat Inf Res.* 2021;29(5):715-726. doi: [10.1007/s41324-021-00387-5](https://doi.org/10.1007/s41324-021-00387-5)
8. Sarkar SK. COVID-19 susceptibility mapping using multicriteria evaluation. *Disaster Med Public Health Prep.* 2020;14(4):521-537. doi: [10.1017/dmp.2020.175](https://doi.org/10.1017/dmp.2020.175)

9. **Aabed K, Lashin MMA.** An analytical study of the factors that influence COVID-19 spread. *Saudi J Biol Sci.* 2021;28(2):1177-1195. doi: [10.1016/j.sjbs.2020.11.067](https://doi.org/10.1016/j.sjbs.2020.11.067)
10. **Tian T, Zhang J, Hu L, et al.** Risk factors associated with mortality of COVID-19 in 3125 counties of the United States. *Infect Dis Poverty.* 2021;10(1):3. doi: [10.1186/s40249-020-00786-0](https://doi.org/10.1186/s40249-020-00786-0)
11. **Sarkar SK, Khan MMH.** Impact of COVID-19 on PM_{2.5} pollution in fast-growing megacity Dhaka, Bangladesh. *Disaster Med Public Health Prep.* 2021;30:1-4. doi: [10.1017/dmp.2021.131](https://doi.org/10.1017/dmp.2021.131)
12. **Yu W, Rohli KE, Yang S, et al.** Impact of obesity on COVID-19 patients. *J Diabetes Complications.* 2021;35(3):107817. doi: [10.1016/j.jdiacomp.2020.107817](https://doi.org/10.1016/j.jdiacomp.2020.107817)
13. **Yaylali A.** Factors affecting the number of Covid-19 cases and the death rate: empirical evidence from German states. Published online 2020. Accessed June 15, 2022. https://papers.ssrn.com/sol3/papers.cfm?abstract_id=3617986
14. **Baik I.** Region-specific COVID-19 risk scores and nutritional status of a high-risk population based on individual vulnerability assessment in the national survey data. *Clin Nutr.* 2021;23:S0261-5614(21)00093-5. doi: [10.1016/j.clnu.2021.02.019](https://doi.org/10.1016/j.clnu.2021.02.019)
15. **Rocha R, Atun R, Massuda A, et al.** Effect of socioeconomic inequalities and vulnerabilities on health-system preparedness and response to COVID-19 in Brazil: a comprehensive analysis. *Lancet Glob Health.* 2021;(21):1-11. doi: [10.1016/s2214-109x\(21\)00081-4](https://doi.org/10.1016/s2214-109x(21)00081-4)
16. **Christidis P, Christodoulou A.** The predictive capacity of air travel patterns during the global spread of the covid-19 pandemic: risk, uncertainty and randomness. *Int J Environ Res Public Health.* 2020;17(10):3356. doi: [10.3390/ijerph17103356](https://doi.org/10.3390/ijerph17103356)
17. **Wesolowski A, Zu Erbach-Schoenberg E, Tatem AJ, et al.** Multinational patterns of seasonal asymmetry in human movement influence infectious disease dynamics. *Nat Commun.* 2017;8(1):2069. doi: [10.1038/s41467-017-02064-4](https://doi.org/10.1038/s41467-017-02064-4)
18. **Brannen DE, Alhammad A, Branum M, et al.** International air travel to Ohio, USA, and the impact on Malaria, Influenza, and Hepatitis A. *Scientifica (Cairo).* 2016;2016:8258946. doi: [10.1155/2016/8258946](https://doi.org/10.1155/2016/8258946)
19. **Babu GR, Khetrpal S, John DA, et al.** Pandemic preparedness and response to COVID-19 in South Asian countries. *Int J Infect Dis.* 2021;104:169-174. doi: [10.1016/j.ijid.2020.12.048](https://doi.org/10.1016/j.ijid.2020.12.048)
20. **Brannen DE, McDonnell MA, Schmitt A.** Organizational culture on community health outcomes after the 2009 H1N1 pandemic. *J Organ Cult Commun Confl.* 2013;17(1):1-18. Accessed April 29, 2022. <https://search.proquest.com/openview/dd928127b8dcf137d752e8b493952389/1?pq-origsite=gscholar&cbl=38870>
21. **Worldometer.** COVID Live - Coronavirus Statistics. Published 2022. Accessed January 30, 2022. <https://www.worldometers.info/coronavirus/>
22. **Agency for Toxic Substances and Disease.** At a glance: CDC/ATSDR social vulnerability index | place and health | ATSDR. Published 2021. Accessed May 21, 2022. https://www.atsdr.cdc.gov/placeandhealth/svi/at-a-glance_svi.html
23. **Acharya R, Porwal A.** A vulnerability index for the management of and response to the COVID-19 epidemic in India: an ecological study. *Lancet Glob Health.* 2020;8(9):e1142-e1151. doi: [10.1016/S2214-109X\(20\)30300-4](https://doi.org/10.1016/S2214-109X(20)30300-4)
24. **Sharma J, Ravindranath NH.** Applying IPCC 2014 framework for hazard-specific vulnerability assessment under climate change. *Environ Res Commun.* 2019;1(5). doi: [10.1088/2515-7620/ab24ed](https://doi.org/10.1088/2515-7620/ab24ed)
25. **Bjornstad ON, Shea K, Krzywinski M, et al.** Modeling infectious epidemics. *Nat Methods.* 2020;17(5):455-456. doi: [10.1038/s41592-020-0822-z](https://doi.org/10.1038/s41592-020-0822-z)
26. **Bjornstad ON, Shea K, Krzywinski M, et al.** The SEIRS model for infectious disease dynamics. *Nat Methods.* 2020;17(6):557-558. doi: [10.1038/s41592-020-0856-2](https://doi.org/10.1038/s41592-020-0856-2)
27. **Brooks N.** Vulnerability, risk and adaptation: a conceptual framework. Tyndall Centre for Climate Change Research. *Tyndall Cent Clim Chang Res.* 2003;(February):1-20.
28. **Turner BL, Kasperson RE, Matsone PA, et al.** A framework for vulnerability analysis in sustainability science. *Proc Natl Acad Sci U S A.* 2003;100(14):8074-8079. doi: [10.1073/pnas.1231335100](https://doi.org/10.1073/pnas.1231335100)
29. **Cutter SL, Boruff BJ, Shirley WL.** Social vulnerability to environmental hazards. *Soc Sci Q.* 2003;84(2):242-261. doi: [10.1111/1540-6237.8402002](https://doi.org/10.1111/1540-6237.8402002)
30. **Wilhelmi OV, Hayden MH.** Connecting people and place: a new framework for reducing urban vulnerability to extreme heat. *Environ Res Lett.* 2010;5(1). doi: [10.1088/1748-9326/5/1/014021](https://doi.org/10.1088/1748-9326/5/1/014021)
31. **IPCC.** Climate Change 2007: Impacts, Adaptations and Vulnerability. IPCC 2007 Contribution of Working Group II to the Fourth Assessment Report of the Intergovernmental Panel on Climate. Parry M, Canziani O, Palutikof J, Linden P van der, Hanson C, eds. Cambridge: Cambridge University Press; 2007.
32. **Jurgilevich A, Räsänen A, Groundstroem F, et al.** A systematic review of dynamics in climate risk and vulnerability assessments. *Environ Res Lett.* 2017;12(1):13002. doi: [10.1088/1748-9326/aa5508](https://doi.org/10.1088/1748-9326/aa5508)
33. **Morshed MM, Sarkar SK.** Common factors of COVID-19 cases and deaths among the most affected 50 countries. *Diabetes Metab Syndr.* 2021;15(5):102247. doi: [10.1016/j.dsx.2021.102247](https://doi.org/10.1016/j.dsx.2021.102247)
34. **The Economist.** What is the economic cost of covid-19? Published January 9 2021. Accessed February 5, 2022. <https://www.economist.com/finance-and-economics/2021/01/09/what-is-the-economic-cost-of-covid-19>
35. **The Lancet.** Redefining vulnerability in the era of COVID-19. *Lancet.* 2020;395(10230):1089. doi: [10.1016/S0140-6736\(20\)30757-1](https://doi.org/10.1016/S0140-6736(20)30757-1)
36. **Islam ARMT, Talukdar S, Mahato S, et al.** Flood susceptibility modelling using advanced ensemble machine learning models. *Geosci Front.* 2021;12(3). doi: [10.1016/j.gsf.2020.09.006](https://doi.org/10.1016/j.gsf.2020.09.006)
37. **Mahmoudi MR, Heydari MH, Qasem SN, et al.** Principal component analysis to study the relations between the spread rates of COVID-19 in high risks countries. *Alexandria Eng J.* 2021;60(1):457-464. doi: [10.1016/j.aej.2020.09.013](https://doi.org/10.1016/j.aej.2020.09.013)
38. **Williams B, Onsmann A, Brown T.** Exploratory factor analysis: a five-step guide for novices. *J Emerg Prim Heal Care.* 2010;8(3):1-13. doi: [10.33151/ajp.8.3.93](https://doi.org/10.33151/ajp.8.3.93)
39. **World Bank.** World Bank open data indicators. Published 2020. Accessed February 12, 2022. <https://data.worldbank.org/>
40. **ProCon.org.** Global Obesity Levels - Obesity - ProCon.org. Britannica ProCon.org. Published 2020. Accessed February 12, 2022. <https://obesity.procon.org/global-obesity-levels/>
41. **WorldHealthRanking.** Coronary heart disease death rate by country. Published online 2022. Accessed February 12, 2022. <https://www.worldlifeexpectancy.com/cause-of-death/coronary-heart-disease/by-country/>
42. **Arvis J-F, Shepherd B.** The Air Connectivity Index: Measuring integration in the Global Air Transport Network. World Bank Policy Research Working Paper 5722. 2011. Accessed June 15, 2022. <https://elibrary.worldbank.org/doi/abs/10.1596/1813-9450-5722>
43. **Valev N, Bieri F, Varbanova N.** Government effectiveness by country, around the world|TheGlobalEconomy.com. The Global Economy. Published 2015. Accessed February 12, 2022. https://www.theglobaleconomy.com/rankings/wb_government_effectiveness/#USA
44. **United Nations.** Latest Human Development Index Ranking | Human Development Reports. Human Development Reports. Published 2020. Accessed February 12, 2022. <https://hdr.undp.org/en/content/latest-human-development-index-ranking>
45. **Mathieu E, Ritchie H, Ortiz-Ospina E, et al.** A global database of COVID-19 vaccinations. *Nat Hum Behav.* 2021;5(7):947-953. doi: [10.1038/s41562-021-01122-8](https://doi.org/10.1038/s41562-021-01122-8)
46. **Alam MS, Chakraborty T.** Understanding the nexus between public risk perception of COVID-19 and evacuation behavior during cyclone Amphan in Bangladesh. *Heliyon.* 2021;7(7):e07655. doi: [10.1016/j.heliyon.2021.e07655](https://doi.org/10.1016/j.heliyon.2021.e07655)
47. **Duarte P, Riveros-Perez E.** Understanding the cycles of COVID-19 incidence: principal component analysis and interaction of biological and socio-economic factors. *Ann Med Surg (Lond).* 2021;66:102437. doi: [10.1016/j.amsu.2021.102437](https://doi.org/10.1016/j.amsu.2021.102437)

48. **Joshua V, Sylvia Grace J, Godwin Emmanuel J, et al.** Spatial mapping of COVID-19 for Indian states using Principal Component Analysis. *Clin Epidemiol Glob Health*. 2021;10:100690. doi: [10.1016/j.cegh.2020.100690](https://doi.org/10.1016/j.cegh.2020.100690)
49. **Viezzler J, Biondi D.** The influence of urban, socio-economic, and environmental aspects on COVID-19 cases, deaths and mortality: a multi-city case in the Atlantic Forest, Brazil. *Sustain Cities Soc*. 2021;69:102859. doi: [10.1016/j.scs.2021.102859](https://doi.org/10.1016/j.scs.2021.102859)
50. **Dhaka Tribune.** Health minister blames negligence for rise in coronavirus cases. Dhaka Tribune. Published 2021. Accessed February 5, 2022. <https://archive.dhakatribune.com/bangladesh/2021/04/03/health-minister-blames-negligence-for-rise-in-coronavirus-cases>
51. **Shammi M, Bodrud-Doza M, Islam ARMT, et al.** COVID-19 pandemic, socioeconomic crisis and human stress in resource-limited settings: a case from Bangladesh. *Heliyon*. 2020;6(5). doi: [10.1016/j.heliyon.2020.e04063](https://doi.org/10.1016/j.heliyon.2020.e04063)
52. **Anand A, Sandefur J, Subramanian A.** Three new estimates of India's all-cause excess mortality during the COVID-19 pandemic. 2021. Accessed June 15, 2022. www.cgdev.org/publication/three-new-estimates-indias-all-cause-excess-mortality-during-covid-19-pandemic