

GAD symptoms (OR = 5.35; 95% C.I. 2.03-14.15). Respondents who suffered a loss of property because of the wildfire were two times more likely to develop GAD symptoms (OR = 2.36; 95% C.I. 1.01-22.62).

Conclusions: Formulators of policy may mitigate GAD symptoms, particularly after natural disasters, by making long-term mental health counseling available and a key component of post-disaster management, and by investing in the social capital of the people to build resilience and support to deal with the post-disaster mental health effects.

Disclosure of Interest: None Declared

O0064

A Pilot Study Comparing a Community of Practice Group Therapy Program with and without Concurrent Ketamine-assisted Therapy

V. W. L. Tsang^{1*}, R. Moyer², P. Kryskow³, Z. Walsh⁴ and S. Dames³
¹Psychiatry; ²Interdisciplinary Studies Graduate Program, UBC, Vancouver; ³Health and Human Services, Vancouver Island University, Nanaimo and ⁴Department of Psychology, UBC, Kelowna, Canada

*Corresponding author.

doi: 10.1192/j.eurpsy.2023.269

Introduction: Healthcare practitioners (HCPs) are facing a mental health crisis. Group therapies have long been used to treat symptoms associated with PTSD, anxiety and/or depression, however no studies have investigated the role of implementing group therapy with and without ketamine-assisted therapies (KaT).

Objectives: The current study investigated the effects of the Roots to Thrive (RTT) group therapy intervention both with and without adjunctive KaT.

Methods: In the present study we conduct a secondary analysis of data derived from the 12-week group psychotherapy program to that of the same program with adjunct KaT. Participants were administered a series of validated psychiatric assessment tools before and after the 12-weeks. Inclusion criteria included a diagnosis of treatment resistant mental health condition (depression, PTSD and/or generalized anxiety disorder) and a score of 15 or greater on the PTSD Checklist for DSM-5 (PCL-5). To assess the effects of time x group interaction and calculate differences between the RTT only and RTT-KaT subgroups, a repeated measures ANOVA was conducted. Effect sizes were calculated through partial eta-squared.

Results: Forty-nine HCPs with treatment-resistant PTSD, anxiety and/or depression were treated with the RTT group therapy model to target their symptoms. A total of 49 individuals (34 female, 10 male, 3 other) with a median age of 47 years old (SD 14.19) participated in the study. There were no statistically significant differences between RTT only (n=14) and RTT KaT (n=35) subgroups across gender [$X^2(1, N=44) = 2.84, ns$] or age [$F(1, 36) = .257, p = .615$]. From pre- to post-treatment, all patients showed significant reductions in scores of PTSD (from 39.3 to 20.99), depression (from 15.5 to 7.7) and anxiety (from 15.5 to 6.2). Two-way repeated measures ANOVA did not reveal any significant between-group differences between the RTT and RTT-KaT subgroups.

Conclusions: This observational study provides preliminary support for the potential of the RTT community of care model of group therapy and adds to a small but growing body of knowledge on the integration of group therapy and the broad category of psychedelic psychotherapies. Given the rapid proliferation and expansion of KaT clinics throughout North America, the finding that KaT did not appear to impact changes related to the RTT intervention suggests the need for further research to better explain the potential impacts of relational transference between the two groups, and the distinct contributions of ketamine administration in a group therapy context.

Disclosure of Interest: None Declared

O0065

Childhood trauma and anger in adults with and without depressive and anxiety disorders

N. De Bles^{1*}, L. E. Putz¹, N. Rius Ottenheim¹, A. M. van Hemert¹, B. M. Elzinga², B. W. Penninx³ and E. J. Giltay¹

¹Department of Psychiatry, LUMC; ²Department of Clinical Psychology, Leiden University, Leiden and ³Department of Psychiatry, Amsterdam UMC/Vrije Universiteit, Amsterdam, Netherlands

*Corresponding author.

doi: 10.1192/j.eurpsy.2023.270

Introduction: Childhood trauma (CT) is associated with severe sequelae, including personality disorders and stress-related mental health disorders that can perpetuate long into adulthood.

Objectives: We aimed to investigate (1) whether childhood trauma is associated with anger in adulthood, and, if so, (2) to explore which types of childhood trauma predominate in the prediction of anger, and (3) to explore whether the association is independent of psychopathology in a cohort that included participants without lifetime psychiatric disorders, with current or remitted depressive and anxiety disorders, or comorbid depressive and anxiety disorders.

Methods: In the Netherlands Study of Depression and Anxiety (NESDA), childhood trauma was assessed with a semi-structured Childhood Trauma Interview (CTI) at baseline, and analyzed in relation to anger as measured at 4-year follow-up with the Spielberger Trait Anger Subscale (STAS), the Anger Attacks Questionnaire, and cluster B personality traits (i.e., borderline, antisocial) of the Personality Disorder Questionnaire 4 (PDQ-4), using analysis of covariance (ANCOVA) and multivariable logistic regression analyses. Post-hoc analyses comprised cross-sectional regression analyses, using the Childhood Trauma Questionnaire – Short Form (CTQ-SF) obtained at 4-year follow-up.

Results: Participants ($n = 2,276$) were on average 42.1 years ($SD = 13.1$), and 66.3% were female. Childhood trauma showed a dose-response association with all anger constructs. Zooming in, all types of childhood trauma except for sexual abuse were associated with higher levels of trait anger, and a higher prevalence of anger attacks and antisocial personality traits in adulthood, independently of depression and anxiety. Additionally, all types of childhood trauma were significantly associated with borderline personality traits. Cross-sectionally, the effect sizes were larger compared to the analyses with the childhood trauma measured four years prior to the anger measures.