

Dimensions: head and neck 1.25 mm., trunk 3.2 mm., abdominal appendages 0.7 mm., eggstrings 2 mm., second maxillae 4.8 mm. (Fig. 6).

Diagnosis. *Cephalothorax* flattened dorsally; a slight dorsal carapace. *Neck* not segmented. *Trunk* pyriform. *Abdominal appendages* small and stout. *Eggstrings* stout, shorter than trunk. *Second maxillae* extremely long, longer than trunk, straight with expanded apices. *Bulla* resembling that of *Lernaeopoda*. A very small species.

My thanks are due to Miss E. C. Humphreys for kindly executing the figures.

REFERENCES.

- KRØYER (1837). Om Snyltekrebsene, isaer med Hensyn til den dansk Fauna. *Naturhist. Tidsskr.* Copenhagen. Vol. I.
 — (1863). Bidrag til Kundskab om Snyltekrebsene. *Ibid.* 3die Raekke, II Bind. Kjøbenhavn.
 LEIGH-SHARPE (1925). A Revision of the British Species of *Clavella*. *Parasitol.* 17, 194.
 RETZIUS (1829–1831). Beskrifning öfver en ny Scandinavisk *Lernaea* fran Nordsjön, kallad *Lernaea dalmanni*. *Kongl. Vetensk. Akad. Handling.* Stockholm, pp. 109–119, c. tab.; also (in German) *Froriep's Notizen*, 29, 6, Figs. 5–9; also in *Isis*, 1831, p. 1345, pl. IX.
 SCOTT, T. (1900). Notes on some Crustacean parasites of Fishes. *Eighteenth Ann. Report Fishery Board for Scotland*, pp. 144–148.
 SCOTT, T. and A. (1913–1914). The British Parasitic Copepoda. *The Ray Soc.* London.
 WILSON (1915). North American Parasitic Copepods belonging to the *Lernaeopodidae*, with a revision of the entire family. *Proc. U.S. Nat. Mus.* 47, 565–729. Washington.

ERRATUM.

Parasitology, 20, 87. The legend to Fig. 7 should read:

A 1. = antennules; A 2. = antennae.

(MS. received for publication 24. v. 1928.—Ed.)