

modifiable environmental factor. If pesticides are found to alter the immune response to COVID-19 infection and vaccination, these data will provide an evidence base for efforts to reduce pesticide exposure in children.

50

Why are Black and Mexican American children more vulnerable than White children to upper respiratory viral infection?

Darlene Bhavnani, D. Bhavnani, P. Dunphy, Paul Rathouz and E.C. Matsui

University of Texas at Austin

OBJECTIVES/GOALS: There is an excess risk of upper respiratory infection (URI) among Black and Mexican-American children in the US. Factors that underpin these disparities are largely unknown. We evaluated the extent to which socioeconomic status (SES), serum cotinine, obesity, and household size explained the association between race/ethnicity and URI. **METHODS/STUDY POPULATION:** We studied children, 6-17 years of age, who identified as Black, Mexican-American, or White in the National Health and Nutritional Examination Survey (2007-2012). URI was defined as a self-reported cough, cold, phlegm, runny nose, or other respiratory illness (excluding hay fever and allergies) in the past 7 days. The proportion of the association between race/ethnicity and URI explained by SES, serum cotinine, obesity, and household size was estimated as the average causal mediation effect (i.e., the indirect effect of race/ethnicity via the mediator) divided by the total effect of race/ethnicity. The average causal mediation effect was derived from survey weighted logistic regression models adjusted for age and sex. **RESULTS/ANTICIPATED RESULTS:** Black children were nearly 40% and Mexican American children were ~60% more likely to report a URI than those who identified as White (OR, 1.37; 95% CI, 1.06-1.77 and OR, 1.61; 95% CI, 1.21-2.13, respectively). Lower SES explained ~25% of the association between Black and Mexican American identity and URI (percent mediated 24.7; 95% CI, 23.0-26.6 and 26.1; 95% CI, 24.2-28.2, respectively). Obesity explained ~7% of the association between Black and Mexican-American identity and URI (percent mediated, 7.6; 95% CI, 7.3-8.0 and percent mediated, 6.7; 95% CI, 6.4-6.9, respectively). Nicotine exposure explained 8% of the association between Black identity and URI (percent mediated, 7.9; 95% CI, 5.6-10.1). **DISCUSSION/SIGNIFICANCE:** Lower SES explained a quarter of the association between race/ethnicity and URI. Low SES is a broad concept that may work through different mechanisms to lead to disparities in URI by race/ethnicity. Future research is needed to better understand these mechanisms and to identify modifiable aspects that can serve as targets for intervention.

51

School Shootings and Mental Health in the United States

Camerin Rencken¹, Alice Ellyson^{2,3}, Isaac Rhew^{4,5}, Carol A. Davis^{6,7} and Ali Rowhani-Rahbar^{1,4}

¹University of Washington; ²Firearm Injury and Policy Research Program, Seattle, WA, USA; ³Department of Pediatrics, University of Washington, Seattle, WA, USA; ⁴Department of Epidemiology, University of Washington, Seattle, WA, USA; ⁵Department of Psychiatry and Behavioral Sciences, University of Washington, Seattle, WA, USA; ⁶College of Education, University of Washington, Seattle, WA, USA and ⁷School Mental Health Assessment, Research, and Training Center, University of Washington, Seattle, WA, USA

OBJECTIVES/GOALS: It is estimated that 357,000 children have experienced a school shooting since 1999, yet due to limitations in the firearm violence field broadly, the sequelae are not well understood. The objective of this work is to examine the mental health impacts of school shootings, providing insight into the lasting effects of firearm violence on our communities. **METHODS/STUDY POPULATION:** We will first conduct a quasi-experimental study using controlled interrupted time series with repeated cross-sectional data to assess school shootings' impact on US mental health. School shooting data is from the K-12 School Shooting Database, and mental health data will be collected via the Behavioral Risk Factor Surveillance System. Second, we will conduct focus groups with community organizations, school administrators, and the public. Interview guides will be developed to explore the mental health impacts of school shootings, to guide the quantitative results interpretation, and assess educational materials' usefulness. Qualitative analysis will occur in NVivo software with codebook refinement through thematic analysis. Results will be triangulated through convergence coding. **RESULTS/ANTICIPATED RESULTS:** This research is situated within the context of the pervasive mental health challenges in the US, where mental illness poses significant health, social, and economic burdens. Thus, we anticipate finding an association between school shootings and decreased self-reported mental well-being among US adults. Literature suggests that there may be a stronger association among specific subgroups, such as parents with school-aged children or individuals living in close proximity to such incidents. We expect to find heterogeneity in the effect estimate based on school shooting attributes, such as the number of casualties. Through focus groups, we anticipate furthering our comprehension of the broad-ranging effects of school shootings on less quantifiable outcomes and the unique trajectories of recovery. **DISCUSSION/SIGNIFICANCE:** This project will contribute needed information on the impact of school shootings and mental health and assist in reducing the frequency and impact of school shootings. Furthermore, we aim to extend our findings beyond the scientific community, translating them into educational resources advocating for policy and public health interventions.

54

Identifying Geographic Clusters of H. pylori Associated Metastatic Early-Onset Gastric Cancer: A case-control study in Los Angeles

Jessica Sheth Bhutada¹, Syma Iqbal², Myles Cockburn², Chanita Hughes Halbert² and David Freyer¹

¹Children's Hospital Los Angeles / University of Southern California and ²University of Southern California

OBJECTIVES/GOALS: More young adults (age <50 years) are diagnosed with metastatic gastric cancer (mGC) every year. We will evaluate the association between environmental risk factors (including historical racial residential segregation) clinical, pathologic, molecular features and H.pylori associated early-onset mGC (mE-GC). **METHODS/STUDY POPULATION:** This retrospective matched case-control study of patients (1:2 by diagnosis year) with mGC (early-onset [E-GC; <50 years]; vs older-onset [O-GC; >50 years]) from 2000-2022 from the Los Angeles Cancer Surveillance Program (LA-CSP) will be enriched by a chart-abstracted cohort from USC Norris Comprehensive Cancer (NCCC). This annotated database captures sociodemographic, medical, and pathologic features of patients treated for mGC at NCCC. It will link to LA-CSP data exploring neighborhood features (obesity rate, poverty,