

ON A THEOREM OF ERDŐS AND SZEKERES

M. V. Subbarao

The theorem of Erdos and Szekeres [1] referred to in the title may be stated as follows.

THEOREM A. Given a sequence of r distinct real numbers such that the number of terms of every decreasing subsequence is at most m , then there exists an increasing subsequence of more than n terms, where n is the largest integer less than r/m .

An extremely simple and elegant proof of the theorem was given by A. Seidenberg [2]. This note is intended to point out that a result analogous to the above holds under a more general setting.

Let S be a finite sequence of elements, $S = \{a_1, a_2, \dots, a_r\}$ on which a binary relation R is defined such that, for every distinct pair a, b of elements of S , exactly one of the relations $a R b$, $b R a$ holds. We do not require R to be transitive - that is, $a R b$ and $b R c$ may hold, but not $a R c$.

A subsequence $\{b_1, b_2, \dots, b_s\}$ of S is said to be an ascending R -subsequence with first term b_1 provided either $s = 1$, or $s > 1$ and $b_i R b_{i+1}$ holds for $i = 1, 2, \dots, s-1$. It is said to be a descending R -subsequence with last term b_1 if either $s = 1$, or if $s > 1$ and $b_i R b_{i-1}$ holds for $i = s, s-1, \dots, 2$. Thus only single term subsequences are both ascending and descending R -subsequences. Finally, the "length" of a subsequence is defined to be the number of its elements.

We now have

THEOREM B. Let S be a finite sequence of r abstract elements, on which a binary relation R is defined such that for distinct elements a, b in S , exactly one of the properties $a R b$, $b R a$ holds. If every ascending R -subsequence of S has a length not exceeding m , then there exists a descending R -subsequence of length not less than r/m .

The proof, which is essentially similar to Seidenberg's proof of Theorem A, is as follows:

To each element a_i of S we associate an ordered pair of positive integers (m_i, n_i) where m_i (n_i) is the largest of the lengths of all ascending R -subsequences with first term a_i (descending R -subsequences with last term a_i). These integers m_i, n_i exist for all

i ($i = 1, 2, \dots, r$) and are ≥ 1 . Further, for any two distinct elements a_i, a_j of S , the corresponding ordered pairs (m_i, n_i) and (m_j, n_j) are distinct, for if $i < j$ and if $a_i R a_j$ holds, we have $m_i \geq m_j + 1$, while, if $a_j R a_i$ holds, we have $n_i \geq n_j + 1$. The total number of ordered pairs is then r , the number of elements in S . Let $t = \text{Max}_{1 \leq i \leq r} n_i$. Since $\text{Max}_{1 \leq i \leq r} m_i \leq m$, it follows that the number of all the possible ordered pairs (m_i, n_i) is $\leq mt$, and hence $r \leq mt$. Thus, $t \geq r/m$ and the theorem is proved.

REFERENCES

1. P. Erdős and G. Szekeres, A combinatorial problem in geometry. *Composite Math.* 2 (1935) 463-470.
2. A. Seidenberg, A simple proof of a theorem of Erdős and Szekeres. *J. London Math. Soc.* 34 (1959) 352.

University of Missouri
 University of Alberta