

The Journal of Laryngology and Otology

EDITED BY
WALTER HOWARTH

ASSISTANT EDITOR
G. H. BATEMAN

Contents

CANCER OF THE BASE OF THE TONGUE: TREATMENT BY RADIOTHERAPY	M. LEDERMAN ✓
OUR OBSERVATIONS ON THE ENDOLARYNGEAL APPLI- CATION OF RADIOACTIVE ISOTOPES	IVO PADOVAN ✓
THE SO-CALLED JUVENILE ANGIO-FIBROMA OF THE NASOPHARYNX	D. A. OSBORN ✓
AN OPERATION FOR RE-FORMING THE OVAL WINDOW	J. MİDOŃSKI ✓
SECRETORY CYSTS OF THE MAXILLARY ANTRUM AND THEIR RELATION TO THE DEVELOPMENT OF ANTROCHOANAL POLYPI	C. PETER MILLS ✓
MODIFICATIONS IN THE TECHNIQUE OF FENESTRATION	FRANCISCO ANTOLI-CANDELA ✓
CLINICAL RECORDS—	
ON THE TREATMENT OF PAPILOMA OF THE LARYNX IN CHILDREN	JØRGEN ARNVIG ✓
MÉNIÈRE'S SYNDROME IN A SEVEN-YEAR-OLD GIRL	HENNING SØRENSEN ✓
GENERAL NOTES	

London
Headley Brothers Ltd
109 Kingsway WC2

Annual Subscription £4/4/0 net, U.S.A. \$13

Monthly, 10/- net post free

The Journal of Laryngology and Otology

CONTENTS

	PAGE
CANCER OF THE BASE OF THE TONGUE: TREATMENT BY RADIO-THERAPY. M. Lederman (London)	279
OUR OBSERVATIONS ON THE ENDOLARYNGEAL APPLICATION OF RADIOACTIVE ISOTOPES. Ivo Padovan (Zagreb)	289
THE SO-CALLED JUVENILE ANGIO-FIBROMA OF THE NASOPHARYNX. D. A. Osborn (London)	295
AN OPERATION FOR RE-FORMING THE OVAL WINDOW. J. Miodonski (Cracow, Poland)	317
SECRETORY CYSTS OF THE MAXILLARY ANTRUM AND THEIR RELATION TO THE DEVELOPMENT OF ANTROCHOANAL POLYPLI. C. Peter Mills (London)	324
MODIFICATIONS IN THE TECHNIQUE OF FENESTRATION. Francisco Antoli-Candela (Madrid)	335
CLINICAL RECORDS—	
On the Treatment of Papilloma of the Larynx in Children. Jørgen Arnvig (Søndeberg, Denmark)	343
Ménière's Syndrome in a Seven-Year-Old Girl. Henning Sørensen (Aarhus, Denmark)	346
GENERAL NOTES	351

For Advertisement space in this Journal apply to:
HEADLEY BROTHERS Ltd, 109 Kingsway London WC 2

The Journal of Laryngology and Otology

(FOUNDED in 1887 by MORELL MACKENZIE and NORRIS WOLFENDEN)

EDITED BY
WALTER HOWARTH

ASSISTANT EDITOR
G. H. BATEMAN

1. Original Articles are accepted on the condition that they have not been published elsewhere.

2. Manuscripts should be typewritten, on one side only of the paper, and well spaced. Captions to illustrations should be typed on a separate sheet and sent at the same time as original photographs, etc.

The Harvard system of recording references should be used, e.g. GREEN, C., and BROWN, D. (1951) *J. Laryng.*, 65, 33. Abbreviations of Journals should follow the style recommended in *World Medical Periodicals*, published by World Health Organization 1952.

It is most important that authors should verify *personally* the accuracy of every reference before submitting a paper for publication.

3. Galley proofs and engraver's proofs of illustrations are sent to the author. Corrections, which should be kept to a minimum, must be clearly marked, and no extra matter added. Proofs should be returned within 5 days.

4. Illustration blocks are provided free up to the limit of £10 per article; beyond this authors are expected to pay half the cost. Coloured illustrations will be charged in full to authors.

Blocks will normally be held by the Printers for three years after which they will be destroyed. Any author who has borne a part of the cost of his blocks is entitled to have these returned to him, but a request for this must be sent within three years of the appearance of the article, to HEADLEY BROTHERS LTD., 109 Kingsway London, WC2.

5. Orders for reprints should be sent when returning galley proofs, and for this purpose special forms are supplied.

6. Authors of original communications on Oto-Laryngology in other journals are invited to send a copy, or two reprints, to the *Journal of Laryngology*. If they are willing, at the same time, to submit their own abstract (in English, French, Italian, or German) it will be welcomed.

7. Editorial communications may be addressed to THE EDITOR, *Journal of Laryngology*, c/o HEADLEY BROTHERS LTD., 109 Kingsway, London, WC2.

8. The annual subscription is four guineas sterling (U.S.A. \$13) post free, and is payable in advance.


9. Single copies will be on sale at 10s. od. each; copies of parts up to Vol. LXIII may be purchased at 4s. each.

10. All subscriptions, advertising and business communications should be sent to the publishers, HEADLEY BROTHERS LTD., 109 KINGSWAY LONDON, WC2.

United States of America

Orders for this *Journal* may be sent through local bookseller, or to STECHERT-HAFNER, INC., 31-33 East 10th Street, New York, or direct to the publishers, HEADLEY BROTHERS LTD., 109 KINGSWAY LONDON, WC2, England.

**When a
hearing
aid is
advised . .**



**. . you may confidently
recommend your patients
to S. C. INGRAM**


- ★ They can try the different makes of hearing aids
- ★ They get impartial advice on all makes of hearing aids
- ★ Their personal needs are given individual and continued attention
- ★ A full report on each fitting recommended is sent to the patient's ear specialist
- ★ They can compare at one consultation the different makes one with another

S. C. INGRAM Independent Hearing Aid Consultant
On the approved List of the National Institute for the Deaf
 2 SHEPHERD STREET, SHEPHERD MARKET, LONDON, W.1
 Hyde Park 9042

Please mention *The Journal of Laryngology and Otology* when replying to advertisements

**THE
HALLPIKE—**

**BLACKMORE
EAR
MICROSCOPE***



USES

DIAGNOSIS

The excellent optical performance of the Hallpike Ear Microscope gives the user greatly increased confidence in the diagnosis of abnormal conditions; and many disorders may be recognized at an earlier stage than has hitherto been possible with "headlamp and speculum" techniques. Thus slight pulsation of swellings and vascular tumours—of high diagnostic significance and almost impossible to detect by conventional methods—can readily be detected under this microscope.

Use of the Siegle attachment permits observation of small bubbles present in fluid collections within the tympanum; it is also invaluable for demonstrating minute perforations. Where defects of the tympanic membrane are present, the middle-ear cavity can be scrutinized minutely; the detailed structure of the stapes, stapedius tendon, round window and Eustachian orifice—not normally visible by ordinary methods of examination—are seen with quite remarkable clarity when the Ear Microscope is used.

TREATMENT

Manipulative procedures may be undertaken with greater confidence and precision if the Ear Microscope and its special instruments are used.

The currettes enable adherent wax and debris to be removed with the greatest precision from the walls of the deep meatus or from the tympanic membrane itself. Cholesteatomatous masses or neoplasms may be examined critically and biopsy specimens obtained by means of the currettes. These may also be used as delicate probes for palpation of the incus and head of the stapes in certain cases of otosclerosis with retraction and atrophy of the tympanic membrane; in this way unmistakable evidence may be obtained of bony fixation of the footplate. Or again, it is sometimes possible to identify by palpation defects of the bony tympanic ring or meatal wall which may underlie granulations.

Myringotomy or removal of foreign bodies can be undertaken with surprisingly high precision because the optics of the Ear Microscope and the specially angled myringotome blade eliminate all shadows so that the surgeon can see clearly where and what he is cutting. The outfit is invaluable also for post-operative examinations and treatment of mastoid cavities. Granulations can be cauterized and exposed areas of dura mater carefully palpated.

**J. Laryng.* (1953) 67, 108.

**39 WIGMORE STREET
LONDON W.1**




**5241 WHITBY AVENUE
PHILADELPHIA 43, Pa.,
U.S.A.**

Please mention *The Journal of Laryngology and Otology* when replying to advertisements

In schools for the DEAF— multitone Telesonic INDUCTION LOOP SYSTEM brings new advantages

- 1 Can be installed in adjoining classrooms without risk of 'spill over'.
- 2 Combines good hearing with complete freedom of movement. Unaffected by distance between pupil and teacher.
- 3 Clear speech at high amplification.
- 4 Any number of pupils may use the same installation.
- 5 The child can use the instrument as a normal hearing aid outside school.



The Kindergarten Class at the London Residential School for Jewish Deaf Children where all classrooms have been installed with the Multitone Telesonic Induction Loop System

Write for further details to:

multitone

ELECTRIC COMPANY LTD., 12-20 UNDERWOOD STREET, LONDON, N.1.

CRC M23

Please mention *The Journal of Laryngology and Otology* when replying to advertisements


EAR WAX

Removed this easy way

The removal of wax from the external auditory meatus has, in the past, normally entailed attendance by the patient for diagnosis and for the prescription of a suitable loosening agent, and a second attendance a few days later for syringing.

Now, by the use of Cerumol Ear Drops, wax can be removed in most cases at one visit. A few drops of Cerumol can be instilled into the ear and, while another patient is being attended to, the soft cerumen dissolves and the harder wax disimpacts. The wax can then be removed by gentle syringing or with cotton wool. The wax may even be found to run out of the ear on its own accord, in which case patients themselves may instil Cerumol at home, obviating further attendances. Cerumol is anti-bacterial, non-irritating and harmless to the lining of the external auditory meatus or the tympanic membrane.

Cerumol is included in Category No. 4 of the M.O.H. classified list and may be prescribed on N.H.S. Form E.C.10.



CERUMOL
REGD. TRADE MARK

EAR DROPS

for the easier removal of wax

If you wish to test for yourself and have not received recently a 10. c.c. vial please write or telephone direct to:

LABORATORIES FOR APPLIED BIOLOGY LTD.,
91, AMHURST PARK, LONDON, N.16 Tel.: STA 2252


PACKS For Surgery Use:
10 c.c. vial — separate
dropper included
(Basic N.H.S. price 2/8)
for Hospital Use: 2 oz.
and 10 oz. bottles.

Please mention *The Journal of Laryngology and Otology* when replying to advertisements

An outstanding accomplishment
in hearing aid design

Amplivox

"C" SERIES HEARING AIDS



FACILITIES

Individual fitting related to the Audiogram by means of selective compensation. Tone control giving low frequency attenuation combined with an overall response extending to 4000 c/s provides excellent speech discrimination in cases of perceptible deafness.

Two-stage adjustable automatic volume compression.

SPECIFICATION

4 stage transistor circuit. Stable operation from 0° to 40°C.

Miniature earphone with extended response. Magnetic microphone. Telephone inductor. Mercury cell battery —life 100 hours. Size $2\frac{1}{8}$ " x $1\frac{1}{16}$ " x $\frac{15}{32}$ "

Weight 1 oz.

Mirror gold finish.

AMPLIVOX LTD . LONDON . ENGLAND

80 New Bond Street, London, W.1. Hyde Park 9888

WORLD-WIDE DISTRIBUTION AND SERVICE

Please mention *The Journal of Laryngology and Otology* when replying to advertisements



'OTOSPORIN' clears infected ears

The modern method of treatment in otitis externa, chronic suppurative otitis media, and infected mastoid cavities, is 'Otosporin' brand Drops.

'Otosporin' is an aqueous suspension containing both neomycin (5 mgm. per c.c.) and polymyxin B sulphate (10,000 units per c.c.) which are together effective against practically all the bacteria found in ear infections. They are unlikely to induce bacterial resistance or skin sensitisation, and neither of them gives rise to cross-resistance or cross-sensitisation to penicillin and other antibiotics. Their effectiveness is visibly enhanced by the hydrocortisone (5 mgm. per c.c.) in 'Otosporin'; this, by reducing inflammation, not only relieves pain, but provides greater access for the antibiotics.

'OTOSPORIN'^{BRAND} drops
issued in bottles of 5 c.c.



BURROUGHS WELLCOME & CO. (The Wellcome Foundation Ltd.) **LONDON**

Please mention *The Journal of Laryngology* when replying to advertisements