

MOCK THETA DOUBLE SUMS

JEREMY LOVEJOY

CNRS, LIAFA, Université Denis Diderot - Paris 7, Case 7014, 75205 Paris Cedex 13, France
e-mail: lovejoy@math.cnrs.fr

and ROBERT OSBURN

School of Mathematics and Statistics, University College Dublin, Belfield, Dublin 4, Ireland
IHÉS, Le Bois-Marie, 35, route de Chartres, F-91440 Bures-sur-Yvette, France
e-mails: robert.osburn@ucd.ie, osburn@ihes.fr

(Received 16 January 2015; revised 27 July 2015; accepted 5 January 2016;
first published online 10 June 2016)

Abstract. We prove a general result on Bailey pairs and show that two Bailey pairs of Bringmann and Kane are special cases. We also show how to use a change of base formula to pass from the pairs of Bringmann and Kane to pairs used by Andrews in his study of Ramanujan's seventh order mock theta functions. We derive several more Bailey pairs of a similar type and use these to construct a number of new q -hypergeometric double sums which are mock theta functions. Finally, we prove identities between some of these mock theta double sums and classical mock theta functions.

2010 *Mathematics Subject Classification.* Primary: 33D15; Secondary: 11F37.

1. Introduction.

1.1. Bailey pairs. A *Bailey pair* relative to a is a pair of sequences $(\alpha_n, \beta_n)_{n \geq 0}$ satisfying

$$\beta_n = \sum_{k=0}^n \frac{\alpha_k}{(q)_{n-k}(aq)_{n+k}}, \quad (1)$$

or equivalently

$$\alpha_n = (1 - aq^{2n}) \sum_{j=0}^n \frac{(aq)_{n+j-1} (-1)^{n-j} q^{\binom{n-j}{2}}}{(q)_{n-j}} \beta_j. \quad (2)$$

Here, we have used the standard q -hypergeometric notation,

$$(a)_n = (a; q)_n = \prod_{k=1}^n (1 - aq^{k-1}),$$

valid for $n \in \mathbb{N} \cup \{\infty\}$. The *Bailey lemma* says that if (α_n, β_n) is a Bailey pair relative to a , then so is (α'_n, β'_n) , where

$$\alpha'_n = \frac{(\rho_1)_n(\rho_2)_n(aq/\rho_1\rho_2)^n}{(aq/\rho_1)_n(aq/\rho_2)_n} \alpha_n \tag{3}$$

and

$$\beta'_n = \sum_{k=0}^n \frac{(\rho_1)_k(\rho_2)_k(aq/\rho_1\rho_2)_{n-k}(aq/\rho_1\rho_2)^k}{(aq/\rho_1)_n(aq/\rho_2)_n(q)_{n-k}} \beta_k. \tag{4}$$

A useful limiting form of the Bailey lemma is found by putting (3) and (4) into (1) and letting $n \rightarrow \infty$, giving

$$\sum_{n \geq 0} (\rho_1)_n(\rho_2)_n(aq/\rho_1\rho_2)^n \beta_n = \frac{(aq/\rho_1)_\infty(aq/\rho_2)_\infty}{(aq)_\infty(aq/\rho_1\rho_2)_\infty} \sum_{n \geq 0} \frac{(\rho_1)_n(\rho_2)_n(aq/\rho_1\rho_2)^n}{(aq/\rho_1)_n(aq/\rho_2)_n} \alpha_n. \tag{5}$$

For more on Bailey pairs and the Bailey lemma, see [1, 2, 17].

This paper has its origins in the following two Bailey pairs discovered by Bringmann and Kane [9]. First, (a_n, b_n) is a Bailey pair relative to 1, where

$$a_{2n} = (1 - q^{4n})q^{2n^2-2n} \sum_{j=-n}^{n-1} q^{-2j^2-2j}, \tag{6}$$

$$a_{2n+1} = -(1 - q^{4n+2})q^{2n^2} \sum_{j=-n}^n q^{-2j^2}, \tag{7}$$

and

$$b_n = \frac{(-1)^n(q; q^2)_{n-1}}{(q)_{2n-1}} \chi(n \neq 0)^1, \tag{8}$$

and second, (α_n, β_n) is a Bailey pair relative to q , where

$$\alpha_{2n} = \frac{1}{1 - q} \left(q^{2n^2+2n} \sum_{j=-n}^{n-1} q^{-2j^2-2j} + q^{2n^2} \sum_{j=-n}^n q^{-2j^2} \right), \tag{9}$$

$$\alpha_{2n+1} = -\frac{1}{1 - q} \left(q^{2n^2+4n+2} \sum_{j=-n}^n q^{-2j^2} + q^{2n^2+2n} \sum_{j=-n-1}^n q^{-2j^2-2j} \right), \tag{10}$$

and

$$\beta_n = \frac{(-1)^n(q; q^2)_n}{(q)_{2n+1}}. \tag{11}$$

¹As usual, $\chi(X) = 1$ if X is true and 0 if X is false.

These are highly reminiscent of three Bailey pairs discovered by Andrews [3] in his study of Ramanujan’s seventh order mock theta functions. Namely, he showed that $(\mathcal{A}_n(0), \mathcal{B}_n(0))$ and $(\mathcal{A}_n(1), \mathcal{B}_n(1))$ form Bailey pairs relative to 1, where

$$\mathcal{A}_{2n}(0) = q^{3n^2+n} \sum_{j=-n}^n q^{-j^2} - q^{3n^2-n} \sum_{j=-n+1}^{n-1} q^{-j^2}, \tag{12}$$

$$\mathcal{A}_{2n+1}(0) = -q^{3n^2+4n+1} \sum_{j=-n-1}^n q^{-j^2-j} + q^{3n^2+2n} \sum_{j=-n}^{n-1} q^{-j^2-j}, \tag{13}$$

$$\mathcal{B}_n(0) = \frac{1}{(q^{n+1})_n}, \tag{14}$$

$$\mathcal{A}_{2n}(1) = -(1 - q^{4n})q^{3n^2-2n} \sum_{j=-n}^{n-1} q^{-j^2-j}, \tag{15}$$

$$\mathcal{A}_{2n+1}(1) = (1 - q^{4n+2})q^{3n^2+n} \sum_{j=-n}^n q^{-j^2}, \tag{16}$$

and

$$\mathcal{B}_n(1) = \frac{1}{(q^n)_n} \chi(n \neq 0), \tag{17}$$

while $(\mathcal{A}_n(2), \mathcal{B}_n(2))$ is a Bailey pair relative to q , where

$$\mathcal{A}_{2n}(2) = \frac{1}{1 - q} \left(q^{3n^2+2n} \sum_{j=-n}^{n-1} q^{-j^2-j} + q^{3n^2+n} \sum_{j=-n}^n q^{-j^2} \right), \tag{18}$$

$$\mathcal{A}_{2n+1}(2) = -\frac{1}{1 - q} \left(q^{3n^2+5n+2} \sum_{j=-n}^n q^{-j^2} + q^{3n^2+4n+1} \sum_{j=-n-1}^n q^{-j^2-j} \right), \tag{19}$$

and

$$\mathcal{B}_n(2) = \frac{1}{(q^{n+1})_{n+1}}. \tag{20}$$

Our first goal in this paper is to prove the following results, which will lead to more Bailey pairs like those of Bringmann–Kane and Andrews. Note that Theorem 1.3 is simply an application of Theorem 1.1 followed by an application of Theorem 1.2.

THEOREM 1.1. *If (α_n, β_n) is a Bailey pair relative to 1 with $\alpha_0 = \beta_0 = 1$, then (α'_n, β'_n) is also a Bailey pair relative to 1, where $\alpha'_0 = \beta'_0 = 0$,*

$$\alpha'_{2n} = -(1 - q^{4n})q^{2n^2-2n} \sum_{j=0}^{n-1} q^{-2j^2-2j} \alpha_{2j+1}, \tag{21}$$

$$\alpha'_{2n+1} = -(1 - q^{4n+2})q^{2n^2} \sum_{j=0}^n q^{-2j^2} \alpha_{2j}, \tag{22}$$

and for $n \geq 1$,

$$\beta'_n = -\frac{\beta_{n-1}}{1 - q^{2n-1}}. \tag{23}$$

THEOREM 1.2. *Suppose that (α_n, β_n) is a Bailey pair relative to 1 with $\alpha_0 = \beta_0 = 0$. Then (α'_n, β'_n) is a Bailey pair relative to q , where*

$$\alpha'_n = \frac{1}{1 - q} \left(-\frac{\alpha_{n+1}}{1 - q^{2n+2}} + \frac{q^{2n}\alpha_n}{1 - q^{2n}} \right) \tag{24}$$

and

$$\beta'_n = -\beta_{n+1}. \tag{25}$$

THEOREM 1.3. *If (α_n, β_n) is a Bailey pair relative to 1 with $\alpha_0 = \beta_0 = 1$, then (α''_n, β''_n) is a Bailey pair relative to q , where*

$$\alpha''_{2n} = \frac{1}{1 - q} \left(q^{2n^2} \sum_{j=0}^n q^{-2j^2} \alpha_{2j} - q^{2n^2+2n} \sum_{j=0}^{n-1} q^{-2j^2-2j} \alpha_{2j+1} \right), \tag{26}$$

$$\alpha''_{2n+1} = \frac{1}{1 - q} \left(q^{2n^2+2n} \sum_{j=0}^n q^{-2j^2-2j} \alpha_{2j+1} - q^{2n^2+4n+2} \sum_{j=0}^n q^{-2j^2} \alpha_{2j} \right), \tag{27}$$

and

$$\beta''_n = \frac{\beta_n}{1 - q^{2n+1}}. \tag{28}$$

An application of Theorem 1.1 or 1.3 to a “typical” Bailey pair (from Slater’s list [16], for example) will give a positive definite quadratic form in the power of q occurring in α_n . However, there are a few cases where we obtain an indefinite quadratic form. For example, using Theorems 1.1 and 1.3 and the following Bailey pair relative to 1 from Slater’s list [16, p. 468],

$$\alpha_n = \begin{cases} 1, & \text{if } n = 0, \\ 2(-1)^n, & \text{otherwise} \end{cases}$$

and

$$\beta_n = \frac{(-1)^n}{(q^2; q^2)_n},$$

we recover the Bailey pairs of Bringmann and Kane in (6)–(11). Some other examples are recorded in Corollaries 2.2–2.4. Andrews’ Bailey pairs in (15)–(20) do not seem to be simple applications of Theorems 1.1–1.3, but they can be deduced from (6)–(11) using a change of base formula. We discuss this in Section 3. For another treatment of these pairs, see [6].

1.2. Mock theta functions. An important difference between the pairs of Andrews and those of Bringmann–Kane is that the former yield mock theta functions when substituted into (5), while the latter do not. However, as we showed in [13], the Bailey pairs of Bringmann and Kane *do* give rise to mock theta functions after an appropriate application of the Bailey lemma. These mock theta functions are q -hypergeometric double sums.

To recall them, we need some special functions. We use the classical theta series

$$j(x, q) := \sum_{n \in \mathbb{Z}} (-x)^n q^{\binom{n}{2}} = (x)_\infty (q/x)_\infty (q)_\infty,$$

and for brevity, we write $J_m := J_{m,3m}$ with $J_{a,m} := j(q^a, q^m)$, and $\bar{J}_{a,m} := j(-q^a, q^m)$. We also use the Hecke-type series

$$f_{a,b,c}(x, y, q) := \left(\sum_{r,s \geq 0} - \sum_{r,s < 0} \right) (-1)^{r+s} x^r y^s q^{a\binom{r}{2} + brs + c\binom{s}{2}}, \tag{29}$$

which is an indefinite theta series when $ac < b^2$. Finally, we employ the Appell–Lerch series

$$m(x, q, z) := \frac{1}{j(z, q)} \sum_{r \in \mathbb{Z}} \frac{(-1)^r q^{\binom{r}{2}} z^r}{1 - q^{r-1} xz}, \tag{30}$$

where $x, z \in \mathbb{C}^* := \mathbb{C} \setminus \{0\}$ with neither z nor xz an integral power of q . The Appell–Lerch series is a “mock Jacobi form” which specializes to a mock theta function when x and z are of the form $\zeta q^{\frac{m}{n}}$ with ζ a root of unity [18, 19]. Recall that a mock theta function is the holomorphic part of a weight $1/2$ harmonic weak Maass form $f(\tau)$ (as usual, $q := e^{2\pi i \tau}$ where $\tau = x + iy \in \mathbb{H}$) whose image of under the operator $\xi_{\frac{1}{2}} := 2iy^{\frac{1}{2}} \frac{\partial}{\partial \bar{\tau}}$ is a unary theta function [15, 18].

The main result in [13] contains identities equivalent to the following.

THEOREM 1.4. [13, Theorem 1.3]

$$\begin{aligned}
 \mathcal{W}_1(q) &:= \sum_{n \geq 1} \sum_{n \geq j \geq 1} \frac{(-1)_j(q; q^2)_{j-1} (-1)^j q^{n^2 + \binom{j+1}{2}}}{(-q)_n (q)_{n-j} (q)_{2j-1}} \\
 &= \frac{-2q^2}{(q)_\infty} f_{3,5,3}(q^5, q^5, q) \\
 &= 4m(-q^{17}, q^{48}, q^{24}) - 4q^{-5}m(-q, q^{48}, q^{24}) - 2q^2 \frac{J_8 J_{12} J_{96} J_{7,16} \bar{J}_{4,24} J_{6,48} J_{30,96}}{J_{24} J_{48} J_{3,8} J_{2,12} J_{14,96} J_{46,96}}, \tag{31}
 \end{aligned}$$

$$\begin{aligned}
 \mathcal{W}_2(q) &:= \sum_{n \geq 1}^* \sum_{n \geq j \geq 1} \frac{(q; q^2)_n (-1)_j (q; q^2)_{j-1} (-1)^{n+j} q^{\binom{j+1}{2}}}{(-q)_n (q)_{n-j} (q)_{2j-1}} \\
 &= \frac{q(q; q^2)_\infty}{(q^2; q^2)_\infty} f_{1,3,1}(-q^2, -q^2, q) \\
 &= 4m(-q, q^8, q^4) + q \frac{J_{1,8}^2 J_{3,8}^3 J_{2,16}}{J_8^4 J_{16}}, \tag{32}
 \end{aligned}$$

$$\begin{aligned}
 \mathcal{W}_3(q) &:= \sum_{n \geq 1} \sum_{n \geq j \geq 1} \frac{(q; q^2)_n (-1; q^2)_j (q^2; q^4)_{j-1} (-1)^{n+j} q^{n^2 + j^2 + j}}{(-q^2; q^2)_n (q^2; q^2)_{n-j} (q^2; q^2)_{2j-1}} \\
 &= \frac{2q^3 (q; q^2)_\infty}{(q^2; q^2)_\infty} f_{1,2,1}(-q^7, -q^7, q^4) \\
 &= 4m(-q, q^{12}, q^4) + 2q^3 \frac{\bar{J}_{1,12}^2}{J_{1,4}}, \tag{33}
 \end{aligned}$$

$$\begin{aligned}
 \mathcal{W}_4(q) &:= \sum_{n \geq 0} \sum_{n \geq j \geq 0} \frac{(-q)_j (q; q^2)_j (-1)^j q^{n^2 + n + \binom{j+1}{2}}}{(-q)_n (q)_{n-j} (q)_{2j+1}} \\
 &= \frac{1}{(q)_\infty} f_{3,5,3}(q^3, q^3, q) \\
 &= -2q^{-4}m(-q^5, q^{48}, q^{24}) - 2q^{-2}m(-q^{11}, q^{48}, q^{24}) \\
 &\quad + \frac{J_8 J_{12} J_{96} J_{3,16} \bar{J}_{4,24} J_{6,48} J_{18,96} J_{30,96}}{J_{24} J_{48} J_{1,8} J_{2,12} J_{6,96} J_{26,96} J_{38,96}}. \tag{34}
 \end{aligned}$$

In particular, $\mathcal{W}_1(q) - \mathcal{W}_4(q)$ are mock theta functions. We remark that the series defining $\mathcal{W}_2(q)$ does not converge. However, similar to the classical sixth order mock theta function $\mu(q)$ [5], the sequence of even partial sums and the sequence of odd partial sums both converge. We define $\mathcal{W}_2(q)$ as the average of these two values. This averaging is denoted here and throughout by the notation \sum^* .

The second goal of this paper is to use Bailey pairs arising from Theorems 1.1 and 1.3 to obtain many more mock theta functions like $\mathcal{W}_1(q) - \mathcal{W}_4(q)$. Just as with the pairs of Bringmann and Kane, this first requires one application of the Bailey lemma, and so the mock theta functions we obtain are q -hypergeometric double sums. We record these mock theta functions in three separate results, corresponding to three sets of

Bailey pairs. We first express the double sums in terms of the indefinite theta series (29) and then in terms of the Appell–Lerch series (30).

THEOREM 1.5. *The following are mock theta functions.*

$$\begin{aligned}
 \mathcal{M}_1(q) &:= \sum_{n \geq 1} \sum_{n \geq j \geq 1} \frac{(-1)_j (-1)^j q^{n^2 + \binom{j}{2}}}{(-q)_n (q)_{n-j} (q^2; q^2)_{j-1} (1 - q^{2j-1})} \\
 &= -\frac{2q}{(q)_\infty} f_{3,5,3}(q^4, q^6, q) \\
 &= 4q^{-3} m(-q^7, q^{48}, q^{24}) + 4m(-q^{25}, q^{48}, q^{-24}) - 2 \\
 &\quad - 2 \frac{J_8 J_{12} J_{96} J_{1,16} \bar{J}_{4,24} J_{6,48} J_{18,96}}{J_{24} J_{48} J_{3,8} J_{2,12} J_{2,96} J_{34,96}}, \tag{35}
 \end{aligned}$$

$$\begin{aligned}
 \mathcal{M}_2(q) &:= \sum_{n \geq 1}^* \sum_{n \geq j \geq 1} \frac{(q; q^2)_n (-1)_j (-1)^{n+j} q^{\binom{j}{2}}}{(-q)_n (q)_{n-j} (q^2; q^2)_{j-1} (1 - q^{2j-1})} \\
 &= \frac{(q; q^2)_\infty}{(q^2; q^2)_\infty} f_{1,3,1}(-q, -q^3, q) \\
 &= 4m(-q^5, q^8, q^4) - 2 - \frac{J_{1,8}^3 J_{3,8}^2 J_{6,16}}{J_8^4 J_{16}}, \tag{36}
 \end{aligned}$$

$$\begin{aligned}
 \mathcal{M}_3(q) &:= \sum_{n \geq 1} \sum_{n \geq j \geq 1} \frac{(q; q^2)_n (-1; q^2)_j (-1)^{n+j} q^{n^2 + j^2 - j}}{(-q^2; q^2)_n (q^2; q^2)_{n-j} (q^4; q^4)_{j-1} (1 - q^{4j-2})} \\
 &= \frac{2q(q; q^2)_\infty}{(q^2; q^2)_\infty} f_{1,2,1}(-q^5, -q^9, q^4) \\
 &= 4m(-q^7, q^{12}, q^4) - 2 + 2 \frac{J_{12}^3 \bar{J}_{5,12}}{J_{4,12} \bar{J}_{1,12} \bar{J}_{3,12}}, \tag{37}
 \end{aligned}$$

$$\begin{aligned}
 \mathcal{M}_4(q) &:= \sum_{n \geq 0} \sum_{n \geq j \geq 0} \frac{(-1)^j q^{n^2 + n + \binom{j}{2}}}{(-q)_n (q)_{n-j} (q)_j (1 - q^{2j+1})} \\
 &= \frac{1}{(q)_\infty} f_{3,5,3}(q^2, q^4, q) \\
 &= 2m(-q^{29}, q^{48}, q^{24}) - 1 - 2q^{-1} m(-q^{13}, q^{48}, q^{-24}) \\
 &\quad + q \frac{J_8 J_{12} J_{96}^3 J_{5,16} \bar{J}_{4,24} J_{6,48} J_{18,48}}{J_{24} J_{48}^2 J_{1,8} J_{2,12} J_{10,96} J_{22,96} J_{42,96}}, \tag{38}
 \end{aligned}$$

$$\begin{aligned}
 \mathcal{M}_5(q) &:= \sum_{n \geq 0} \sum_{n \geq j \geq 0} \frac{(-1)^j q^{\binom{n+1}{2} + \binom{j}{2}}}{(q)_{n-j} (q)_j (1 - q^{2j+1})} \\
 &= \frac{(-q)_\infty}{(q)_\infty} f_{1,2,1}(q, q^3, q^2) \\
 &= 2m(q^5, q^6, q^2) - 1 - q \frac{J_6^3}{J_{2,6} J_{3,6}}. \tag{39}
 \end{aligned}$$

THEOREM 1.6. *The following are mock theta functions.*

$$\begin{aligned}
 \mathcal{M}_6(q) &:= \sum_{n \geq 1} \sum_{n \geq j \geq 1} \frac{(-1)^j q^{n^2 + \binom{j+1}{2}}}{(q)_{n-j} (q)_{j-1} (1 - q^{2j-1})} \\
 &= -\frac{q^2}{(q)_\infty} f_{3,7,3}(q^5, q^6, q) \\
 &= m(-q^{49}, q^{120}, q^{-3}) - q^{-3} m(-q^{89}, q^{120}, q^{-3}) + q^{-3} \\
 &\quad + q^{-14} m(-q^{119}, q^{120}, q^3) - q^{-14} \\
 &\quad - q^{-1} m(-q^{79}, q^{120}, q^3) + q^{-1} - q^{-11} \frac{J_{12,48} J_{16,40} J_{2,20} J_{3,40} \bar{J}_{17,40} J_{40}}{J_1 J_{3,120} \bar{J}_{6,40} J_{20} J_{80}} \\
 &\quad + q^{-4} \frac{J_{24,48} J_{1,40} J_{4,40} \bar{J}_{1,40} J_{8,20} \bar{J}_{4,40} J_{18,40} J_{80}}{J_1 J_{3,120} \bar{J}_{6,40} \bar{J}_{2,40} J_{20}^2 J_{40}} \\
 &\quad + q^{-12} \frac{J_{24,48} J_{1,40} J_{4,40} \bar{J}_{1,40} J_{8,20} \bar{J}_{16,40} J_{42,80}^2}{J_1 J_{3,120} \bar{J}_{6,40} \bar{J}_{2,40} J_{20}^2 J_{80}}, \tag{40}
 \end{aligned}$$

$$\begin{aligned}
 \mathcal{M}_7(q) &:= \sum_{n \geq 1} \sum_{n \geq j \geq 1} \frac{(-1)_n (-1)^j q^{\binom{n+1}{2} + \binom{j+1}{2}}}{(q)_{n-j} (q)_{j-1} (1 - q^{2j-1})} \\
 &= -\frac{2q^2 (-q)_\infty}{(q)_\infty} f_{1,3,1}(q^4, q^5, q^2) \\
 &= 2q^{-1} m(-q, q^{16}, q^{-1}) - 2q^{-1} \frac{J_{4,8} J_{16,32} J_{1,16} J_{14,32}}{J_{1,2} \bar{J}_{2,16} \bar{J}_{0,16}}, \tag{41}
 \end{aligned}$$

$$\begin{aligned}
 \mathcal{M}_8(q) &:= 2 \sum_{n \geq 1}^* \sum_{n \geq j \geq 1} \frac{(q; q^2)_n (-1)^{n+j} q^{\binom{j+1}{2}}}{(q)_{n-j} (q)_{j-1} (1 - q^{2j-1})} \\
 &= \frac{(q; q^2)_\infty}{(q^2; q^2)_\infty} f_{1,5,1}(-q^2, -q^3, q) \\
 &= 2m(-q^7, q^{24}, q^6) + 2q^{-2} m(-q, q^{24}, q^{-6}) + q \frac{J_1 J_{3,8} J_{2,16}}{J_2 J_{16}}, \tag{42}
 \end{aligned}$$

$$\begin{aligned}
 \mathcal{M}_9(q) &:= \sum_{n \geq 1} \sum_{n \geq j \geq 1} \frac{(q; q^2)_n (-1)^{n+j} q^{n^2 + j^2 + j}}{(q^2; q^2)_{n-j} (q^2; q^2)_{j-1} (1 - q^{4j-2})} \\
 &= \frac{q^3 (q; q^2)_\infty}{(q^2; q^2)_\infty} f_{1,3,1}(-q^7, -q^9, q^4) \\
 &= m(-q^8, q^{32}, q^{-2}) - q^{-1} \frac{J_{64}^2 J_{28,64}}{J_{32} J_{4,64}} + q^{-1} \frac{J_{8,16} J_{32,64} J_{4,32} J_{24,64}}{\bar{J}_{1,4} \bar{J}_{6,32} \bar{J}_{2,32}}, \tag{43}
 \end{aligned}$$

$$\begin{aligned}
 \mathcal{M}_{10}(q) &:= \sum_{n \geq 1} \sum_{n \geq j \geq 1} \frac{(-1)_{2n} (-1)^j q^{n^2 + j^2 + j}}{(q^2; q^2)_{n-j} (q^2; q^2)_{j-1} (1 - q^{4j-2})} \\
 &= -\frac{2q^3 (-q)_\infty}{(q)_\infty} f_{1,5,1}(q^5, q^7, q^2) \\
 &= -2q^{-1} m(-q^{10}, q^{48}, q^{-2}) - 2q^{-4} m(-q^2, q^{48}, q^{-2})
 \end{aligned}$$

$$\begin{aligned}
 & - 4q^{-3} \frac{J_{8,32} J_{20,48} \bar{J}_{22,48} J_{2,24} J_{6,48} J_{96}}{J_{1,2} \bar{J}_{8,48} \bar{J}_{0,48} J_{24} J_{2,48}} \\
 & + 2q^6 \frac{J_{16,32} J_{4,48} \bar{J}_{2,48} J_{10,24} \bar{J}_{4,48} J_{20,48} J_{96}}{J_{1,2} \bar{J}_{8,48} \bar{J}_{0,48} J_{24}^2 J_{48}} \\
 & + 2q^{-4} \frac{J_{16,32} J_{4,48} \bar{J}_{2,48} J_{10,24} \bar{J}_{20,48} J_{44,96}^2}{J_{1,2} \bar{J}_{8,48} \bar{J}_{0,48} J_{24}^2 J_{96}}, \tag{44}
 \end{aligned}$$

$$\begin{aligned}
 \mathcal{M}_{11}(q) & := \sum_{n \geq 0} \sum_{n \geq j \geq 0} \frac{(-1)^j q^{n^2+n+\binom{j+1}{2}}}{(q)_{n-j}(q)_j(1 - q^{2j+1})} \\
 & = \frac{1}{(q)_\infty} f_{3,7,3}(q^3, q^4, q) \\
 & = q^{-8} m(-q^{17}, q^{120}, q^{-3}) + q^{-6} m(-q^{23}, q^{120}, q^3) - q^{-1} m(-q^{47}, q^{120}, q^3) \\
 & \quad + q^{-12} m(-q^7, q^{120}, q^3) + q^{-9} \frac{J_{12,48} J_{1,40} J_{8,40} \bar{J}_{19,40} J_{6,20} \bar{J}_{12,40} J_{18,40} J_{40}^2}{J_1 J_{3,120} \bar{J}_{14,40} \bar{J}_{10,40} J_{20}^3 J_{80}} \\
 & \quad + q^{-4} \frac{J_{12,48} J_{1,40} J_{8,40} \bar{J}_{19,40} J_{6,20} \bar{J}_{8,40} J_{19,40}^2 \bar{J}_{1,40}^2}{J_1 J_{3,120} \bar{J}_{14,40} \bar{J}_{10,40} J_{20}^3 J_{40} J_{80}} \\
 & \quad - q^{-4} \frac{J_{24,48} J_{1,40} J_{12,40} \bar{J}_{1,40} J_{4,20} \bar{J}_{12,40} J_{18,40} J_{80}}{J_1 J_{3,120} \bar{J}_{14,40} \bar{J}_{10,40} J_{20}^2 J_{40}} \\
 & \quad - q^{-8} \frac{J_{24,48} J_{1,40} J_{12,40} \bar{J}_{1,40} J_{4,20} \bar{J}_{8,40} J_{38,80}^2}{J_1 J_{3,120} \bar{J}_{14,40} \bar{J}_{10,40} J_{20}^2 J_{80}}, \tag{45}
 \end{aligned}$$

$$\begin{aligned}
 \mathcal{M}_{12}(q) & := \sum_{n \geq 0} \sum_{n \geq j \geq 0} \frac{(-q)_n (-1)^j q^{\binom{n+1}{2} + \binom{j+1}{2}}}{(q)_{n-j}(q)_j(1 - q^{2j+1})} \\
 & = \frac{(-q)_\infty}{(q)_\infty} f_{1,3,1}(q^2, q^3, q^2) \\
 & = -q^{-1} m(-q^3, q^{16}, q) + q \frac{J_{4,8} J_{16,32} J_{5,16} J_{6,32}}{J_{1,2} \bar{J}_{6,16} \bar{J}_{4,16}}. \tag{46}
 \end{aligned}$$

THEOREM 1.7. *The following are mock theta functions.*

$$\begin{aligned}
 \mathcal{M}_{13}(q) & := \sum_{n \geq 1} \sum_{n \geq j \geq 1} \frac{(-1)^j q^{n^2+\binom{j}{2}}}{(q)_{n-j}(q)_{j-1}(1 - q^{2j-1})} \\
 & = -\frac{q}{(q)_\infty} f_{3,7,3}(q^4, q^7, q) \\
 & = m(-q^{59}, q^{120}, q^{-9}) - q^{-7} m(-q^{19}, q^{120}, q^{-9}) - q^{-4} m(-q^{29}, q^{120}, q^9) \\
 & \quad - q^{-10} m(-q^{11}, q^{120}, q^{-9}) - q^{-8} \frac{J_{12,48} J_{3,40} J_{16,40} \bar{J}_{17,40} J_{2,20} \bar{J}_{4,40} J_{14,40} J_{40}^2}{J_1 J_{9,120} \bar{J}_{10,40} \bar{J}_{2,40} J_{20}^3 J_{80}} \\
 & \quad - q^{-9} \frac{J_{12,48} J_{3,40} J_{16,40} \bar{J}_{17,40} J_{2,20} \bar{J}_{16,40} J_{17,40}^2 \bar{J}_{3,40}^2}{J_1 J_{9,120} \bar{J}_{10,40} \bar{J}_{2,40} J_{20}^3 J_{40} J_{80}}
 \end{aligned}$$

$$\begin{aligned}
 &+ q^{-2} \frac{J_{24,48} J_{3,40} J_{4,40} \bar{J}_{3,40} J_{8,20} \bar{J}_{4,40} J_{14,40} J_{80}}{J_1 J_{9,120} \bar{J}_{10,40} \bar{J}_{2,40} J_{20}^2 J_{40}} \\
 &+ q^{-10} \frac{J_{24,48} J_{3,40} J_{4,40} \bar{J}_{3,40} J_{8,20} \bar{J}_{16,40} J_{34,80}^2}{J_1 J_{9,120} \bar{J}_{10,40} \bar{J}_{2,40} J_{20}^2 J_{80}}, \tag{47}
 \end{aligned}$$

$$\begin{aligned}
 \mathcal{M}_{14}(q) &:= \sum_{n \geq 1} \sum_{n \geq j \geq 1} \frac{(-1)_n (-1)^j q^{\binom{n+1}{2} + \binom{j}{2}}}{(q)_{n-j} (q)_{j-1} (1 - q^{2j-1})} \\
 &= -\frac{2q(-q)_\infty}{(q)_\infty} f_{1,3,1}(q^3, q^6, q^2) \\
 &= 2m(-q^7, q^{16}, q^{-3}) - 2 \frac{J_{4,8} J_{16,32} J_{1,16} J_{14,32}}{J_{1,2} \bar{J}_{2,16} \bar{J}_{4,16}}, \tag{48}
 \end{aligned}$$

$$\begin{aligned}
 \mathcal{M}_{15}(q) &:= 2 \sum_{n \geq 1}^* \sum_{n \geq j \geq 1} \frac{(q; q^2)_n (-1)^{n+j} q^{\binom{j}{2}}}{(q)_{n-j} (q)_{j-1} (1 - q^{2j-1})} \\
 &= \frac{(q; q^2)_\infty}{(q^2; q^2)_\infty} f_{1,5,1}(-q, -q^4, q) \\
 &= 2m(-q^{13}, q^{24}, q^2) - 2q^{-1} m(-q^5, q^{24}, q^2) + \frac{J_1 J_{1,8} J_{6,16}}{J_2 J_{16}}, \tag{49}
 \end{aligned}$$

$$\begin{aligned}
 \mathcal{M}_{16}(q) &:= \sum_{n \geq 1} \sum_{n \geq j \geq 1} \frac{(q; q^2)_n (-1)^{n+j} q^{n^2+j^2-j}}{(q^2; q^2)_{n-j} (q^2; q^2)_{j-1} (1 - q^{4j-2})} \\
 &= \frac{q(q; q^2)_\infty}{(q^2; q^2)_\infty} f_{1,3,1}(-q^5, -q^{11}, q^4) \\
 &= -q^{-1} m(-q^8, q^{32}, q^{-6}) + \frac{1}{2} + \frac{J_{32}^3 J_{10,32} \bar{J}_{6,32}}{J_{6,32} J_{16,32} \bar{J}_{0,32} \bar{J}_{10,32}} + q^{-1} \frac{J_{8,16} J_{32,64} J_{4,32} J_{24,64}}{\bar{J}_{1,4} \bar{J}_{2,32} \bar{J}_{10,32}}, \tag{50}
 \end{aligned}$$

$$\begin{aligned}
 \mathcal{M}_{17}(q) &:= \sum_{n \geq 1} \sum_{n \geq j \geq 1} \frac{(-1)_{2n} (-1)^j q^{n^2+j^2-j}}{(q^2; q^2)_{n-j} (q^2; q^2)_{j-1} (1 - q^{4j-2})} \\
 &= -\frac{2q(-q)_\infty}{(q)_\infty} f_{1,5,1}(q^3, q^9, q^2) \\
 &= 2m(-q^{22}, q^{48}, q^{-6}) + 2q^{-1} m(-q^{14}, q^{48}, q^{-6}) \\
 &\quad - 2q^3 \frac{J_{8,32} J_{20,48} \bar{J}_{18,48} J_{2,24} \bar{J}_{4,48} J_{12,48} J_{48}^2}{J_{1,2} \bar{J}_{16,48} \bar{J}_{8,48} J_{24}^3 J_{96}} \\
 &\quad - 2q^{-1} \frac{J_{8,32} J_{20,48} \bar{J}_{18,48} J_{2,24} \bar{J}_{20,48} J_{18,48}^2 \bar{J}_{6,48}^2}{J_{1,2} \bar{J}_{16,48} \bar{J}_{8,48} J_{24}^3 J_{48} J_{96}} \\
 &\quad + 2q^{10} \frac{J_{16,32} J_{4,48} \bar{J}_{6,48} J_{10,24} \bar{J}_{4,48} J_{12,48} J_{96}}{J_{1,2} \bar{J}_{16,48} \bar{J}_{8,48} J_{24}^2 J_{48}} \\
 &\quad + 2 \frac{J_{16,32} J_{4,48} \bar{J}_{6,48} J_{10,24} \bar{J}_{20,48} J_{36,96}^2}{J_{1,2} \bar{J}_{16,48} \bar{J}_{8,48} J_{24}^2 J_{96}}, \tag{51}
 \end{aligned}$$

$$\begin{aligned}
 \mathcal{M}_{18}(q) &:= \sum_{n \geq 0} \sum_{n \geq j \geq 0} \frac{(-1)^j q^{n^2+n+\binom{j}{2}}}{(q)_{n-j}(q)_j(1-q^{2j+1})} \\
 &= \frac{1}{(q)_\infty} f_{3,7,3}(q^2, q^5, q) \\
 &= m(-q^{67}, q^{120}, q^{-9}) + q^{-9}m(-q^{13}, q^{120}, q^9) - q^{-2}m(-q^{37}, q^{120}, q^9) \\
 &\quad - q^{-1}m(-q^{43}, q^{120}, q^{-9}) + q^{-7} \frac{J_{12,48}J_{3,40}J_{8,40}\bar{J}_{17,40}J_{6,20}\bar{J}_{12,40}J_{14,40}J_{40}^2}{J_1J_9,120\bar{J}_{18,40}\bar{J}_{6,40}J_{20}^3J_{80}} \\
 &\quad + q^{-4} \frac{J_{12,48}J_{3,40}J_{8,40}\bar{J}_{17,40}J_{6,20}\bar{J}_{8,40}J_{17,40}^2\bar{J}_{3,40}^2}{J_1J_9,120\bar{J}_{18,40}\bar{J}_{6,40}J_{20}^3J_{40}J_{80}} \\
 &\quad - q^{-3} \frac{J_{24,48}J_{3,40}J_{12,40}\bar{J}_{3,40}J_{4,20}\bar{J}_{12,40}J_{14,40}J_{80}}{J_1J_9,120\bar{J}_{18,40}\bar{J}_{6,40}J_{20}^2J_{40}} \\
 &\quad - q^{-7} \frac{J_{24,48}J_{3,40}J_{12,40}\bar{J}_{3,40}J_{4,20}\bar{J}_{8,40}J_{34,80}^2}{J_1J_9,120\bar{J}_{18,40}\bar{J}_{6,40}J_{20}^2J_{80}}, \tag{52}
 \end{aligned}$$

$$\begin{aligned}
 \mathcal{M}_{19}(q) &:= \sum_{n \geq 0} \sum_{n \geq j \geq 0} \frac{(-q)_n(-1)^j q^{\binom{n+1}{2}+\binom{j}{2}}}{(q)_{n-j}(q)_j(1-q^{2j+1})} \\
 &= \frac{(-q)_\infty}{(q)_\infty} f_{1,3,1}(q, q^4, q^2) \\
 &= m(-q^{11}, q^{16}, q^{-3}) + q \frac{J_{4,8}J_{16,32}J_{5,16}J_{6,32}}{J_{1,2}\bar{J}_{8,16}\bar{J}_{2,16}}. \tag{53}
 \end{aligned}$$

It will have been noticed that some of the expressions in Theorems 1.5–1.7 are considerably more involved than others. For instance, equation (52) involves four Appell–Lerch series and four modular forms while equation (53) involves only one of each. This depends on the indefinite theta function $f_{n,n+p,n}(x, y, q)$. In general, the number of Appell–Lerch series grows with n and the number of modular forms grows with p .

The final goal of the paper is to give identities involving some of the double sums in Theorems 1.5–1.7 and “classical” mock theta functions. Namely, we express the double sums $\mathcal{M}_2(q)$, $\mathcal{M}_5(q)$, $\mathcal{M}_9(q)$, and $\mathcal{M}_{16}(q)$ in terms of the mock theta functions

$$T_0(q) := \sum_{n \geq 0} \frac{q^{(n+1)(n+2)}(-q^2; q^2)_n}{(-q; q^2)_{n+1}},$$

$$\omega(q) := \sum_{n \geq 0} \frac{q^{2n(n+1)}}{(q; q^2)_{n+1}^2},$$

$$A(q) := \sum_{n \geq 0} \frac{q^{n+1}(-q^2; q^2)_n}{(q; q^2)_{n+1}},$$

and

$$U_1(q) := \sum_{n \geq 0} \frac{q^{(n+1)^2}(-q; q^2)_n}{(-q^2; q^4)_{n+1}},$$

of “orders” 8, 3, 2, and 8, respectively (see [11]). A similar identity was found in [13], namely

$$\mathcal{W}_2(q) = 2qT_1(q) - qS_1(q)$$

where

$$S_1(q) := \sum_{n \geq 0} \frac{q^{n(n+2)}(-q; q^2)_n}{(-q^2; q^2)_n}$$

and

$$T_1(q) := \sum_{n \geq 0} \frac{q^{n(n+1)}(-q^2; q^2)_n}{(-q; q^2)_{n+1}}$$

are mock theta functions of order 8 [11].

COROLLARY 1.8. *We have the following identities.*

$$\mathcal{M}_2(q) = 4T_0(q) + 2 - 2 \frac{J_8^3 J_{4,8}}{J_{2,8}^2 \bar{J}_{1,8}} + \frac{J_{2,4} J_{8,16} J_{1,8} J_{6,16}}{\bar{J}_{1,4} \bar{J}_{1,8} \bar{J}_{3,8}}, \tag{54}$$

$$\mathcal{M}_5(q) = 1 + q\omega(q), \tag{55}$$

$$\mathcal{M}_9(q) = -A(-q^8) + \frac{J_{32}^3 J_{14,32} \bar{J}_{10,32}}{J_{16,32} J_{2,32} \bar{J}_{6,32} \bar{J}_{8,32}} - q^{-1} \frac{J_{64}^2 J_{28,64}}{J_{32} J_{4,64}} + q^{-1} \frac{J_{8,16} J_{32,64} J_{4,32} J_{24,64}}{\bar{J}_{1,4} \bar{J}_{6,32} \bar{J}_{2,32}}, \tag{56}$$

$$\mathcal{M}_{16}(q) = \frac{1}{2} + q^{-1} U_1(q^8) - q^{-1} \frac{J_{32}^3 \bar{J}_{10,32} J_{14,32}}{\bar{J}_{16,32} J_{6,32} J_{8,32} \bar{J}_{2,32}} + \frac{J_{32}^3 J_{10,32} \bar{J}_{6,32}}{J_{6,32} J_{16,32} \bar{J}_{0,32} \bar{J}_{10,32}} \tag{57}$$

$$+ q^{-1} \frac{J_{8,16} J_{32,64} J_{4,32} J_{24,64}}{\bar{J}_{1,4} \bar{J}_{2,32} \bar{J}_{10,32}}. \tag{58}$$

The paper is organized as follows. In Section 2, we prove Theorems 1.1 and 1.2 and record some corollaries. In Section 3, we establish a change of base lemma and deduce Andrews’ Bailey pairs from those of Bringmann and Kane. In Section 4, we recall important work of Hickerson and Mortenson on mock theta functions [12] and then prove Theorems 1.5–1.7 and Corollary 1.8.

In [14], we consider applications of Theorems 1.1–1.3 to q -hypergeometric double sums related to real quadratic fields, in the spirit of [4].

2. Proofs of Theorems 1.1 and 1.2. *Proof of Theorem 1.1.* First note that the sequence α'_n in (21) and (22) is uniquely defined by $\alpha'_0 = 0$, $\alpha'_1 = -(1 - q^2)$, and

$$\frac{\alpha'_{n+2}}{1 - q^{2n+4}} - \frac{q^{2n} \alpha'_n}{1 - q^{2n}} = -\alpha_{n+1}. \tag{59}$$

Suppose that the β'_n are given by (23). Then the corresponding α'_n satisfy the initial conditions. Moreover, using (2) we have

$$\begin{aligned} \frac{\alpha'_{n+2}}{1 - q^{2n+4}} - \frac{q^{2n}\alpha'_n}{1 - q^{2n}} &= \sum_{j=1}^{n+2} \frac{(q)_{n+j+1}q^{\binom{n-j+2}{2}}(-1)^{n+j}}{(q)_{n-j+2}} \beta'_j - \sum_{j=1}^n \frac{(q)_{n+j-1}q^{\binom{n-j}{2}+2n}(-1)^{n+j}}{(q)_{n-j}} \beta'_j \\ &= \sum_{j=1}^{n+2} \frac{(q)_{n+j-1}(-1)^{n+j}q^{\binom{n-j}{2}+2n}}{(q)_{n-j+2}} \left((1 - q^{n+j})(1 - q^{n+j+1})q^{-2j+1} \right. \\ &\quad \left. - (1 - q^{n-j+2})(1 - q^{n-j+1}) \right) \beta'_j \\ &= - \sum_{j=1}^{n+2} \frac{q^{\binom{n-j}{2}+2n}(-1)^{n+j}(q)_{n+j-1}}{(q)_{n-j+2}} \left((1 - q^{2n+2})(1 - q^{-2j+1}) \right) \beta'_j \\ &= (1 - q^{2n+2}) \sum_{j=0}^{n+1} \frac{q^{\binom{n-j-1}{2}+2n-2j-1}(q)_{n+j}(-1)^{n+j+1}}{(q)_{n-j+1}} (1 - q^{2j+1}) \beta'_{j+1} \\ &= -(1 - q^{2n+2}) \sum_{j=0}^{n+1} \frac{q^{\binom{n+1-j}{2}}(q)_{n+j}(-1)^{n+j+1}}{(q)_{n+1-j}} \beta_j \\ &= -\alpha_{n+1}. \end{aligned}$$

□

Proof of Theorem 1.2. Let $a = q$ and let β'_n be defined as in (25). Then

$$\begin{aligned} \alpha'_n &= \frac{(-1)^n}{1 - q} \sum_{j=0}^n \frac{(q)_{n+j}(-1)^j q^{\binom{n-j}{2}}}{(q)_{n-j}} \beta'_j (1 - q^{n+j+1} + q^{n+j+1}(1 - q^{n-j})) \\ &= \frac{(-1)^n}{1 - q} \left(\sum_{j=0}^n \frac{(q)_{n+j+1}(-1)^j q^{\binom{n-j}{2}}}{(q)_{n-j}} \beta'_j + \sum_{j=0}^{n-1} \frac{(q)_{n+j}(-1)^j q^{\binom{n-j}{2}+n+j+1}}{(q)_{n-j-1}} \beta'_j \right) \\ &= \frac{1}{1 - q} \left(- \sum_{j=1}^{n+1} \frac{(q)_{n+j}(-1)^{n+j+1} q^{\binom{n-j+1}{2}}}{(q)_{n-j+1}} \beta_j + \sum_{j=1}^n \frac{(q)_{n+j-1}(-1)^{n-j} q^{\binom{n-j}{2}+2n}}{(q)_{n-j}} \beta_j \right) \\ &= \frac{1}{1 - q} \left(- \frac{\alpha_{n+1}}{1 - q^{2n+2}} + \frac{q^{2n}\alpha_n}{1 - q^{2n}} \right), \end{aligned}$$

which establishes the result. □

Note that the proof of Theorem 1.1 implies an inverse result. Since the α'_n are uniquely defined by the α_n in (59) together with the initial conditions $\alpha'_0 = 0$ and $\alpha'_1 = -(1 - q^2)$, we have the following.

THEOREM 2.1. *If (α'_n, β'_n) form a Bailey pair relative to 1 with $\alpha'_0 = 0$ and $\alpha'_1 = -(1 - q^2)$, then (α_n, β_n) also form a Bailey pair relative to 1, where $\alpha_0 = \beta_0 = 1$,*

$$\alpha_n = \frac{-1}{1 - q^{2n+2}} \alpha'_{n+1} + \frac{q^{2n-2}}{1 - q^{2n-2}} \alpha'_{n-1}, \tag{60}$$

and

$$\beta_n = -(1 - q^{2n+1})\beta'_{n+1}. \tag{61}$$

We finish this section with three corollaries of Theorems 1.1 and 1.3, giving three sets of two Bailey pairs involving indefinite quadratic forms. These come from three Bailey pairs in Slater’s list [16]. These are not the only three pairs from Slater’s list which lead to indefinite quadratic forms in Theorems 1.1 and 1.3, but we have limited ourselves to those we will use in the sequel.

First, on p. 468 of [16] we find the Bailey pair relative to 1,

$$\alpha_n = \begin{cases} 1, & \text{if } n = 0, \\ (-1)^n(q^n + q^{-n}), & \text{otherwise,} \end{cases}$$

and

$$\beta_n = \frac{(-1)^n q^{-n}}{(q^2; q^2)_n}.$$

Applying Theorems 1.1 and 1.3 we have the following.

COROLLARY 2.2. *The sequences (a_n, b_n) form a Bailey pair relative to 1, where*

$$a_{2n} = (1 - q^{4n})q^{2n^2-2n+1} \sum_{j=-n}^{n-1} q^{-2j^2}, \tag{62}$$

$$a_{2n+1} = -(1 - q^{4n+2})q^{2n^2} \sum_{j=-n}^n q^{-2j^2-2j}, \tag{63}$$

and

$$b_n = \begin{cases} 0, & \text{if } n = 0, \\ \frac{(-1)^n q^{-n+1}}{(q^2; q^2)_{n-1}(1 - q^{2n-1})}, & \text{otherwise,} \end{cases} \tag{64}$$

and the sequences (α_n, β_n) form a Bailey pair relative to q , where

$$\alpha_{2n} = \frac{1}{1 - q} \left(q^{2n^2} \sum_{j=-n}^n q^{-2j^2-2j} + q^{2n^2+2n+1} \sum_{j=-n}^{n-1} q^{-2j^2} \right), \tag{65}$$

$$\alpha_{2n+1} = -\frac{1}{1 - q} \left(q^{2n^2+2n+1} \sum_{j=-n-1}^n q^{-2j^2} + q^{2n^2+4n+2} \sum_{j=-n}^n q^{-2j^2-2j} \right), \tag{66}$$

and

$$\beta_n = \frac{(-1)^n q^{-n}}{(q^2; q^2)_n(1 - q^{2n+1})}. \tag{67}$$

Next, on p. 468 of [16] we find the Bailey pair relative to 1,

$$\alpha_n = \begin{cases} 1, & \text{if } n = 0, \\ (-1)^n q^{-\binom{n+1}{2}}(1 + q^n), & \text{otherwise,} \end{cases}$$

and

$$\beta_n = \frac{(-1)^n q^{-\binom{n+1}{2}}}{(q)_n}.$$

Applying Theorems 1.1 and 1.3, we have the following.

COROLLARY 2.3. *The sequences (a_n, b_n) form a Bailey pair relative to 1, where*

$$a_{2n} = (1 - q^{4n})q^{2n^2-2n} \sum_{j=-n}^{n-1} q^{-4j^2-3j}, \tag{68}$$

$$a_{2n+1} = -(1 - q^{4n+2})q^{2n^2} \sum_{j=-n}^n q^{-4j^2-j}, \tag{69}$$

and

$$b_n = \begin{cases} 0, & \text{if } n = 0, \\ \frac{(-1)^n q^{-\binom{n}{2}}}{(q)_{n-1}(1 - q^{2n-1})}, & \text{otherwise,} \end{cases} \tag{70}$$

and the sequences (α_n, β_n) form a Bailey pair relative to q , where

$$\alpha_{2n} = \frac{1}{1 - q} \left(q^{2n^2} \sum_{j=-n}^n q^{-4j^2-j} + q^{2n^2+2n} \sum_{j=-n}^{n-1} q^{-4j^2-3j} \right), \tag{71}$$

$$\alpha_{2n+1} = -\frac{1}{1 - q} \left(q^{2n^2+2n} \sum_{j=-n-1}^n q^{-4j^2-3j} + q^{2n^2+4n+2} \sum_{j=-n}^n q^{-4j^2-j} \right), \tag{72}$$

and

$$\beta_n = \frac{(-1)^n q^{-\binom{n+1}{2}}}{(q)_n(1 - q^{2n+1})}. \tag{73}$$

Finally, on p. 468 of [16] we find the Bailey pair relative to 1,

$$\alpha_n = \begin{cases} 1, & \text{if } n = 0, \\ (-1)^n q^{-n(n+3)/2}(1 + q^{2n}), & \text{otherwise,} \end{cases}$$

and

$$\beta_n = \frac{(-1)^n q^{-n(n+3)/2}}{(q)_n}.$$

Applying Theorems 1.1 and 1.3, we find the following.

COROLLARY 2.4. *The sequences (a_n, b_n) form a Bailey pair relative to 1, where*

$$a_{2n} = (1 - q^{4n})q^{2n^2-2n+1} \sum_{j=-n}^{n-1} q^{-4j^2-j}, \tag{74}$$

$$a_{2n+1} = -(1 - q^{4n+2})q^{2n^2} \sum_{j=-n}^n q^{-4j^2-3j}, \tag{75}$$

and

$$b_n = \begin{cases} 0, & \text{if } n = 0, \\ \frac{(-1)^n q^{-\binom{n+1}{2}}}{(q)_{n-1}(1-q^{2n-1})}, & \text{otherwise,} \end{cases} \tag{76}$$

and the sequences (α_n, β_n) form a Bailey pair relative to q , where

$$\alpha_{2n} = \frac{1}{1-q} \left(q^{2n^2} \sum_{j=-n}^n q^{-4j^2-3j} + q^{2n^2+2n+1} \sum_{j=-n}^{n-1} q^{-4j^2-j} \right), \tag{77}$$

$$\alpha_{2n+1} = -\frac{1}{1-q} \left(q^{2n^2+2n+1} \sum_{j=-n-1}^n q^{-4j^2-j} + q^{2n^2+4n+2} \sum_{j=-n}^n q^{-4j^2-3j} \right), \tag{78}$$

and

$$\beta_n = \frac{(-1)^n q^{-n(n+3)/2}}{(q)_n(1-q^{2n+1})}. \tag{79}$$

3. The seventh order Bailey pairs of Andrews. We begin with a change of base lemma for Bailey pairs. For other results of this nature, see [7] and [8]. Throughout this section, we emphasize the base by saying that a pair of sequences satisfying (1) is a Bailey pair relative to (a, q) .

LEMMA 3.1. *If (α_n, β_n) is a Bailey pair relative to $(1, q)$, then (α'_n, β'_n) is a Bailey pair relative to $(1, q^2)$, where*

$$\alpha'_n = \frac{1}{2}(1 + q^{2n})q^{n^2-n}\alpha_n \tag{80}$$

and

$$\beta'_n = \frac{1}{(-1)_{2n}} \sum_{k=0}^n \frac{q^{k^2-k}}{(q^2; q^2)_{n-k}} \beta_k. \tag{81}$$

Proof. We will need the fact that

$$(q^{-n})_k = \frac{(q)_n}{(q)_{n-k}} (-1)^k q^{\binom{k}{2} - nk} \tag{82}$$

along with the identity

$$\sum_{k=0}^{n-r} \frac{(-1)^k q^{2nk} (q^{-(n-r)})_k (-q^{-(n-r)})_k}{(q)_k (q^{2r+1})_k} = \frac{(1 + q^{2r})(q)_{2r} (-1)_{2n}}{2(q^2; q^2)_{n+r}}, \tag{83}$$

which follows from a short calculation using the case $z = -q^{2n}$, $a = q^{-n+r}$, $b = -q^{-n+r}$, and $c = q^{2r+1}$ of the second Heine transformation [10],

$$\sum_{n \geq 0} \frac{(a)_n (b)_n}{(c)_n (q)_n} z^n = \frac{(c/b)_\infty (bz)_\infty}{(c)_\infty (z)_\infty} \sum_{n \geq 0} \frac{(\frac{abz}{c})_n (b)_n}{(bz)_n (q)_n} \left(\frac{c}{b}\right)^n. \tag{84}$$

Now, beginning with (81), we have

$$\begin{aligned} \beta'_n &= \frac{1}{(-1)_{2n}} \sum_{k=0}^n \frac{q^{k^2-k}}{(q^2; q^2)_{n-k}} \beta_k \\ &= \frac{1}{(-1)_{2n}} \sum_{k=0}^n \frac{q^{k^2-k}}{(q^2; q^2)_{n-k}} \sum_{r=0}^k \frac{1}{(q)_{k-r} (q)_{k+r}} \alpha_r \\ &= \frac{1}{(-1)_{2n}} \sum_{r=0}^n \alpha_r \sum_{k=r}^n \frac{q^{k^2-k}}{(q^2; q^2)_{n-k}} \frac{1}{(q)_{k-r} (q)_{k+r}} \\ &= \frac{1}{(-1)_{2n}} \sum_{r=0}^n \alpha_r \sum_{k=0}^{n-r} \frac{q^{k^2+2kr+r^2-k-r}}{(q^2; q^2)_{n-k-r} (q)_k (q)_{k+2r}} \\ &= \frac{1}{(-1)_{2n}} \sum_{r=0}^n \frac{q^{r^2-r}}{(q)_{2r}} \alpha_r \sum_{k=0}^{n-r} \frac{q^{k^2+2kr-k}}{(q^2; q^2)_{n-k-r} (q)_k (q^{2r+1})_k} \\ &= \frac{1}{(-1)_{2n}} \sum_{r=0}^n \frac{q^{r^2-r}}{(q)_{2r} (q^2; q^2)_{n-r}} \alpha_r \sum_{k=0}^{n-r} \frac{(-1)^k q^{2nk} (q^{-2(n-r)}; q^2)_k}{(q)_k (q^{2r+1})_k} \tag{by (82)} \\ &= \frac{1}{(-1)_{2n}} \sum_{r=0}^n \frac{q^{r^2-r}}{(q)_{2r} (q^2; q^2)_{n-r}} \alpha_r \sum_{k=0}^{n-r} \frac{(-1)^k q^{2nk} (q^{-(n-r)})_k (-q^{-(n-r)})_k}{(q)_k (q^{2r+1})_k} \\ &= \sum_{r=0}^n \frac{(1 + q^{2r}) q^{r^2-r}}{2(q^2; q^2)_{n-r} (q^2; q^2)_{n+r}} \alpha_r \tag{by (83)}. \end{aligned}$$

This implies the statement of the theorem. □

Now, we insert the Bailey pair in (6)–(8) into Lemma 3.1. We obtain a Bailey pair relative to $(1, q^2)$, where

$$a'_{2n} = \frac{1}{2} (1 - q^{8n}) q^{6n^2-4n} \sum_{j=-n}^{n-1} q^{-2j^2-2j},$$

$$a'_{2n+1} = -\frac{1}{2}(1 - q^{8n+4})q^{6n^2+2n} \sum_{j=-n}^n q^{-2j^2},$$

$b'_0 = 0$, and for $n \geq 1$,

$$\begin{aligned} b'_n &= \frac{1}{(-1)_{2n}} \sum_{k=1}^n \frac{(-1)^k q^{k^2-k} (q; q^2)_{k-1}}{(q)_{2k-1} (q^2; q^2)_{n-k}} \\ &= \frac{-1}{(-1)_{2n}} \sum_{k=0}^{n-1} \frac{(-1)^k q^{k^2+k} (q; q^2)_k}{(q)_{2k+1} (q^2; q^2)_{n-k-1}} \\ &= \frac{-1}{(-1)_{2n} (q^2; q^2)_{n-1} (1-q)} \sum_{k=0}^{n-1} \frac{q^{2nk} (q^{-2(n-1)}; q^2)_k (q; q^2)_k}{(q^2; q^2)_k (q^3; q^2)_k} \quad (\text{by (82)}) \\ &= \frac{-1}{(-1)_{2n} (q; q^2)_n} \\ &= \frac{-1}{2(q^{2n}; q^2)_n}, \end{aligned}$$

where the penultimate equality follows from the case $q = q^2$, $n = n - 1$, $a = q$, and $c = q^3$ of the q -Chu–Vandermonde summation [10]

$$\sum_{k=0}^n \frac{(a)_k (q^{-n})_k}{(q)_k (c)_k} \left(\frac{cq^n}{a}\right)^k = \frac{(c/a)_n}{(c)_n}.$$

Now multiplying (a'_n, b'_n) by -2 and replacing q by $q^{1/2}$ gives Andrews’ Bailey pair (15)–(17). Finally, multiplying (15)–(17) by -1 , then applying Theorem 1.2 and Theorem 2.1 gives the pairs (18)–(20) and (12)–(14), respectively.

4. Proofs of Theorems 1.5–1.7 and Corollary 1.8. The approach for proving Theorems 1.5–1.7 is as follows. We first apply (3) and (4) to Corollaries 2.2–2.4 to obtain new Bailey pairs, then use (5) in various ways to obtain identities expressing q -hypergeometric double sums in terms of the indefinite theta series (29). Next, to deduce that these q -hypergeometric double sums are mock theta functions, we apply the following three explicit results of Hickerson and Mortenson which express (29) in terms of the Appell–Lerch series (30). Define

$$\begin{aligned} &g_{a,b,c}(x, y, q, z_1, z_0) \\ &:= \sum_{t=0}^{a-1} (-y)^t q^{c\binom{t}{2}} j(q^{bt}x, q^a) m \left(-q^{a\binom{b+1}{2} - c\binom{a+1}{2} - t(b^2-ac)} \frac{(-y)^a}{(-x)^b}, q^{a(b^2-ac)}, z_0 \right) \\ &+ \sum_{t=0}^{c-1} (-x)^t q^{a\binom{t}{2}} j(q^{bt}y, q^c) m \left(-q^{c\binom{b+1}{2} - a\binom{c+1}{2} - t(b^2-ac)} \frac{(-x)^c}{(-y)^b}, q^{c(b^2-ac)}, z_1 \right). \end{aligned} \tag{85}$$

Following [12], we use the term “generic” to mean that the parameters do not cause poles in the Appell–Lerch sums or in the quotients of theta functions.

THEOREM 4.1. [12, Theorem 1.6] *Let n be a positive integer. For generic $x, y \in \mathbb{C}^*$*

$$f_{n,n+1,n}(x, y, q) = g_{n,n+1,n}(x, y, q, y^n/x^n, x^n/y^n).$$

THEOREM 4.2. [12, Theorem 1.9] *Let n be an odd positive integer. For generic $x, y \in \mathbb{C}^*$*

$$f_{n,n+2,n}(x, y, q) = g_{n,n+2,n}(x, y, q, y^n/x^n, x^n/y^n) - \Theta_{n,2}(x, y, q)$$

where

$$\Theta_{n,2}(x, y, q) := \frac{y^{(n+1)/2} J_{2n,4n} J_{4(n+1),8(n+1)} j(y/x, q^{4(n+1)}) j(q^{n+2}xy, q^{4(n+1)}) j(q^{2n}/x^2y^2, q^{8(n+1)})}{q^{(n^2-3)/2} x^{(n-3)/2} j(y^n/x^n, q^{4n(n+1)}) j(-q^{n+2}x^2, q^{4(n+1)}) j(-q^{n+2}y^2, q^{4(n+1)})}.$$

THEOREM 4.3. [12, Theorem 1.11] *Let n be an odd positive integer. For generic $x, y \in \mathbb{C}^*$*

$$f_{n,n+4,n}(x, y, q) = g_{n,n+4,n}(x, y, q, y^n/x^n, x^n/y^n) - \Theta_{n,4}(x, y, q)$$

where

$$\Theta_{n,4}(x, y, q) := \frac{q^{-(n^2+n-3)} x^{-(n-3)/2} y^{(n+1)/2} j(y/x, q^{4(2n+4)})}{j(y^n/x^n, q^{4n(2n+4)}) j(-q^{2n+8}x^4, q^{4(2n+4)}) j(-q^{2n+8}, q^{4(2n+4)})} \left\{ J_{4n,16n} S_1 - q J_{8n,16n} S_2 \right\},$$

$$S_1 := \frac{j(q^{6n+16}x^2y^2, q^{4(2n+4)}) j(-q^{2(2n+4)}y/x, q^{4(2n+4)}) j(q^{n+4}xy, q^{2(2n+4)})}{J_{2(2n+4)}^3 J_{8(2n+4)}} \cdot \left\{ j(-q^{2n+8}x^2y^2, q^{4(2n+4)}) j(q^{2(2n+4)}, q^{4(2n+4)}) J_{4(2n+4)}^2 \right.$$

$$\left. + \frac{q^{n+4}x^2 j(-q^{6n+16}x^2y^2, q^{4(2n+4)}) j(q^{2(2n+4)}y/x, q^{4(2n+4)})^2 j(-y/x, q^{4(2n+4)})^2}{J_{4(2n+4)}} \right\}$$

and

$$S_2 := \frac{j(q^{2n+8}x^2y^2, q^{4(2n+4)}) j(-y/x, q^{4(2n+4)}) j(q^{3n+8}xy, q^{2(2n+4)})}{J_{2(2n+4)}^2} \cdot \left\{ \frac{q^{n+1} j(-q^{2n+8}x^2y^2, q^{4(2n+4)}) j(q^{2(2n+4)}y^2/x^2, q^{4(2n+4)}) J_{8(2n+4)}}{y J_{4(2n+4)}} \right.$$

$$\left. + \frac{qxy j(-q^{6n+16}x^2y^2, q^{4(2n+4)}) j(q^{4(2n+4)}y^2/x^2, q^{8(2n+4)})^2}{J_{8(2n+4)}} \right\}.$$

Finally, we use the fact that specializations of Appell–Lerch series are well known to be mock theta functions [18], [19, Ch. 1].

To simplify expressions arising in Theorems 4.1–4.3, we require certain facts about $j(x, q)$ and $m(x, q, z)$. From the definition of $j(x, q)$, we have

$$j(q^n x, q) = (-1)^n q^{-\binom{n}{2}} x^{-n} j(x, q) \tag{86}$$

where $n \in \mathbb{Z}$ and

$$j(x, q) = j(q/x, q) = -xj(x^{-1}, q). \tag{87}$$

Next, some relevant properties of the sum $m(x, q, z)$ are given in the following (see (3.2b), (3.2c) of Proposition 3.1, (3.3) of Corollary 3.2 and Theorem 3.3 in [12]).

PROPOSITION 4.4. For generic $x, z, z_0, z' \in \mathbb{C}^*$,

$$m(x, q, z) = x^{-1} m(x^{-1}, q, z^{-1}), \tag{88}$$

$$m(qx, q, z) = 1 - xm(x, q, z), \tag{89}$$

$$m(q, q^2, -1) = \frac{1}{2} \tag{90}$$

and

$$m(x, q, z) = m(x, q, z_0) + \frac{z_0 J_1^3 j(z/z_0, q) j(xz z_0, q)}{j(z_0, q) j(z, q) j(xz z_0, q) j(xz, q)}. \tag{91}$$

We are now ready to proceed to the proofs of Theorems 1.5–1.7.

Proof of Theorem 1.5. Applying equations (3) and (4) with $(a, \rho_1, \rho_2) = (1, -1, \infty)$ to (62)–(64) and $(q, -q, \infty)$ to (65)–(67) gives a Bailey pair relative to 1,

$$a'_{2n} = 2(1 - q^{2n}) q^{4n^2 - n + 1} \sum_{j=-n}^{n-1} q^{-2j^2}, \tag{92}$$

$$a'_{2n+1} = -2(1 - q^{2n+1}) q^{4n^2 + 3n + 1} \sum_{j=-n}^n q^{-2j^2 - 2j}, \tag{93}$$

and

$$b'_n = \frac{1}{(-q)_n} \sum_{j=1}^n \frac{(-1)_j (-1)^j q^{\binom{j}{2} + 1}}{(q)_{n-j} (q^2; q^2)_{j-1} (1 - q^{2j-1})}, \tag{94}$$

and a Bailey pair relative to q ,

$$\alpha'_{2n} = \frac{1}{1 - q} \left(q^{4n^2 + n} \sum_{j=-n}^n q^{-2j^2 - 2j} + q^{4n^2 + 3n + 1} \sum_{j=-n}^{n-1} q^{-2j^2} \right), \tag{95}$$

$$\alpha'_{2n+1} = -\frac{1}{1 - q} \left(q^{4n^2 + 5n + 2} \sum_{j=-n-1}^n q^{-2j^2} + q^{4n^2 + 7n + 3} \sum_{j=-n}^n q^{-2j^2 - 2j} \right), \tag{96}$$

and

$$\beta'_n = \frac{1}{(-q)_n} \sum_{j=0}^n \frac{(-1)^j q^{\binom{j}{2}}}{(q)_{n-j}(q)_j(1 - q^{2j+1})}, \tag{97}$$

respectively. Now to prove (35), we insert the Bailey pair (a'_n, b'_n) from equations (92)–(94) into (5) with $\rho_1, \rho_2 \rightarrow \infty$. This gives

$$\begin{aligned} q\mathcal{M}_1(q) &= \sum_{n \geq 0} q^{n^2} b'_n(q) = \frac{1}{(q)_\infty} \sum_{n \geq 0} q^{n^2} a'_n(q) \\ &= \frac{1}{(q)_\infty} \left(\sum_{n \geq 0} q^{4n^2} a'_{2n}(q) + \sum_{n \geq 0} q^{4n^2+4n+1} a'_{2n+1}(q) \right) \\ &= \frac{2}{(q)_\infty} \left(\sum_{n \geq 0} q^{8n^2-n+1} \sum_{j=-n}^{n-1} q^{-2j^2} - \sum_{n \geq 0} q^{8n^2+n+1} \sum_{j=-n}^{n-1} q^{-2j^2} \right. \\ &\quad \left. - \sum_{n \geq 0} q^{8n^2+7n+2} \sum_{j=-n}^n q^{-2j^2-2j} + \sum_{n \geq 0} q^{8n^2+9n+3} \sum_{j=-n}^n q^{-2j^2-2j} \right). \end{aligned}$$

After replacing n with $-n$ in the second sum and n with $-n - 1$ in the fourth sum, we let $n = (r + s + 1)/2, j = (r - s - 1)/2$ in the first two sums and $n = (r + s)/2, j = (r - s)/2$ in the latter two sums to find

$$\begin{aligned} \mathcal{M}_1(q) &= \frac{2q}{(q)_\infty} \left(\left(\sum_{\substack{r,s \geq 0 \\ r \not\equiv s \pmod{2}}} - \sum_{\substack{r,s < 0 \\ r \not\equiv s \pmod{2}}} \right) q^{\frac{3}{2}r^2+5rs+\frac{9}{2}r+\frac{3}{2}s^2+\frac{5}{2}s} \right. \\ &\quad \left. - \left(\sum_{\substack{r,s \geq 0 \\ r \equiv s \pmod{2}}} - \sum_{\substack{r,s < 0 \\ r \equiv s \pmod{2}}} \right) q^{\frac{3}{2}r^2+5rs+\frac{5}{2}r+\frac{3}{2}s^2+\frac{9}{2}s} \right) \tag{98} \\ &= -\frac{2q}{(q)_\infty} \left(\left(\sum_{r,s \geq 0} - \sum_{r,s < 0} \right) q^{\frac{3}{2}r^2+5rs+\frac{9}{2}r+\frac{3}{2}s^2+\frac{5}{2}s} \right) \\ &= -\frac{2q}{(q)_\infty} f_{3,5,3}(q^6, q^4, q). \end{aligned}$$

By (98), Theorem 4.2 with $n = 3$ and (85)–(87),

$$\begin{aligned} \mathcal{M}_1(q) &= -2q^{-10}m(-q^{-7}, q^{48}, q^6) + 2q^{-23}m(-q^{-23}, q^{48}, q^6) + 2m(-q^{25}, q^{48}, q^{-6}) \\ &\quad - 2q^{-10}m(-q^{-7}, q^{48}, q^{-6}) + \frac{2q}{(q)_\infty} \Theta_{3,2}(q^6, q^4, q). \tag{99} \end{aligned}$$

Now, we simplify (99) using (88), (89) and (91) to obtain

$$\begin{aligned} \mathcal{M}_1(q) &= 4q^{-3}m(-q^7, q^{48}, q^6) + 4m(-q^{25}, q^{48}, q^6) - 2 + \frac{2q}{(q)_\infty} \Theta_{3,2}(q^6, q^4, q) \\ &\quad + 2q^{-3} \frac{J_{48}^3 J_{12,48} \bar{J}_{7,48}}{J_{6,48}^2 \bar{J}_{1,48} \bar{J}_{13,48}} + 2 \frac{J_{48}^3 J_{12,48} \bar{J}_{25,48}}{J_{6,48}^2 \bar{J}_{19,48} \bar{J}_{31,48}} \\ &= 4q^{-3}m(-q^7, q^{48}, q^{24}) + 4m(-q^{25}, q^{48}, q^{-24}) - 4q^3 \frac{J_{48}^3 J_{18,48} \bar{J}_{11,48}}{J_{6,48} J_{24,48} \bar{J}_{13,48} \bar{J}_{17,48}} \\ &\quad - 4 \frac{J_{48}^3 J_{18,48} \bar{J}_{7,48}}{J_{6,48} J_{24,48} \bar{J}_{1,48} \bar{J}_{17,48}} - 2 + \frac{2q}{(q)_\infty} \Theta_{3,2}(q^6, q^4, q) \\ &\quad + 2q^{-3} \frac{J_{48}^3 J_{12,48} \bar{J}_{7,48}}{J_{6,48}^2 \bar{J}_{1,48} \bar{J}_{13,48}} + 2 \frac{J_{48}^3 J_{12,48} \bar{J}_{25,48}}{J_{6,48}^2 \bar{J}_{19,48} \bar{J}_{31,48}}. \end{aligned}$$

Comparing with (35), we are left with a modular identity to verify. Such a verification can always be done using a finite computation. This, and similar computations in this paper, were carried out using computer software packages available at <http://www.qseries.org/fgarvan>.

This proves identity (35) and shows that $\mathcal{M}_1(q)$ is a mock theta function.

As equations (36)–(39) are handled similarly, we briefly sketch the relevant details. For equations (36) and (37), we again use the Bailey pair (92)–(94) in (5), with $(\rho_1, \rho_2, q) = (\sqrt{q}, -\sqrt{q}, q)$ and (q, ∞, q^2) and the above argument to obtain

$$\mathcal{M}_2(q) = \frac{(q; q^2)_\infty}{(q^2; q^2)_\infty} f_{1,3,1}(-q, -q^3, q)$$

and

$$\mathcal{M}_3(q) = \frac{2q(q; q^2)_\infty}{(q^2; q^2)_\infty} f_{1,2,1}(-q^5, -q^9, q^4).$$

One then proceeds with Theorems 4.2 and 4.1, respectively, and simplifies.

For (38) and (39), we use the Bailey pair (95)–(97) in (5), with $(\rho_1, \rho_2, q) = (\infty, \infty, q)$ and $(-q, \infty, q)$ to get

$$\mathcal{M}_4(q) = \frac{1}{(q)_\infty} f_{3,5,3}(q^2, q^4, q)$$

and

$$\mathcal{M}_5(q) = \frac{(-q)_\infty}{(q)_\infty} f_{1,2,1}(q, q^3, q^2).$$

Applying Theorems 4.2 and 4.1, respectively, and continuing as above yields the result. □

Proof of Theorem 1.6. Applying equations (3) and (4) with $(a, \rho_1, \rho_2) = (1, \infty, \infty)$ to (68)–(70) and (q, ∞, ∞) to (71)–(73) gives a Bailey pair relative to 1,

$$a'_{2n} = (1 - q^{4n})q^{6n^2-2n} \sum_{j=-n}^{n-1} q^{-4j^2-3j}, \tag{100}$$

$$a'_{2n+1} = -(1 - q^{4n+2})q^{6n^2+4n+1} \sum_{j=-n}^n q^{-4j^2-j}, \tag{101}$$

and

$$b'_n = \sum_{j=1}^n \frac{(-1)^j q^{\binom{j+1}{2}}}{(q)_{n-j}(q)_{j-1}(1 - q^{2j-1})}, \tag{102}$$

and a Bailey pair relative to q ,

$$\alpha'_{2n} = \frac{1}{1 - q} \left(q^{6n^2+2n} \sum_{j=-n}^n q^{-4j^2-j} + q^{6n^2+4n} \sum_{j=-n}^{n-1} q^{-4j^2-3j} \right), \tag{103}$$

$$\alpha'_{2n+1} = -\frac{1}{1 - q} \left(q^{6n^2+8n+2} \sum_{j=-n-1}^n q^{-4j^2-3j} + q^{6n^2+10n+4} \sum_{j=-n}^n q^{-4j^2-j} \right), \tag{104}$$

and

$$\beta'_n = \sum_{j=0}^n \frac{(-1)^j q^{\binom{j+1}{2}}}{(q)_{n-j}(q)_j(1 - q^{2j+1})}, \tag{105}$$

respectively. For the identities (40)–(44) in Theorem 1.6, we use the Bailey pair (100)–(102) in (5) with $(\rho_1, \rho_2, q) = (\infty, \infty, q)$, $(-1, \infty, q)$, $(\sqrt{q}, -\sqrt{q}, q)$, (q, ∞, q^2) , and $(-1, -q, q^2)$ to obtain

$$\mathcal{M}_6(q) = -\frac{q^2}{(q)_\infty} f_{3,7,3}(q^5, q^6, q),$$

$$\mathcal{M}_7(q) = -\frac{2q^2(-q)_\infty}{(q)_\infty} f_{1,3,1}(q^4, q^5, q^2),$$

$$\mathcal{M}_8(q) = \frac{q(q; q^2)_\infty}{2(q^2; q^2)_\infty} f_{1,5,1}(-q^2, -q^3, q),$$

$$\mathcal{M}_9(q) = \frac{q^3(q; q^2)_\infty}{(q^2; q^2)_\infty} f_{1,3,1}(-q^7, -q^9, q^4),$$

$$\mathcal{M}_{10}(q) = -\frac{2q^3(-q)_\infty}{(q)_\infty} f_{1.5,1}(q^5, q^7, q^2),$$

respectively. One then applies Theorems 4.2 and 4.3 and proceeds as in the proof of (35). The identities (45) and (46) follow similarly but with the Bailey pair (103)–(105) in (5) with $(\rho_1, \rho_2, q) = (\infty, \infty, q)$ and $(-q, \infty, q)$ yielding

$$\mathcal{M}_{11}(q) = \frac{1}{(q)_\infty} f_{3,7,3}(q^3, q^4, q)$$

and

$$\mathcal{M}_{12}(q) = \frac{(-q)_\infty}{(q)_\infty} f_{1,3,1}(q^2, q^3, q^2).$$

One now applies Theorems 4.3 and 4.2, respectively, and simplifies. □

Proof of Theorem 1.7. Applying (3) and (4) with $(a, \rho_1, \rho_2) = (1, \infty, \infty)$ to (74)–(76) and (q, ∞, ∞) to (77)–(79), we obtain a Bailey pair relative to 1,

$$a'_{2n} = (1 - q^{4n})q^{6n^2-2n+1} \sum_{j=-n}^{n-1} q^{-4j^2-j}, \tag{106}$$

$$a'_{2n+1} = -(1 - q^{4n+2})q^{6n^2+4n+1} \sum_{j=-n}^n q^{-4j^2-3j}, \tag{107}$$

and

$$b'_n = \sum_{j=1}^n \frac{(-1)^j q^{\binom{j}{2}+1}}{(q)_{n-j}(q)_{j-1}(1 - q^{2j-1})}, \tag{108}$$

and a Bailey pair relative to q

$$\alpha'_{2n} = \frac{1}{1 - q} \left(q^{6n^2+2n} \sum_{j=-n}^n q^{-4j^2-3j} + q^{6n^2+4n+1} \sum_{j=-n}^{n-1} q^{-4j^2-j} \right), \tag{109}$$

$$\alpha'_{2n+1} = -\frac{1}{1 - q} \left(q^{6n^2+8n+3} \sum_{j=-n-1}^n q^{-4j^2-j} + q^{6n^2+10n+4} \sum_{j=-n}^n q^{-4j^2-3j} \right), \tag{110}$$

and

$$\beta'_n = \sum_{j=0}^n \frac{(-1)^j q^{\binom{j}{2}}}{(q)_{n-j}(q)_j(1 - q^{2j+1})}, \tag{111}$$

respectively. For the identities (47)–(51) in Theorem 1.7, we use the Bailey pair (106)–(108) in (5) with $(\rho_1, \rho_2, q) = (\infty, \infty, q), (-1, \infty, q), (\sqrt{q}, -\sqrt{q}, q), (q, \infty, q^2),$ and $(-1, -q, q^2)$ to obtain

$$\mathcal{M}_{13}(q) = -\frac{q}{(q)_\infty} f_{3,7,3}(q^4, q^7, q),$$

$$\mathcal{M}_{14}(q) = -\frac{2q(-q)_\infty}{(q)_\infty} f_{1,3,1}(q^3, q^6, q^2),$$

$$\mathcal{M}_{15}(q) = \frac{(q; q^2)_\infty}{2(q^2; q^2)_\infty} f_{1,5,1}(-q, -q^4, q),$$

$$\mathcal{M}_{16}(q) = \frac{q(q; q^2)_\infty}{(q^2; q^2)_\infty} f_{1,3,1}(-q^5, -q^{11}, q^4),$$

$$\mathcal{M}_{17}(q) = -\frac{2q(-q)_\infty}{(q)_\infty} f_{1,5,1}(q^3, q^9, q^2),$$

respectively. One then applies Theorems 4.2 and 4.3 and proceeds as above where (90) is used for $\mathcal{M}_{16}(q)$. The identities (52) and (53) follow similarly but with the Bailey pair (109)–(111) in (5) with $(\rho_1, \rho_2, q) = (\infty, \infty, q)$ and $(-q, \infty, q)$ yielding

$$\mathcal{M}_{18}(q) = \frac{1}{(q)_\infty} f_{3,7,3}(q^2, q^5, q)$$

and

$$\mathcal{M}_{19}(q) = \frac{(-q)_\infty}{(q)_\infty} f_{1,3,1}(q, q^4, q^2).$$

One applies Theorems 4.3 and 4.2, respectively, and simplifies. □

Proof of Corollary 1.8. The identities in Corollary 1.8 are established by comparing the expressions in Theorems 1.5–1.7 with those for classical mock theta functions. We sketch the details. Equations (5.37) and (5.8) in [12] state

$$T_0(q) = -m(-q^3, q^8, q^2)$$

and

$$\omega(q) = -2q^{-1}m(q, q^6, q^2) + \frac{J_6^3}{J_2 J_{3,6}}.$$

By (88), (89), (91), (36) and (39), (55) and (56) follow. Equations (5.1) and (5.40) in [12] state

$$A(q) = -m(q, q^4, q^2)$$

and

$$U_1(q) = -m(-q, q^4, -q^2).$$

By (91), (43) and (50), (57) and (58) follow. \square

ACKNOWLEDGEMENTS. The authors thank Eric Mortenson for helping with the simplification of certain expressions in Theorems 1.5–1.7. They also thank Dennis Stanton for information concerning Section 3. Robert Osburn would like to thank the Institut des Hautes Études Scientifiques for their support during the completion of this paper. This material is based upon work supported by the National Science Foundation under Grant No. 1002477.

REFERENCES

1. G.E. Andrews, Multiple series Rogers-Ramanujan identities, *Pacific J. Math.* **114** (1984), 267–283.
2. G.E. Andrews, q -Series: Their Development and Application in Analysis, Number Theory, Combinatorics, Physics, and Computer Algebra, in *Regional Conference Series in Mathematics*, vol. 66 (American Mathematical Society, Providence, RI, 1986).
3. G.E. Andrews, The fifth and seventh order mock theta functions, *Trans. Amer. Math. Soc.* **293** (1986), 113–134.
4. G.E. Andrews, F. Dyson and D. Hickerson, Partitions and indefinite quadratic forms, *Invent. Math.* **91** (1988), no. 3, 391–407.
5. G.E. Andrews and D. Hickerson, Ramanujan’s “lost” notebook. VII. The sixth order mock theta functions, *Adv. Math.* **89** (1991), no. 1, 60–105.
6. G.E. Andrews, Bailey pairs with free parameters, mock theta functions and tubular partitions, *Ann. Combin.* **18** (2014), 563–578.
7. A. Berkovich and S.O. Warnaar, Positivity preserving transformations for q -binomial coefficients, *Trans. Amer. Math. Soc.* **357** (2005), 2291–2351.
8. D. Bressoud, M.E.H. Ismail and D. Stanton, Change of base in Bailey pairs, *Ramanujan J.* **4** (2000), 435–453.
9. K. Bringmann and B. Kane, Multiplicative q -hypergeometric series arising from real quadratic fields, *Trans. Amer. Math. Soc.* **363**(4) (2011), 2191–2209.
10. G. Gasper and M. Rahman, *Basic Hypergeometric Series*, 2nd edition. Encyclopedia of Mathematics and its Applications, **96** (Cambridge University Press, Cambridge, 2004).
11. B. Gordon and R.J. McIntosh, A survey of classical mock theta functions, in *Partitions, q -series, and Modular Forms*, Developments in Mathematics, vol. 23 (Springer, New York, 2012), 95–144.
12. D. Hickerson and E. Mortenson, Hecke-type double sums, Appell-Lerch sums, and mock theta functions, I, *Proc. London Math. Soc.* **109** (2014), 382–422.
13. J. Lovejoy and R. Osburn, q -hypergeometric double sums as mock theta functions, *Pacific J. Math.* **264** (2013), no. 1, 151–162.
14. J. Lovejoy and R. Osburn, Real quadratic double sums, *Indag. Math.* **26**(4) (2015), 697–712.
15. K. Ono, Unearthing the visions of a master: harmonic Maass forms and number theory, in *Proceedings of the 2008 Harvard-MIT Current Developments in Mathematics Conference* (International Press, Somerville, MA, 2009), 347–454.
16. L.J. Slater, A new proof of Rogers’s transformations of infinite series, *Proc. London Math. Soc.* **53**(2) (1951), 460–475.
17. S. O. Warnaar, Algebraic combinatorics and applications, in *50 years of Bailey’s lemma*, (Göbweinstein, 1999) (Springer, Berlin, 2001), 333–347.
18. D. Zagier, Ramanujan’s mock theta functions and their applications (after Zwegers and Ono-Bringmann), *Astérisque* **326** (2009), 143–164.
19. S. Zwegers, Mock Theta Functions, PhD Thesis (Universiteit Utrecht, 2002).