

the USA 4%-6%. Existing studies have reported that people are switching to vaping because it is felt to be safer than smoking.

However there is also emerging evidence that this mode of consumption is increasingly being used as it is discreet and much less easy to detect, hence sometimes referred to as stealth-vaping. This appears to be driving a switch to vaping to administer substances other than nicotine, notably, but not exclusively cannabis, including concentrated forms of Tetrahydrocannabinol (THC) and synthetic cannabinoids. Anecdotally this practice is known to be occurring in psychiatric inpatient settings.

This is against a backdrop of the uncertain long-term effects of vaping and the emergence of case reports of the death of otherwise healthy young persons after using ENDS to consume cannabis.

Method. Search strategy: MEDLINE, EMBASE, Cochrane Database of Systematic Reviews, Grey Literature using Medical Subject Headings (MeSH), text words relating to vaping of drugs and hand searching journals.

Statistical methods: Synthesis of data was performed using inverse variance with double arcsine transformation in MetaXL. Heterogeneity was assessed with the Cochran's Q and I².

Result. From 970 abstracts, 61 papers were selected for full text review, 18 met the inclusion criteria. The total study population for the outcome of ENDS nicotine users who also use ENDS for the consumption of illicit substances was 9098. There was significant heterogeneity with a random effects model prevalence of 17% (95%CI 7%-32%). The total study population for the outcome of cannabis users who use ENDS to consume cannabis was 52708. There was significant heterogeneity with a random effects model prevalence of 23% (95%CI 12%-37%).

Conclusion. The use of ENDS to consume illicit substances is concerning as it appears to be relatively common practice. This was most notable in studies of existing cannabis users, younger people and medical marijuana users.

Given the uncertainty of long term health consequences and poor understanding of sudden death in some users, this study highlights an emerging and substantial public health concern.

Currently there is a paucity of primary studies to elucidate the impact on health.

Self-harm on a specialist adult eating disorder unit: a retrospective cohort study of patient characteristics and outcomes

Leah Holm-Mercer*, Douglas Kohler and Agnes Ayton

Oxford Health NHS Foundation Trust

*Corresponding author.

doi: 10.1192/bjo.2021.687

Aims. Deliberate self-harm (DSH) is common but rarely studied among inpatients with eating disorders. We sought to investigate the frequency of DSH among inpatients in a specialist adult eating disorders unit, and the association of DSH with comorbidities and treatment outcomes. We also investigated changes in these parameters during the pandemic.

Method. We included the records of 70 patients consecutively admitted to Cotswold House in Oxford between April 2018 and November 2020. Data were analysed using Microsoft Excel using descriptive statistics. For comparisons, student T-tests were used for continuous variables and Chi-square tests used for categorical variables.

Result. 99% of patients were female; their ages ranged from 17 to 67 years (mean 30.7). 81% had a primary diagnosis of anorexia nervosa, and 67% had a history of DSH prior to admission.

There was a total of 100 incidences of DSH, of which 12% required transfer to a general hospital for medical treatment.

Frequency of self-harm decreased with time throughout admission (17% self-harming on admission, vs 7% at discharge, $p = 0.043$).

Compared to those with no history of DSH, patients who self-harmed during admission were more likely to be detained under the Mental Health Act (45% vs 17.4%, $p = 0.003$), and to have psychiatric comorbidities (85% vs 35%, $p = 0.001$). Patients whose self-harm required transfer for general hospital treatment had a lower mean discharge BMI (18.18kg/m² vs 20.23kg/m², $p = 0.039$), longer admission (105.9 days vs 78.1 days, $p = 0.037$), and gained weight at a slower rate (0.26kg/m²/week vs 0.43kg/m²/week, $p = 0.048$) than those who did not require transfer.

During the pandemic, the frequency of DSH doubled on the ward. Overall outcomes were similar, however mean length of admission was lower during the pandemic (67.83 vs 89.94 days, $p = 0.046$), and patients regained weight more rapidly (0.43kg/m²/week vs 0.28kg/m²/week, $p = 0.003$) than prior to it.

Conclusion. Self-harm during admission was seen in 29% of patients and was associated with the presence of comorbid psychiatric diagnoses. The frequency of DSH much reduced between admission and discharge, suggesting a beneficial effect of treatment. Medical transfer for DSH, considered as a proxy measure for severity, predicted poorer outcomes in weight restoration. We also noted an increase in rates of DSH during the pandemic, which may have resulted from a combination of increased psychosocial stressors and a reduction in admission capacity in eating disorder units.

Psychiatric presentations of patients with COVID-19: a retrospective review of 100 consecutive patients seen by liaison psychiatry services

Yousaf Iqbal^{1*}, Peter Haddad², Javed Latoo², Majid Alabdulla², Sultan Albrahim¹, Rajeev Kumar² and Ovais Wadoo¹

¹Hamad medical corporation and ²Hamad medical corporation, Qatar University

*Corresponding author.

doi: 10.1192/bjo.2021.688

Aims. Coronavirus disease 2019 (COVID-19) is associated with higher rates of psychiatric morbidity due to various factors, including quarantine, social isolation, stigma, financial difficulties and direct and indirect central nervous system impact of severe acute respiratory syndrome coronavirus 2 (SARS-Cov-2).

This study aimed to describe the psychiatric morbidity of patients with COVID-19 referred to liaison psychiatry services in Qatar.

Method. This study was a retrospective review of patient records of the first 100 consecutive SARS-Cov-2 positive patients referred to liaison psychiatry services. The study was approved by the Hamad Medical Corporation Institutional Review Board (IRB) (MRC-05-072). Data were analysed using descriptive statistics.

Result. The majority ($n = 92$) of 100 included patients were male and median age was 43 years. Patients were of diverse background with majority of South Asian (Indian, Pakistani, Bengali, Nepalese, and Afghan) ($n = 60$), followed by Qatari ($n = 18$) background. Mean length of hospital stay was 26.51 days.

35 patients had severe or critical COVID-19 pneumonia, and 67 had at least one underlying physical comorbidity. Significant psychosocial stressors other than positive SARS-Cov-2 status, including lockdown, quarantine, finances and relationships issues were identified in 48 patients.