

Magneto-hydrostatic equilibrium in starspots: dependences on color (T_{eff}) and surface gravity (g)

S. P. Rajaguru and S. S. Hasan

Indian Institute of Astrophysics, Bangalore - 560034, India,
email: rajaguru@iap.res.in

Abstract. Temperature contrasts and magnetic field strengths of sunspot umbrae broadly follow the thermal-magnetic relationship obtained from magneto-hydrostatic equilibrium. Using a compilation of recent observations, especially in molecular bands, of temperature contrasts of starspots in cool stars, and a grid of Kurucz stellar model atmospheres constructed to cover layers of sub-surface convection zone, we examine how the above relationship scales with effective temperature (T_{eff}), surface gravity g and the associated changes in opacity of stellar photospheric gas. We calculate expected field strengths in starspots and find that a given relative reduction in temperatures (or the same darkness contrasts) yield increasing field strengths against decreasing T_{eff} due to a combination of pressure and opacity variations against T_{eff} .

Keywords. Sun: magnetic fields, sunspots, stars: magnetic fields, stars: spots, stars: activity

1. Introduction

Despite a lack of deductive magnetohydrodynamic explanation for the formation and the equilibrium of a sunspot, extensive observations in combination with magneto-hydrostatic models have provided reasonable understanding of the thermal-magnetic structure of sunspots in the observable layers. Reduced temperature and gas pressure inside sunspots dictate dominantly the thermal-magnetic relationship derived from the magneto-hydrostatic balance. Such a relationship causes the Wilson depression – geometrical depression of the observable optical depth unity level within a sunspot –, which provides an explanation for the field strengths observed in sunspots and also relates the intensity (or the brightness) contrasts to field strengths. Here we examine how such a thermal-magnetic relationship scales with the stellar parameters, viz. the effective temperature T_{eff} and surface gravity g as well as the associated changes in the opacity of the stellar photospheric gas. We then discuss the implications of such scalings for the interpretations of observed field strengths. We also discuss how such scalings could be crucial players in the activity related photospheric brightness variations and their correlation with other activity measures.

2. Method of calculation

Thermal-magnetic relationship for starspots is obtained from a simplified magneto-hydrostatic (MHS) condition, that relates the magnetic field strength to the temperature at the axis of a vertical column of magnetic field (Solanki *et al.* 1993)). The radial component of the magneto-hydrostatic force-balance equation (Maltby 1977, Solanki *et al.* 1993), after neglecting the magnetic curvature force and with the use of the equation of state $P = R\rho T/\mu$, yields the thermal-magnetic relation,

$$\frac{T(r, z)}{T_e(z)} = \frac{\mu(r, z)\rho_e(z)}{\mu_e(z)\rho(r, z)} \left[1 - \frac{B_z^2(r, z)}{8\pi P_e(z)} \right], \tag{2.1}$$

where P, T, ρ, μ and R denote the gas pressure, temperature, density, mean molecular weight and the gas constant respectively. The subscript e stands for the external atmosphere. B_z is the vertical component of magnetic field and r is the radial distance from the center of the spot and z is the depth measured from continuum optical depth $\tau_{c,e}=1$. We further neglect the r -dependence of quantities in the above equation, thus the calculated quantities refer to the axis of the spot, and we refer B_z simply as B hereafter. Eqn. 2.1 is valid for each level z . The variation of external atmospheric quantities with depth z are determined by g and T_{eff} of the parent star and are taken from the grid of Kurucz stellar models constructed to cover deeper regions of the convection zone using the ATLAS9 stellar atmosphere code (Kurucz 2001). We prescribe temperature contrasts for starspots under two cases, case (i): use the empirical relation $\Delta T = (590 \cdot \log g) - 680K$ (O’Neal *et al.* 1996) to determine the effective temperature of the spot $T_{eff,spt}$ in a parent star characterised by g and T_{eff} : $T_{spt}(\tau_c = 2/3) = T_{eff,spt} = T_{eff} - \Delta T$, case (ii): the temperature contrasts are independent of g and are of a constant ratio of T_{eff} : $\Delta T = 0.3T_{eff}$ (Berdyugina 2005).

Hydrostatic equilibrium inside the spot yields, to a very good approximation (Cox and Giuli 1968),

$$P(\tau_c = 2/3) = \frac{2}{3} \frac{g}{\kappa_R(\rho, T_{eff,spt})}. \tag{2.2}$$

The density inside the spot at $\tau_c = 2/3$ is determined by solving the above equation with the use of Rosseland mean opacities κ_R from the tables of Kurucz (1993) and Alexander and Ferguson (1994). Saha’s equation is used to determine the mean molecular weight. The density difference between the $\tau_c = 2/3$ and $\tau_c = 1$ levels within the spot is assumed to be negligible, i.e., $\rho(\tau_c = 1) \approx \rho(\tau_c = 2/3)$. Using the assumption that $\rho(\tau_c = 1) = \rho_e(z = Z_w)$, i.e., the densities inside and outside the spot are equal at the




Figure 1. Variation of field strength B (solid lines) and the Wilson depression Z_w (dashed lines) against $\log(g)$ for $T_{eff} = 6000K$

level of Wilson depression Z_w (Maltby 1977), we find the depth z at which the above equality is satisfied in the parent stellar model. This gives the value of Z_w , and now we determine the only unknown quantity B at this level using Eqn. 2.1.

3. Results and discussions

Figures 1 and 2 summarise the main results of our study. Each figure shows the variation of observable field strengths $B(Z_w)$ on the spot axis as a function of $\log(g)$ for a particular T_{eff} for the two distinct kinds of variation of spot temperature given as case (i) (*red curves*) and (ii) (*green curves*) in the previous Section. The Wilson depressions, Z_w , are shown as dashed curves and the B values as solid curves. Z_w are scaled in terms of pressure scale heights H_p at the $\tau_c=1$ level in the quiet photospheres. Fig. 1, for $T_{eff} = 6000K$, shows that the B values are not very different for the two cases above, and moreover the Wilson depressions Z_w are of the order of scale heights. In contrast, Figure 2, for $T_{eff} = 5000K$, shows that the results for the above two cases are very




Figure 2. Same as Fig. 1 but for $T_{eff} = 5500K$ (top) and $5000K$ (bottom)

different: the gravity dependent temperature contrasts (as implied by the observed relation of O'Neal *et al.* (1996)) requires very strong magnetic fields for starspot equilibrium, which stems from the large values of Z_w . On the other hand, a constant value of temperature reduction (case ii above) over the range of surface gravity values used yields B values within the observed ranges (Saar 1990, Solanki 1992). Consequently, a relatively less temperature reduction and hence a less amount of gas evacuation is sufficient to attain a given value of field strength in cooler stars than that in stars hotter than about 5500K. In other words, spots of a given field strength would appear much darker in a star of $T_{eff} = 6000K$ than those in $T_{eff} = 5000K$.

According to Lockwood *et al.* (1992), younger and faster rotating stars are 'spot dominated', i.e. more flux in spots than small-scale fields and faculae, and hence grow darker as activity increases. If such a 'spot domination' is purely dependent on age (rotation) but not on color (spectral type), then it could be thought of as a phenomenon not contradictory with the g dependent temperature contrasts for spots derived by O'Neal *et al.* (1996). However, our results in Figure 2 imply unrealistically large B and Z_w values for spots in such cases. On the other hand, consistency between Lockwood *et al.*'s results requiring 'spot domination' for younger stars and the situation of case (ii) of our results requires that there be a color T_{eff} dependence of spot properties in addition to the age dependence.

Alternatively, results of Lockwood *et al.* (1992), but without the requirement of 'spot domination' in younger stars, could be made consistent with our case (ii) of less gas evacuated spots if the small-scale fields forming the 'faculae' too are of such less evacuated state. This would imply that faculae in younger and cooler stars are less bright and therefore these stars grow darker as activity increases. Interestingly, the superadiabaticity that drives the convective collapse of small-scale flux tubes indeed linearly decrease with T_{eff} and is found not to intensify such fields to a high degree of evacuation as in stars hotter than about 5000 K (Rajaguru *et al.* 2002). Hence, it would appear that the near-surface thermal structure of cool stars crucially determine the key properties of magnetic structures small and large. We conclude that the scaling of thermal and magnetic properties of starspots with both the gravity g (age) and T_{eff} (color) are crucial for a consistent interpretation of observed correlations between activity measures that sample the different heights in the outer atmospheres.

Acknowledgements

The presentation of this paper in the IAU Symposium 273 was possible due to partial support from the National Science Foundation grant numbers ATM 0548260, AST 0968672 and NASA - Living With a Star grant number 09-LWSTR09-0039.

References

- Alexander, D. R. & Ferguson, J. W. 1994, *ApJ*, 437, 879
 Berdyugina, S. 2005, *Living Rev. Solar Phys.*, 2, 8; URL: www.livingreviews.org/lrsp-2008-8
 Cox, J. P. & Giuli, R. T. 1968, *Principles of Stellar Structure, Vol. 2*, New York: Gordon & Breach, p590
 Kurucz, R. L. 1993, *ATLAS9 Stellar Atmosphere Programs and 2 km/s grid, Kurucz CD-ROM No.13. Cambridge, Mass.: Smithsonian Astrophysical Observatory.*
 Kurucz, R. L. 2001, *private communication*
 Lockwood, G. W., Skiff, B. A., Baliunas, S. L., & Radick, R. R. 1992, *Nature*, 360, 653
 Maltby, P. 1977, *Solar Phys.*, 57, 335

- O'Neal, D., Saar, S. H., & Neff, J. E. 1996, *ApJ*, 463, 766
- Rajaguru, S. P., Kurucz, R. L., & Hasan, S. S. 2002, *ApJ*, 565, L101
- Saar, S. H. 1990, in *IAU Symposium No. 138 Solar Photosphere: Structure, Convection, and Magnetic Fields*, ed. J. O. Stenflo (Kluwer: Dordrecht), p427
- Solanki, S. K., Walther, U., & Livingston, W. 1993, *A&A*, 277, 639