

NEW



LASER AND PARTICLE BEAMS

Physics of High Energy Densities

Editor in Chief

Heinrich Hora, *Department of Theoretical Physics, University of New South Wales*

Laser and Particle Beams is a new international journal which covers the generation of high intensity laser and particle beams, their interaction with materials and the physics of systems of very high densities. The fields covered have applications to nuclear fusion and to material treatment, laboratory astrophysics, plasmas and spectroscopy at extreme conditions. As well as publishing original articles the journal also publishes occasional review articles and surveys of research at particular laboratories.

Volume 1 Number 1 includes the following articles

D. D. RYUTOV and G. V. STUPAKOV Generation of High Power Ion Beams by the Oscillating Electrons

K. I. GOLDEN Strong Ion Coupling Effects at High Density of Inertial Confinement Fusion Plasmas

S. NAKAI, K. IMASAKI, S. MIYAMOTO, S. HIGAKI, T. OZAKI, A. YOSHINOCHI, H. FUJITA, K. MIMA, K. NISHIHARI, I. YABE, S. IDO, Y. OHGAKI and

C. YAMANAKA Inertial Confinement Fusion Research by Particle Beams

P. LÄDRACH A Model for Heating and Expansion of Picosecond Laser-Plasma Including Ponderomotive Force Effects

B. I. HENRY Spectral Line Limiting and Polarization Shift in Plasmas of High Particle and Energy Density
Book Reviews

Volume 1: February, May, August and November 1983

£50.00 per volume Single parts: £15.00

Airmail £15.00 extra

For further information on this and other Cambridge Journals please write to Aileen Askwith, Journals Publicity at the address below. All enquiries in North America should be sent to Cambridge University Press, 32 East 57th Street, New York, N.Y. 10022, U.S.A.

CAMBRIDGE UNIVERSITY PRESS

The Edinburgh Building, Shaftesbury Road, Cambridge CB2 2RU, England

JOURNAL OF PLASMA PHYSICS

Volume 30 Part 2 October 1983

CONTENTS

Linear conversion in a magnetized plasma with density gradient parallel to the magnetic field E. MJØLHUS	page 179
Nonlinear Rayleigh–Taylor instability in hydromagnetics B. B. CHAKRABORTY, A. R. NAYAK AND H. K. S. IYENGAR	193
Effect of finite spectral width on the modulational instability of Langmuir waves J. C. BHAKTA AND D. MAJUMDER	203
Nonlinear variations of the reflexion coefficient for lower-hybrid heating V. A. PETRŽÍLKA, R. KLÍMA AND P. PAVLO	211
A retarded time superposition principle and the relativistic collision operator K. HIZANIDIS, K. MOLVIG AND K. SWARTZ	223
On the negative resistance of double layers M. A. RAADU AND M. B. SILEVITCH	249
On the existence of a free boundary solution of the Grad–Shafranov equation NOBUO FUJII AND MASAHIDE HIRAI	255
Guided modes between upper-hybrid and plasma resonance G. J. MORALES AND J. E. MAGGS	267
Large-amplitude disturbances caused by ion-rich sheath motion A. J. COATES, P. D. EDGLEY AND J. E. ALLEN	275
The effect of thermal anisotropy on the propagation of whistler waves in mixed hot–cold electron plasmas HIROMITSU HAMABATA	291
Asymptotic solution for field-line reconnection. Compressible case of Petschek’s model V. S. SEMENOV, I. V. KUBYSHKIN AND M. F. HEYN	303
Field-line reconnection in the two-dimensional asymmetric case V. S. SEMENOV, I. V. KUBYSHKIN, M. F. HEYN AND H. K. BIERNAT	321

© Cambridge University Press 1983

CAMBRIDGE UNIVERSITY PRESS

THE PITT BUILDING, TRUMPINGTON STREET, CAMBRIDGE CB2 1RP
32 EAST 57TH STREET, NEW YORK, N.Y. 10022

Printed in Great Britain at the University Press, Cambridge