

Factors associated with non-optimal resource utilization of air ambulance for interfacility transfer of injured patients

Andrew Quirion , MD*; Mahvareh Aghari†; Brodie Nolan , MD†‡

CLINICIAN'S CAPSULE

What is known about the topic?

Early access of trauma patients to trauma centres and definitive care improves patient outcomes.

What did this study ask?

This study examines risk factors for non-optimal air transport of trauma patients via a provincial air medical transport organization.

What did this study find?

Nursing station as a sending facility, advance/primary care paramedics, and transport between 08:00 and 00:00 were risk factors for non-optimal transport.

Why does this study matter to clinicians?

These risk factors can be studied to attempt to minimize them and decrease time to definitive care for trauma patients.

0000 (OR 1.40 and 1.54, respectively). The median delay to arrival to receiving facility if a patient had a non-optimal resource use was 40 minutes.

Conclusions: Three main risk factors were identified in this study. We believe that nursing stations as a sending facility and type of paramedics crew transporting patients resulted in non-optimal resource utilization primarily due to triage of lower acuity patients. However the timing of day is more likely to be a resource availability issue and something that can be further studied and potentially improved moving forward.

RÉSUMÉ

Objectif: L'accès rapide à un traitement indiqué est associé à une amélioration des résultats chez les blessés. L'étude avait donc pour but de cerner des facteurs de risque d'utilisation non optimale des ressources, liés aux patients, aux établissements et aux ambulanciers paramédicaux, dans les mutations de blessés adultes, entre établissements, par ambulance aérienne, vers un centre de soins.

Méthode: Il s'agit d'une étude de cohortes, rétrospective, menée chez des adultes transportés de toute urgence vers un autre établissement, par Ornge. Ont été recueillis des données démographiques ainsi que des renseignements sur l'état clinique, l'établissement d'origine, le transport et les compétences des ambulanciers paramédicaux. Il y a eu analyse des données à l'aide d'un modèle de régression logistique.

Résultats: Sur 1777 transports de blessés par Ornge, 805 étaient considérés comme non optimaux, d'après l'analyse des données. Ceux chez qui le transport était jugé optimal étaient plus âgés que les autres et sous ventilation mécanique. Les facteurs de risque de transport non optimal comprenaient les mutations de patients depuis des postes de soins infirmiers (risque relatif approché [RRA] : 1,94), celles effectuées en présence de paramédicaux en soins primaires ou en soins avancés (RRA : 6,57 et 1,44, respectivement) et celles effectuées entre 8:00 et 17:00 et entre 17:00 et 00:00 (RRA : 1,40 et 1,54, respectivement). Le retard médian avant l'arrivée à l'établissement

ABSTRACT

Objective: Timely access to definitive care is associated with improved outcomes in trauma patients. The goal of this study is to identify patient, institutional and paramedic risk factors for non-optimal resource utilization for interfacility transfers of injured adult patients transported by air ambulance to a LTC.

Methods: This is a retrospective cohort study of adult emergent interfacility transports via Ornge with data collected on patient demographics, clinical status, sending facilities, transport details and paramedic qualifications. A logistic regression model was used to analyze data.

Results: 1777 injured patients undergoing transport with Ornge were analyzed with 805 of these undergoing non-optimal transport. Patients who had an optimal resource use were found to be older and mechanically ventilated. Risk factors increasing odds of non-optimal transport included patients transported from a nursing station (OR 1.94), transport with primary or advanced care paramedics (OR 6.57 and 1.44, respectively) and transport between both 0800-1700 and 1700-

From *Department of Medicine, University of Toronto, Toronto ON; †Ornge, Toronto ON; and the ‡Department of Medicine, St. Michael's Hospital, Toronto ON.

Correspondence to: Dr. Andrew Quirion, 201-3 Market Street, Toronto, ON M5E 0A3; Email: andrew.quirion@mail.utoronto.ca

d'accueil, dans les cas d'utilisation non optimale des ressources, était de 40 minutes.

Conclusion: Il s'est dégagé de l'étude trois grands facteurs de risque. Les auteurs sont d'avis que les postes de premiers soins comme établissements de départ et le type d'équipe d'ambulanciers paramédicaux présents durant les transports entraînent une utilisation non optimale des ressources, en

raison surtout du faible degré de gravité au moment du triage. Toutefois, le moment de la journée pose davantage un problème de disponibilité des ressources, point qui mériterait d'être approfondi et qui serait susceptible d'amélioration au fil du temps.

Keywords: EMS, prehospital, trauma

BACKGROUND

Early access to definitive care at a lead trauma centre has been shown to improve patient outcomes.¹⁻⁵ The use of air ambulance has become an integral part of many trauma systems and helps alleviate geographic barriers to rapid trauma centre access. Although the literature regarding the mortality benefit of air ambulance is mixed, there are multiple studies showing improved patient outcomes in the Canadian system.⁴

¹¹ Speed is often cited as the greatest benefit of air medical transport; however, multiple studies have shown improved outcomes despite no time benefit.^{7,8} These studies suggest a higher level of care that includes advanced procedures such as airway management, blood transfusion, and vasopressor use in conjunction with expedited transfer to trauma centres as part of this benefit.^{6-8,11}

There are several reasons why a severely injured patient may initially be brought to the non-trauma hospital, including inappropriate triage and identification of injuries^{12,13}; also, there may not be a trauma centre within an acceptable safe distance to transport. Patients initially brought to a non-trauma centre undergo an interfacility transfer to be taken to a specialized trauma centre for advanced care. Air ambulance services play an integral role in facilitating the interfacility transport. Delays during the interfacility transfer process resulting in delays to definitive care can have negative impacts on patient outcomes.¹⁻⁵ Previous studies have identified modifiable delays to interfacility transfer including the sending physician doing a procedure, waiting to meet a land emergency medical service (EMS) crew, and delays for diagnostic imaging.⁵

Often, there are times when the closest aircraft or most optimal type of resource (i.e., fixed wing or rotor wing) for transport is unavailable. This can occur for multiple reasons such as the optimal resource being already busy with another patient transport or is unavailable because

of weather or maintenance.⁵ When a non-optimal resource is used, there is an inherent delay to the transport of that patient; as such, there may be critically ill or injured patients who deteriorate as a result of this delay.

The objective of this study was to identify patient-, institutional-, and paramedic-level characteristics associated with non-optimal resource utilization for injured adult patients undergoing emergent interfacility transfer to a trauma centre. A better understanding of patient, institutional, and paramedic characteristics that have non-optimal resource utilization may assist with targeted interventions to reduce any delays and expedite transfer of injured patients to definitive care.

METHODS

Setting

Ornge is the sole provider of air ambulance and critical care transport in Ontario, Canada. Ornge operates a total of 12 bases, 9 of which operate fixed- or rotor-wing aircrafts (Figure 1). Ornge provides service to more than 14 million people over an area of operations that spans more than one million square kilometres. Ornge transports injured patients to and from a multitude of health care facilities varying from nine tertiary trauma centres to small remote nursing stations. Most nursing stations are serviced by fixed-wing resources and have longer transport times, often more than 1-2 hours each way and require a land ambulance transport from the landing site to the nursing station. Ornge has the largest fleet of air ambulances in Canada and includes eight fixed-wing Pilatus Next Generation PC-12 airplanes and 12 Leonardo AW-139 helicopters. In addition, Ornge operates 13 Crestline Commander land ambulances. Aircrafts are staffed with two paramedics and two pilots. A transport medicine physician is also available and provides online



Figure 1. Map of Ontario showing Ornge fixed- and rotor-wing bases.

medical control using direct telecommunication with paramedics. Paramedic staff include critical care paramedics, who are trained to administer vasopressors, blood products, and sedation, as well as to provide complex interventions including intubation, needle thoracostomy, and cricothyrotomy.

Study design

This was a retrospective cohort study of data collected in an internal Ornge database. Ethics approval was given by the Sunnybrook Health Sciences Research Ethics Board.




Figure 2. Cohort creation.

Study participants

Patients were identified from an Ornge database over a five-year period from January 1, 2013, to December 31, 2017. Patients were included as outlined below (Figure 2). All adult (aged 18 years and older) patients who underwent emergent interfacility transport for traumatic injuries during this time period were assessed for inclusion. Patients were excluded from the study if they were under 18 years of age, were transported for medical reason, were an urgent or non-urgent priority, were transported by an aircraft that

was mobile at the time of request, or had an optimal resource determined to be a land resource.

Data sources

Data were collected from two health administrative databases at Ornge. The first database used, Flight Vector, contains all information related to sending and receiving facilities, dispatch times, vital signs at time of request to transport, reason for transport, and patient age and sex. The second database used was the electronic patient care

record (ePCR), which is a computerized charting system for the paramedics. Data captured in this database include patient's vital signs while under the care of paramedics, interventions done by paramedics, time to complete patient transport, reason for transport, and a narrative text of the patient transport. Classification of a resource as optimal or non-optimal was done using time data collected in Flight Vector. This is further described below.

Patient, institutional, and paramedic characteristics that were examined included: age, sex, transporting base (name), sending facility/hospital (name), date and time of transport request, date and time paramedics arrive at patient bedside, season of transport, time of transport (day, evening, and night), vital signs on arrival of paramedics, and paramedic interventions (intubation, vasopressor use, cardiopulmonary resuscitation [CPR], and blood transfusion). Additionally, classification of sending facility (academic, community >100 beds, community <100 beds, and nursing station) was done by manually identifying the number of beds available at each sending hospital.

Defining optimal resource

The process of defining optimal resource utilization is outlined in [Figure 3](#). All possible interfacility transfers were determined using the study cohort by grouping all patient transfers that had identical sending and receiving facilities.

A unique resource was defined as a specific type of aircraft (rotor or fixed wing) that was attached to a specific base; however, multiples of the same type of aircraft within a single base were considered one resource. For example, in the Thunder Bay base, there are two fixed-wing resources and a single rotor-wing resource. In these methods, there was a distinction between fixed-wing and rotor-wing resources, but two fixed-wing resources were considered the same "unique resource" for this methodology. Then, each unique resource that transported the sending-receiving pair was identified. Resources were considered unique if they differed in base or mode of transport (rotor wing, fixed wing, or land). For each unique sending-receiving pair, estimated transfer times for each resource were calculated using a modular method by breaking down the total time to definitive care into intervals based on the major steps of the transfer process.¹⁵ The median times for these intervals was then calculated and summed for each unique resource. The fastest resource was the one that had the minimum

sum of medians. This modular process of the sum of median time intervals has been used in previous studies to estimate optimal resource use for air ambulance services.¹⁵ Furthermore, the same methodology has been used internally at Ornge as a decision support tool to aid planning of resource utilization.

The optimal resource was determined to be the fastest asset (rotor-wing, fixed-wing, or land ambulance) from the closest base. If the distance between sending and receiving was less than 100 km and the mean time difference between land and air resources was less than 10 minutes, then the fastest land resource was considered the optimal resource. These distance cut-offs are used internally at Ornge for flight planning and resource allocation. If the resource that serviced the transfer was mobile at the time of call request, it was removed from the study cohort since time comparisons would be unreliable without an accurate acceptance location. If an optimal resource was determined to be a land resource, they were removed from the final study cohort.

Statistical methods

Descriptive statistics

Descriptive statistics were used to assess the distribution for all variables of interest in each group. Continuous factors were assessed for normality by evaluating kurtosis and skewness and are summarized as means and standard deviations (SD) or medians and interquartile range (IQR) for normal and non-normally distributed data, respectively, and categorical variables are displayed as counts and percentages.

Unadjusted bivariate analyses were performed to assess differences between patients with a non-optimal transfer strategy and patients with an optimal transfer strategy. Continuous variables were compared using the independent samples Student's *t*-test for normally distributed variables and Wilcoxon rank-sum test for non-normally distributed variables. Categorical variables were compared using the Chi-square test. Two-sided *p*-values were reported in all cases, and *p*-values <0.05 were considered significant for all analyses.

Risk factors associated with non-optimal resource use

Patient, institutional, and paramedic characteristics were examined to assess association with non-optimal




Figure 3. Optimal resource utilization categorization.

resource use. These included patient age, sex, vital signs (heart rate, respiratory rate, systolic blood pressure, Glasgow Coma Scale [GCS], and oxygen saturation), ventilator dependence, sending facility type, season, time of day, and paramedic level of care. As this model was exploratory, there was no single key predictor variable; thus, all the above characteristics were all assessed for possible inclusion in the final model. A backward selection technique using a *p*-value cut-off of 0.2 was used to determine which variables to include in the final analysis.

An adjusted logistic regression model was used to explore factors associated with a non-optimal transfer strategy. Patients with any missing data for one or more of the predictor variables of interest were excluded from the final model. Missing data resulted in exclusion of <5% of observations. Lastly, a generalized estimating equation (GEE) logistic regression model accounting for clustering by sending facility was performed.

Assumptions for logistic regression were then assessed, including adequate model fit and absence of influential observations. Model fit was assessed with *c*-statistic. Multicollinearity was assessed using a variation inflation factor (VIF) of four as the cut-off for

exclusion. A 50-50 split sample technique comparing difference in *c*-statistics was used to assess internal validity.

All statistical analyses were conducted using SAS Studio, version 3.4 (SAS Institute, North Carolina, USA).

Missing data

All data were reviewed for completeness and to ensure there were no implausible values. If the optimal resource was missing or unable to be calculated or if the resource was mobile at the time of transfer request, they were removed. Lastly, if the optimal resource was deemed to be a land resource, they were also removed from further analysis.

RESULTS

There were a total of 31,123 adult emergent interfacility transports with Ornge between January 1, 2013, and December 31, 2017. Of these, 24,608 were older than 18 years of age, with 2,735 of those being trauma patients. Of these patients, a total of 1,777 had fixed- or rotor-wing transport as their optimal method, with

972 receiving optimal transport and 805 receiving non-optimal resource use (Figure 2).

The patients receiving optimal resource use were older, with a median age of 48, as compared with 45 (Table 1). A higher percentage of patients who underwent optimal resource utilization were mechanically ventilated, as compared with those who underwent non-optimal resource utilization (24.4% v. 17.4%, respectively) and had a GCS of less than 8 (30.3% v. 24.2%, respectively).

An adjusted logistical regression model accounting for clustering by sending facility was used to determine which factors were associated with higher odds of non-optimal transport (Table 2). Patients being transported from a nursing station had higher odds of non-optimal resource use, as compared with an academic centre (odds ratio [OR] 1.94). Transport with primary or advanced care paramedics had a higher odds of non-optimal resource use, as compared with critical care paramedics (OR 6.57 and 1.44, respectively). As compared with time of transport between 00:00 and 08:00, both daytime (08:00–17:00) and evening (17:00–00:00) times of transport had higher odds of non-optimal resource use (OR 1.40 and 1.54, respectively). There was no significant difference in OR for age, sex, need for mechanical ventilation or time of year.

Only 32.9% of patients transported from a nursing station were brought directly to a trauma centre for assessment, as compared with more than 90% of all transports from academic and community hospitals. Most of the time patients transported from a nursing station were transported to a facility with computed tomography (CT) scanning capabilities if they were not transported to a trauma centre. Patients transported from academic and community centres who were not transported to a trauma centre were transported to centres with neurosurgical capabilities for isolated head injuries. The median delay to arrival to receiving facility if a patient had a non-optimal resource use was 40 minutes (IQR 18–47 min).

DISCUSSION

Multiple studies have shown that early transport of trauma patients to definitive care improves patient outcomes.^{1–4,16–18} In Ontario, air ambulance covers a massive land mass, with the majority of tertiary centres in its southern region.^{16,17} Previous studies have examined

when delays occurred during transport for the various types of Helicopter Emergency Medical Services transports of trauma patients; some of the major delays included weather, refueling and mechanical checks, and procedures at the sending facility and land EMS transport to the sending facility from the aircraft.⁵ Our study adds to this existing literature by identifying patient-, institutional-, and paramedic-level factors associated with non-optimal resource use.

In our study, resource was defined as a two-level variable: originating base and category of aircraft (fixed wing v. rotor wing). Further, the variable “category of aircraft” was not independent of the variable “base,” as certain bases had only specific categories of aircraft. Thunder Bay has two fixed-wing aircrafts and one rotor-wing aircraft; Timmins and Sioux Lookout operate only one fixed-wing aircraft each. Kenora, Toronto, London, Sudbury, Ottawa, and Moosonee operate only a rotor-wing aircraft.

Our study identified three factors associated with non-optimal resource use to emergent interfacility transfer of injured patients by air ambulance: size of sending facility, paramedic level of care, and time of day.

The type of sending facility was associated with non-optimal resource use. In comparison with academic centres, patients being transferred from nursing stations had a higher odds of non-optimal resource use. This is likely multifactorial. Many of the nursing stations in Ontario are remote, with long distances and, thus, long transport times for patients undergoing interfacility transport. Because of this, once an asset is engaged in transporting a patient, it is likely to be unavailable for a number of hours to complete that initial patient transport. As this resource would then be unavailable to transport another patient during this period, a non-optimal resource would be deployed. Furthermore, most nursing stations are further north and more prone to inclement weather than southern Ontario. It is possible that a non-optimal resource was deployed as it was able to transport that patient as the optimal resource was unable to do so because of weather conditions along that route.

A non-optimal resource was used for 45% of all emergent interfacility transports. Advanced and critical care resources are limited across the province, with the same resources being used to transport emergent transports (i.e., going for life-saving interventions or higher-level care) and non-urgent transports (i.e., repatriation to closer hospital with no escalation of care). Currently, if there are no emergent transports pending, advanced

Table 1. Baseline characteristics of study cohort

Patient characteristics	Total cohort (N = 1,777)	Optimal resource utilization (N = 972)	Non-optimal resource utilization (N = 805)	p-value
Age, median (IQR)	46 (30,62)	48 (31,63)	45 (29,60)	0.039 [†]
Sex, n (%)				
Male	1305 (73.4)	723 (74.4)	582 (72.4)	0.34 [‡]
Female	471 (26.5)	249 (25.6)	222 (27.6)	
Heart rate >100 or <50, n (%)	559 (31.5)	320 (32.9)	239 (29.7)	0.14 [‡]
Respiratory rate >29 or <10, n (%)	207 (11.7)	108 (11.1)	99 (12.3)	0.44 [‡]
Systolic blood pressure <90, n (%)	137 (7.7)	77 (7.9)	60 (7.5)	0.71 [‡]
Oxygen saturation <90, n (%)	199 (11.2)	107 (11.1)	92 (11.4)	0.78 [‡]
Glasgow Coma Scale, n (%)				0.011 [‡]
13–15	1259 (70.9)	660 (67.9)	599 (74.4)	
9–12	28 (1.6)	17 (1.8)	11 (1.4)	
<8	490 (27.5)	295 (30.3)	195 (24.2)	
Sending facility, n (%)				<0.001 [‡]
Academic	59 (3.3)	35 (59.3)	24 (40.7)	
Community hospital >100 beds	676 (38.2)	401 (59.3)	275 (40.7)	
Community hospital <100 beds	843 (47.3)	463 (54.9)	380 (45.1)	
Nursing station	198 (11.2)	70 (35.2)	128 (64.3)	
Mechanically ventilated, n (%)	377 (21.2)	237 (24.4)	140 (17.4)	<0.001 [‡]
Vasopressors, n (%)	95 (5.4)	60 (6.2)	35 (4.4)	0.09 [‡]
Blood transfusion, n (%)	139 (7.8)	85 (8.7)	54 (6.7)	0.11 [‡]

IQR = interquartile range; Missing = values missing in original dataset.
[†]Student's t-test
[‡]Wilcoxon rank sum
[‡]Chi-square

Table 2. Adjusted logistic regression models of odds of having non-optimal transfer strategy with and without accounting for clustering by sending facility

Characteristic	Adjusted OR (95% CI)	p-value
Age	1.00 (0.99, 1.00)	0.32
Sex (ref = male)	0.92 (0.75, 1.12)	0.41
Mechanically vented	0.86 (0.66, 1.13)	0.28
Sending facility (ref = academic)		
Community >100 beds	1.02 (0.64, 1.61)	0.94
Community <100 beds	1.08 (0.66, 1.79)	0.75
Nursing station	1.94 (1.14, 3.32)	0.015
Level of care required (ref = critical care crew)		
Primary care crew	6.57 (2.99, 14.50)	<0.001
Advanced care crew	1.44 (1.10, 1.89)	0.007
Season (ref = summer)		
Fall	1.01 (0.77, 1.33)	0.92
Winter	1.04 (0.78, 1.39)	0.79
Spring	0.99 (0.76, 1.29)	0.93
Time of day (ref = 00:00–07:59)		
08:00–16:59	1.40 (1.04, 1.89)	0.027
17:00–23:59	1.54 (1.15, 2.05)	0.003

CI = confidence interval; OR = odds ratio; ref = reference category.

and critical care resources are deployed to service these non-urgent transports when they require a higher level of care during transport. These non-urgent transports may prevent an asset from being available to transport an emergent patient, resulting in a non-optimal resource being used. One strategy to mitigate this may be to reserve advanced and critical care resources for emergent transports only. This could be done by not sending an available resource on a non-urgent transport if it is the only resource available in a select area of the province. Additionally, partnering with local hospitals to send nurse escorts to provide a higher level of care during transports may allow patients to be transported with primary paramedic crews safely, again allowing for advanced and critical care paramedics to prioritize emergent patient transports.

Not all patients were transported directly to a trauma centre. As mentioned earlier, some injured patients are transported for advanced diagnostic imaging such as a CT scan that is unavailable at their current health care facility. For many of these patients, it may be reasonable to have them transported for a CT scan at a non-trauma

centre to assess them for significant injuries rather than transport them all directly to a trauma centre to avoid overburdening the regional trauma centre.

Time of day was also associated with optimal resource use with transports between 08:00–17:00 and 17:00–00:00 having more non-optimal transports. This may be because of fewer new transports booked overnight, resulting in higher aircraft availability and optimal resources being assigned.

There were several limitations for our study. The first being that this was a retrospective study that can often lead to missing data and exclusion of participants. Because of the retrospective aspect of our analysis, we were unable to determine the location of a resource at the time a transfer was requested unless that resource was at the base. Therefore, if a resource was mobile at the time of request, we were unable to include these transfers in our study as we could not reliably identify whether they would have been the optimal resource. Another limitation of this study was that we were unable to identify the specific reasons why a non-optimal resource was used. Unfortunately, the databases used do not provide reason for resource assignment or make note of inclement weather or mechanical or staffing issues. Lastly, our data was limited to a prehospital database; we are unable to report other variables of interest such as severity of illness or injuries identified. This also limited our ability to explore the impact of non-optimal resource use on in-hospital outcomes and mortality. However, it can be inferred from the literature that severely injured patients have increased mortality when they experience delays to definitive care and transport to a trauma centre.^{1–4}

CONCLUSION

In conclusion, there were three factors associated with non-optimal resource use to emergent interfacility transfer of injured patients by air ambulance. These included the size of sending facility, paramedic level of care, and time of day. Further research is required to explore specific causes of non-optimal resource use of air ambulance and to assess for any impact on patient-centred outcomes.

Acknowledgements: Meetings presented at: Canadian Association of Emergency Physicians Annual Conference 2019.

Competing interest: None.

REFERENCES

1. MacKenzie EJ, Rivara FP, Jurkovich GJ, et al. A national evaluation of the effect of trauma-center care on mortality [Reduced Mortality in Severely Injured Patients]. *N Engl J Med* 2006;354(4):366–78.
2. Demetriades D, Martin M, Salim A, et al. Relationship between American College of Surgeons trauma center designation and mortality in patients with severe trauma (injury severity score >15). *J Am Coll Surg* 2006; 202(2):212–5.
3. Haas B, Jurkovich GJ, Wang J, et al. Survival advantage in trauma centers: expeditious intervention or experience? *J Am Coll Surg* 2009;208(1):28–36.
4. Haas B, Stukel TA, Gomez D, et al. The mortality benefit of direct trauma center transport in a regional trauma system: a population-based analysis. *J Trauma Acute Care Surg* 2012;72(6):1510–15.
5. Nolan B, Tien H, Sawadsky B, et al. Comparison of helicopter emergency medical services transport types and delays on patient outcomes at two level I trauma centers. *Prehosp Emerg Care* 2017;21(3):327–33.
6. Chen X, Gestring ML, Rosengart MR, et al. Speed is not everything: identifying patients who may benefit from helicopter transport despite faster ground transport. *J Trauma Acute Care Surg* 2018;84(4):549–57.
7. Brown JB, Gestring ML, Guyette FX, et al. Helicopter transport improves survival following injury in the absence of a time-saving advantage. *Surgery* 2016;159(3):947–59.
8. Taylor CB, Stevenson M, Jan S, et al. A systematic review of the costs and benefits of helicopter emergency medical services. *Injury* 2010;41(1):10–20.
9. Buchanan IM, Coates A, Sne N. Does mode of transport confer a mortality benefit in trauma patients? Characteristics and outcomes at an Ontario lead trauma hospital. *CJEM* 2016;18(5):363–9.
10. Chen X, Gestring ML, Rosengart MR, et al. Logistics of air medical transport: when and where does helicopter transport reduce prehospital time for trauma. *J Trauma Acute Care Surg* 2018;85:174–81.
11. Thomas F, Wisham J, Clemmer TP, Orme Jr JF, Larsen KG. Outcome, transport times, and costs of patients evacuated by helicopter versus fixed-wing aircraft. *West J Med* 1990;153(1):40–3.
12. Brown JB, Forsythe RM, Stassen NA, Gestring ML. The National Trauma Triage Protocol: can this tool predict which patients with trauma will benefit from helicopter transport? *J Trauma Acute Care Surg* 2012;73(2):319–25.
13. Lehmann RK, Arthurs ZM, Cuadrado DG, et al. Trauma team activation: simplified criteria safely reduces overtriage. *Am J Surg* 2007;193(5):630–5.
14. Desmettre T, Yeguiayan J, Coadou H, et al. Impact of emergency medical helicopter transport directly to a university hospital trauma center on mortality of severe blunt trauma. *Crit Care* 2012;16:R170.
15. Giang CW, Donmez B, Fatahi A, Ahgari M, MacDonald RD. Supporting air versus ground vehicle decisions for interfacility medical transport using historical data. *IEEE Trans Hum Mach Syst* 2014;44(1):55–65.

16. Baxt WG, Moody P, Cleveland HC, et al. Hospital-based rotorcraft aeromedical emergency care services and trauma mortality: a multicenter study. *Ann Emerg Med* 1985;14(9):859–64.
17. Galvagno SM, Haut ER, Zafar SN, et al. Association between helicopter vs ground emergency medical services and survival for adults with major trauma. *JAMA* 2012;307(15):1602–10.
18. Doucet J, Bulger E, Sanddal N, et al. Appropriate Use of Helicopter Emergency Medical Services for Transport of Trauma Patients: Guidelines from the Emergency Medical System Subcommittee, Committee on Trauma, American College of Surgeons. *J Trauma Acute Care Surg* 2013; 75(4): 734–741.
19. Stewart K, Garwe T, Bhandari N, Danford B, Albrecht R. Factors associated with the use of helicopter inter-facility transport of trauma patients to tertiary trauma centers within an organized rural trauma system. *Prehosp Emerg Care* 2016;20:601–8.