
INSTRUCTIONS TO CONTRIBUTORS

The *Journal of Helminthology* publishes papers on all aspects of animal parasitic helminths, particularly those of medical or veterinary importance.

Manuscripts, which must be in English or French (with an English summary), should be addressed to:

The Editor, Journal of Helminthology
London School of Hygiene and Tropical Medicine
Keppel Street, London WC1E 7HT
England

Two copies of a typescript, on size A4 paper with double spacing, should be submitted. Papers should be preceded by a short abstract and will normally have the following sections: brief introduction (unheaded); Materials and Methods; Results; Discussion; Acknowledgements; References. However, the form of the paper may vary, depending on its subject matter; recent past issues should be consulted for a suitable form. Illustrations should be drawn in Indian ink, preferably not more than double the final size. Care should be taken that all illustrations fit into the format of the Journal. The maximum size an illustration may be printed is 13.5 × 18 cm. Where many separate drawings are made, some indication of how they may be grouped to make a corporate plate without undue wastage of space should be indicated. Some indication of scale should normally be given on the figure. Photocopies of illustrations should also be enclosed for refereeing purposes. Lettering and numbering, which must be of a high standard, should be added by the author, with due regard for subsequent reduction.

Photographs should be glossy prints, preferably of the same size as they are to appear in the Journal (maximum size 13.5 × 18 cm). Composite prints must be mounted and can have the separate photographs abutting; they will then have a separating line inserted by the printers. All figures and letters on photographs must be inserted by the author.

Information should not be repeated in the text and in tables or figures. The legends to tables and to figures should be sufficiently detailed for the information to be understood without reference to the text.

References should be given in alphabetical order with the full title of the journal. The following are examples:

DUKE, B. O. L. (1971) The ecology of onchocerciasis in man and animals. In *Ecology and physiology of parasites* (ed A. M. Fallis) pp. 213-222. Adam Hilger Ltd.: London.

JAMES, C. and WEBBE, G. (1973) A comparison of Egyptian and East African strains of *Schistosoma haematobium*. *Journal of Helminthology*, **47**, 49-59.

50 offprints are provided free of charge; additional copies may be ordered at the proof stage.

Contents

	Pages
An appreciation of Academician K. I. Skrjabin, founder of the Soviet Helminthological School, on the occasion of the centenary of his birth	171-172
Factors affecting the acquisition of resistance against <i>Schistosoma mansoni</i> in the mouse. I. Demonstration of resistance to reinfection using a model system that involves perfusion of mice within three weeks of challenge. M. DOENHOFF <i>et al.</i>	173-186
Factors affecting the acquisition of resistance against <i>Schistosoma mansoni</i> in the mouse. II. The time at which resistance to reinfection develops. E. LONG, M. DOENHOFF and J. BAIN	187-191
Experimental infection of <i>Ancylostoma caninum</i> in mice. III. Distribution of larvae after repeated exposures to infection. M. K. BHOPALE and G. N. JOHRI	193-198
Balanotaeniidae fam. n. and <i>Balanotaenia newguineensis</i> sp. n. (Cestoidea; Caryophyllidea) from <i>Tandanus</i> (Siluriformes: Plotosidae) in New Guinea. J. S. MACKIEWICZ and D. BLAIR	199-203
Tapeworms of the Chaco Boreal, Paraguay, with two new species. G. D. SCHMIDT and R. L. MARTIN	205-209
<i>Allohymenolepis palawanensis</i> sp. n. (Cyclophyllidea: Hymenolepididae) from the Philippine bird, <i>Nectarinia jugularis</i> (Tweeddale 1878). T. L. DEARDORFF, G. D. SCHMIDT and R. E. KUNTZ	211-213
<i>Paratrichosoma crocodilus</i> n. gen. n. sp. (Nematoda: Trichosomoididae) from the skin of the New Guinea crocodile. R. W. ASHFORD and R. MULLER	215-220
The influence of physical factors on the behaviour and infectivity of miracidia of <i>Schistosoma mansoni</i> and <i>S. haematobium</i> . III. Effect of contact time and dispersion in static and flowing waters. C. JAMES and S. K. PRAH	221-226
Studies with <i>Brugia pahangi</i> . 18. Anthelmintic effects of stibocaptate. Research Note. D. A. DENHAM, R. R. SUSWILLO and R. ROGERS	227-228
Occurrence of <i>Angiostrongylus cantonensis</i> in the heart of a spider monkey. Research Note. R. C. KO	229
Revision of the genus <i>Paracooperia</i> Travassos, 1935 (Nematoda: Trichostrongylidae). L. M. GIBBONS	231-249
Life history of <i>Genarchopsis goppo</i> Ozaski (Trematoda: Hemiuridae) from the freshwater fish <i>Channa punctata</i> R. MADHAVI	251-259
SUPPLEMENTARY REVIEW ARTICLE	
Ascaridoid nematodes of amphibians and reptiles: <i>Gedoelstascaris</i> n. g. and <i>Ortleppascaris</i> n. g. J. F. A. SPRENT	261-282