



COMPOSITIO MATHEMATICA

Tangent cones to generalised theta divisors and generic injectivity of the theta map

George H. Hitching and Michael Hoff

Compositio Math. **153** (2017), 2643–2657.

[doi:10.1112/S0010437X17007539](https://doi.org/10.1112/S0010437X17007539)



FOUNDATION
COMPOSITIO
MATHEMATICA



LONDON
MATHEMATICAL
SOCIETY
EST. 1865



Tangent cones to generalised theta divisors and generic injectivity of the theta map

George H. Hitching and Michael Hoff

ABSTRACT

Let C be a Petri general curve of genus g and E a general stable vector bundle of rank r and slope $g - 1$ over C with $h^0(C, E) = r + 1$. For $g \geq (2r + 2)(2r + 1)$, we show how the bundle E can be recovered from the tangent cone to the generalised theta divisor Θ_E at \mathcal{O}_C . We use this to give a constructive proof and a sharpening of Brivio and Verra’s theorem that the theta map $SU_C(r) \dashrightarrow |r\Theta|$ is generically injective for large values of g .

1. Introduction

Let C be a nonhyperelliptic curve of genus g and $L \in \text{Pic}^{g-1}(C)$ a line bundle with $h^0(C, L) = 2$ corresponding to a general double point of the Riemann theta divisor Θ . It is well known that the projectivised tangent cone to Θ at L is a quadric hypersurface R_L of rank less than or equal to 4 in the canonical space $|K_C|^*$, which contains the canonically embedded curve.

Quadrics arising from tangent cones in this way have been much studied: Green [Gre84] showed that the R_L span the space of all quadrics in $|K_C|^*$ containing C ; and both Kempf and Schreyer [KS88] and Ciliberto and Sernesi [CS92] have used the quadrics R_L in various ways to give new proofs of Torelli’s theorem.

In another direction: via the Riemann–Kempf singularity theorem [Kem73], one sees that the rulings on R_L cut out the linear series $|L|$ and $|K_C L^{-1}|$ on the canonical curve. Thus the data of the tangent cone and the canonical curve allow one to reconstruct the line bundle L . In this paper we study a related construction for vector bundles of higher rank.

Let $V \rightarrow C$ be a semistable vector bundle of rank r and integral slope h . We consider the set

$$\{M \in \text{Pic}^{g-1-h}(C) : h^0(C, V \otimes M) \geq 1\}. \quad (1)$$

It is by now well known that for general V , this is the support of a divisor Θ_V algebraically equivalent to a translate of $r \cdot \Theta$. If V has trivial determinant, then in fact Θ_V , when it exists, belongs to $|r\Theta|$.

For general V , the projectivised tangent cone $\mathcal{T}_M(\Theta_V)$ to Θ_V at a point M of multiplicity $r + 1$ is a determinantal hypersurface of degree $r + 1$ in $|K_C|^*$ (see, for example, Casalaina-Martin and Teixidor i Bigas [CT11]). Our first main result (§ 3.2) is a construction which from $\mathcal{T}_M(\Theta_V)$

Received 3 November 2016, accepted in final form 9 June 2017, published online 13 September 2017.

2010 Mathematics Subject Classification 14H60, 14H51, 14M12, 14B10, 13D10 (primary), 14C34 (secondary).

Keywords: vector bundles, generalised theta divisors, Brill–Noether theory.

The first named author gratefully acknowledges financial support from Høgskolen i Oslo Oppstarts- og Akkvisisjonsmidler 2015. He also thanks the Universität des Saarlandes for hospitality. The second named author was partially supported by the DFG grant SPP 1489 Schr. 307/5-2.

This journal is © Foundation Compositio Mathematica 2017.

recovers the bundle $V \otimes M$, up to the involution $V \otimes M \mapsto K_C \otimes M^{-1} \otimes V^*$. This is valid whenever $V \otimes M$ and $K_C \otimes M^{-1} \otimes V^*$ are globally generated.

We apply this construction to give an improvement of a result of Brivio and Verra [BV12]. To describe this application, we need to recall some more objects. Write $SU_C(r)$ for the moduli space of semistable bundles of rank r and trivial determinant over C . The association $V \mapsto \Theta_V$ defines a map

$$\mathcal{D}: SU_C(r) \dashrightarrow |r\Theta| = \mathbb{P}^{r^g-1}, \quad (2)$$

called the *theta map*. Drezet and Narasimhan [DN89] showed that the line bundle associated to the theta map is the ample generator of the Picard group of $SU_C(r)$. Moreover, the indeterminacy locus of \mathcal{D} consists of those bundles $V \in SU_C(r)$ for which (1) is the whole Picard variety. This has been much studied; see, for example, Pauly [Pau10], Popa [Pop99] and Raynaud [Ray82].

Brivio and Verra [BV12] showed that \mathcal{D} is generically injective for a general curve of genus $g \geq \binom{3r}{r} - 2r - 1$, partially answering a conjecture of Beauville [Bea06, §6]. We apply the aforementioned construction to give the following sharpening of Brivio and Verra's result.

THEOREM 1.1. *For $r \geq 2$ and C a Petri general curve of genus $g \geq (2r+2)(2r+1)$, the theta map (2) is generically injective.*

In addition to giving the statement for several new values of g when $r \geq 3$ (our lower bound for g depends quadratically on r rather than exponentially), our proof is constructive, based on the method mentioned above for explicitly recovering the bundle V from the tangent cone to the theta divisor at a point of multiplicity $r+1$. This gives a new example, in the context of vector bundles, of the principle apparent in [KS88] and [CS92] that the geometry of a theta divisor at a sufficiently singular point can encode essentially all the information of the bundle and/or the curve.

Our method works for $r=2$, but in this case much more is already known: Narasimhan and Ramanan [NR69] showed, for $g=2$ and $r=2$, that \mathcal{D} is an isomorphism $SU_C(2) \xrightarrow{\sim} \mathbb{P}^3$, and van Geemen and Izadi [vGI01] generalised this statement to nonhyperelliptic curves of higher genus. Note that our proof of Theorem 4.1 is not valid for hyperelliptic curves (see Remark 4.3).

Here is a more detailed summary of the paper. In §2, we study semistable bundles E of slope $g-1$ for which the *Petri trace map*

$$\bar{\mu}: H^0(C, E) \otimes H^0(C, K_C \otimes E^*) \rightarrow H^0(C, K_C)$$

is injective. A bundle E with this property will be called *Petri trace injective*. We prove that for large enough genus, the theta divisor of a generic $V \in SU_C(r)$ contains a point M of multiplicity $r+1$ such that $V \otimes M$ and $K_C \otimes M^{-1} \otimes V^*$ are Petri trace injective and globally generated.

Now suppose that E is a vector bundle of slope $g-1$ with $h^0(C, E) \geq r+1$. If Θ_E is defined, then the tangent cone to Θ_E at \mathcal{O}_C is a determinantal hypersurface in $|K_C|^* = \mathbb{P}^{g-1}$ containing the canonical embedding of C . We prove (Proposition 3.3 and Corollary 3.5) that if C is a general curve of genus $g \geq (2r+2)(2r+1)$, and E a globally generated Petri trace injective bundle of rank r and slope $g-1$ with $h^0(C, E) = r+1$, then the bundle E can be reconstructed up to the involution $E \mapsto K_C \otimes E^*$ from a certain determinantal representation of the tangent cone to Θ_E at \mathcal{O}_C . By a classical result of Frobenius (whose proof we sketch in Proposition 3.7), any two such representations are equivalent up to transpose. The generic injectivity of the theta map for a Petri general curve (Theorem 4.1) can then be deduced by combining these facts and the statement in §2 that the theta divisor of a general $V \in SU_C(r)$ contains a suitable point of multiplicity $r+1$.

We assume throughout that the ground field is \mathbb{C} . The reconstruction of E from its tangent cone in § 3.2 is valid for an algebraically closed field of characteristic zero or $p > 0$ not dividing $r + 1$.

2. Singularities of theta divisors of vector bundles

2.1 Petri trace injective bundles

Let C be a projective smooth curve of genus $g \geq 2$. Let $V \rightarrow C$ be a stable vector bundle of rank $r \geq 2$ and integral slope h , and consider the locus

$$\{M \in \text{Pic}^{g-1-h}(C) : h^0(C, V \otimes M) \geq 1\}. \tag{3}$$

If this is not the whole of $\text{Pic}^{g-1-h}(C)$, then it is the support of the theta divisor Θ_V .

The theta divisor of a vector bundle is a special case of a *twisted Brill–Noether locus*

$$B_{1,g-1-h}^n(V) := \{M \in \text{Pic}^{g-1-h}(C) : h^0(C, V \otimes M) \geq n\}. \tag{4}$$

The following is central in the study of these loci (see, for example, Teixidor i Bigas [TiB14, § 1]): For $E \rightarrow C$ a stable vector bundle, we consider the *Petri trace map*:

$$\bar{\mu}: H^0(C, E) \otimes H^0(C, K_C \otimes E^*) \xrightarrow{\mu} H^0(C, K_C \otimes \text{End } E) \xrightarrow{\text{tr}} H^0(C, K_C). \tag{5}$$

Then for $E = V \otimes M$ and $M \in B_{1,g-1-h}^n(V) \setminus B_{1,g-1-h}^{n+1}(V)$, the Zariski tangent space to the twisted Brill–Noether locus $B_{1,g-1-h}^n(V)$ at M is exactly $\text{Im}(\bar{\mu})^\perp$. This motivates the following definition.

DEFINITION 2.1. Suppose $E \rightarrow C$ is a vector bundle with $h^0(C, E) = n \geq 1$. If the map μ above is injective, we will say that E is *Petri injective*. If the composed map $\bar{\mu}$ is injective, we will say that E is *Petri trace injective*.

Remark 2.2. (1) Clearly, a Petri trace injective bundle is Petri injective. For line bundles, the two notions coincide.

(2) Suppose $V \in U_C(r, d)$ where $U_C(r, d)$ is the moduli space of semistable rank r vector bundles of degree d . If $E = V \otimes M$ is Petri trace injective for $M \in \text{Pic}^e(C)$, then $B_{1,e}^n(V)$ is smooth at M and of the expected dimension

$$h^1(C, \mathcal{O}_C) - h^0(C, V \otimes M) \cdot h^1(C, V \otimes M).$$

(3) We will also need to refer to the usual generalised Brill–Noether locus

$$B_{r,d}^n = \{E \in U_C(r, d) : h^0(C, E) \geq n\}.$$

If E is Petri injective then this is smooth and of the expected dimension

$$h^1(C, \text{End } E) - h^0(C, E) \cdot h^1(C, E)$$

at E . See, for example, Grzegorzczuk and Teixidor i Bigas [GT09, § 2].

(4) Petri injectivity and Petri trace injectivity are open conditions on families of bundles $\mathcal{E} \rightarrow C \times B$ with $h^0(C, \mathcal{E}_b)$ constant. Later, we will discuss the sense in which these properties are ‘open’ when $h^0(C, \mathcal{E}_b)$ may vary.

We will also need the notion of a Petri general curve.

DEFINITION 2.3. A curve C is called *Petri general* if every line bundle on C is Petri injective.

By [Gie82], the locus of curves which are not Petri general is a proper subset of the moduli space M_g of curves of genus g , the so-called *Gieseker–Petri locus*. The hyperelliptic locus is contained in the Gieseker–Petri locus. Apart from this, in general not much is known about the components of the Gieseker–Petri locus and their dimensions. For an overview of known results, we refer to [TiB88, Far05, BS11] and the references cited therein.

PROPOSITION 2.4. Suppose V is a stable bundle of rank r and integral slope h . Suppose $M_0 \in \text{Pic}^{g-1-h}(C)$ satisfies $h^0(C, V \otimes M_0) \geq 1$, and furthermore that $V \otimes M_0$ is Petri trace injective. Then the theta divisor $\Theta_V \subset \text{Pic}^{g-1-h}(C)$ is defined. Furthermore, we have equality $\text{mult}_{M_0} \Theta_V = h^0(C, V \otimes M_0)$.

Proof. Write $E := V \otimes M_0$. It is well known that via Serre duality, $\bar{\mu}$ is dual to the cup product map

$$\cup : H^1(C, \mathcal{O}_C) \rightarrow \text{Hom}(H^0(C, E), H^1(C, E)).$$

By hypothesis, therefore, \cup is surjective. Since E has Euler characteristic zero, $h^0(C, E) = h^1(C, E)$. Hence there exists $b \in H^1(C, \mathcal{O}_C)$ such that $\cdot \cup b : H^0(C, E) \rightarrow H^1(C, E)$ is injective. The tangent vector b induces a deformation of M_0 and hence of E , which does not preserve any nonzero section of E . Therefore, the locus

$$\{M \in \text{Pic}^{g-1-h}(C) : h^0(C, V \otimes M) \geq 1\}$$

is a proper sublocus of $\text{Pic}^{g-1-h}(C)$, so Θ_V is defined. Now we can apply Casalaina-Martin and Teixidor i Bigas [CT11, Proposition 4.1] to obtain the desired equality $\text{mult}_{M_0} \Theta_V = h^0(C, V \otimes M_0)$. \square

2.2 Existence of good singular points

In this section, we study global generatedness and Petri trace injectivity of the bundles $V \otimes M$ for $M \in B_{1,g-1}^{r+1}(V)$ for general C and V . The main result of this section is the following theorem.

THEOREM 2.5. Suppose C is a Petri general curve of genus $g \geq (2r+2)(2r+1)$ and $V \in SU_C(r)$ a general bundle. Then there exists $M \in \Theta_V$ such that $h^0(C, V \otimes M) = r+1$, and both $V \otimes M$ and $K_C \otimes M^{-1} \otimes V^*$ are globally generated and Petri trace injective.

The proof of this theorem has several ingredients. We begin by constructing a stable bundle E_0 with some of the properties we are interested in. Let F be a semistable bundle of rank $r-1$ and degree $(r-1)(g-1)-1$, and let N be a line bundle of degree g .

LEMMA 2.6. A general extension $0 \rightarrow F \rightarrow E \rightarrow N \rightarrow 0$ is a stable vector bundle.

Proof. Any subbundle G of E fits into an exact diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & G_1 & \longrightarrow & G & \longrightarrow & N(-D) \longrightarrow 0 \\ & & \downarrow \iota_1 & & \downarrow & & \downarrow \iota_2 \\ 0 & \longrightarrow & F & \longrightarrow & E & \longrightarrow & N \longrightarrow 0 \end{array}$$

where D is an effective divisor on C . If $\iota_2 = 0$, then $\mu(G) = \mu(G_1) \leq \mu(F) < \mu(E)$. Suppose $\iota_2 \neq 0$, and write $s := \text{rk}(G_1)$. If $s \neq 0$, the semistability of F implies that

$$\text{deg}(G_1) \leq s(g - 1) - \frac{s}{r - 1},$$

so in fact $\text{deg}(G_1) \leq s(g - 1) - 1$. As $\text{deg}(N) = g$, we have $\text{deg}(G) \leq (s + 1)(g - 1)$. Thus we need only exclude the case where $\text{deg}(G_1) = s(g - 1) - 1$ and $D = 0$, so $\iota_2 = \text{Id}_N$. In this case, the existence of the above diagram is equivalent to $[E] = (\iota_1)_*[G]$ for some extension G , that is, $[E] \in \text{Im}((\iota_1)_*)$.

It therefore suffices to check that

$$(\iota_1)_*: H^1(C, \text{Hom}(N, G_1)) \rightarrow H^1(C, \text{Hom}(N, F))$$

is not surjective. This follows from the fact, easily shown by a Riemann–Roch calculation, that $h^1(C, \text{Hom}(N, F/G_1)) > 0$.

If $s = 0$, then we need to exclude the lifting of $G = N(-p)$ for all $p \in C$, that is,

$$[E] \notin \bigcup_{p \in C} (\text{Ker}(H^1(C, \text{Hom}(N, F)) \rightarrow H^1(C, \text{Hom}(N(-p), F))))).$$

A dimension count shows that this locus is not dense in $H^1(C, \text{Hom}(N, F))$. □

LEMMA 2.7. *Suppose $h^0(C, N) \geq h^1(C, F)$. Then for a general extension $0 \rightarrow F \rightarrow E \rightarrow N \rightarrow 0$, the coboundary map is surjective.*

Proof. Clearly it suffices to exhibit one extension E_0 with the required property. We write $n := h^1(C, F)$ for brevity.

Let $0 \rightarrow F \rightarrow \tilde{F} \rightarrow \tau \rightarrow 0$ be an elementary transformation with $\text{deg}(\tau) = n$ and such that the image of $\Gamma(C, \tau)$ generates $H^1(C, F)$. We may assume that τ is supported along n general points p_1, \dots, p_n of C which are not base points of $|N|$. Then τ_{p_i} is generated by an element

$$\phi_i \in \left(\frac{F(p_i)}{F} \right)_{p_i}$$

defined up to nonzero scalar multiple. We write $[\phi_i]$ for the class in $H^1(C, F)$ defined by ϕ_i .

Now $h^0(C, N) \geq n$ and the image of C is nondegenerate in $|N|^*$. As the p_i can be assumed to be general, they impose independent conditions on sections of N . We choose sections $s_1, \dots, s_n \in H^0(C, N)$ such that $s_i(p_i) \neq 0$ but $s_i(p_j) = 0$ for $j \neq i$. For $1 \leq i \leq n$, let η_i be a local section of N^{-1} such that $\eta_i(s_i(p_i)) = 1$.

Let $0 \rightarrow F \rightarrow E_0 \rightarrow N \rightarrow 0$ be the extension with class $[E_0]$ defined by the image of

$$(\eta_1 \otimes \phi_1, \dots, \eta_n \otimes \phi_n)$$

by the coboundary map $\Gamma(C, N^{-1} \otimes \tau) \rightarrow H^1(C, N^{-1} \otimes F)$. Then $s_i \cup [E_0] = [\phi_i]$ for $1 \leq i \leq n$. Hence the image of $\cdot \cup [E_0]$ spans $H^1(C, F)$. □

We now make further assumptions on F and N . If $r = 2$, then $g \geq (2r + 2)(2r + 1) = 30$. Hence by the Brill–Noether theory of line bundles on C , we may choose a line bundle F of degree $g - 2$ with $h^0(C, F) = 2$ and $|F|$ base point free. If $r \geq 3$: since $g \geq 3$, we have $(r - 1)(g - 1) - 1 \geq r$. Therefore, by [BBN15, Theorem 5.1] we may choose a semistable bundle F of rank $r - 1$ and degree $(r - 1)(g - 1) - 1$ which is globally generated and satisfies $h^0(C, F) = r$, so $h^1(C, F) = r + 1$.

Furthermore, again by Brill–Noether theory, since $g \geq (2r + 2)(2r + 1)$ we may choose $N \in \text{Pic}^g(C)$ such that $h^0(C, N) = 2r + 2$ and $|N|$ is base point free. By Lemma 2.7, we may choose an $(r + 1)$ -dimensional subspace $\Pi \subset H^0(C, E)$ lifting from $H^0(C, N)$.

PROPOSITION 2.8. *Let F, N and Π be as above, and let $0 \rightarrow F \rightarrow E \rightarrow N \rightarrow 0$ be a general extension. Then the restricted Petri trace map $\Pi \otimes H^0(C, K_C \otimes E^*) \rightarrow H^0(C, K_C)$ is injective.*

Proof. Choose a basis $\sigma_1, \dots, \sigma_{r+1}$ for Π . For each i , write $\tilde{\sigma}_i$ for the image of σ_i in $H^0(C, N)$.

By Lemma 2.7, there is an isomorphism $H^1(C, E) \xrightarrow{\sim} H^1(C, N)$. Hence, by Serre duality, the injection $K_C \otimes N^{-1} \hookrightarrow K_C \otimes E^*$ induces an isomorphism on global sections. Choose a basis $\tau_1, \dots, \tau_{2r+1}$ for $H^0(C, K_C \otimes E^*)$. For each j , write $\tilde{\tau}_j$ for the preimage of τ_j by the aforementioned isomorphism.

For each i and j we have a commutative diagram

$$\begin{array}{ccccc}
 E^* & \xrightarrow{t\sigma_i} & \mathcal{O}_C & \xrightarrow{\tau_j} & K_C \otimes E^* \\
 \uparrow & \nearrow \tilde{\sigma}_i & & \searrow \tilde{\tau}_j & \uparrow \\
 N^{-1} & & & & K_C \otimes N^{-1}
 \end{array}$$

where the top row defines the twisted endomorphism

$$\mu(\sigma_i \otimes \tau_j) \in H^0(C, K_C \otimes \text{End } E^*) = H^0(C, K_C \otimes \text{End } E).$$

Clearly this has rank one. As it factorises via $K_C \otimes N^{-1}$, at a general point of C the eigenspace corresponding to the single nonzero eigenvalue is identified with the fibre of N^{-1} in E^* . Hence the Petri trace $\bar{\mu}(\sigma_i \otimes \tau_j)$ may be identified with the restriction to N^{-1} . By the diagram, we may identify this restriction with

$$\mu_N(\tilde{\sigma}_i \otimes \tilde{\tau}_j) \in H^0(C, K_C),$$

where μ_N is the Petri map of the line bundle N . Since C is Petri, μ_N is injective. Hence the elements $\bar{\mu}(\sigma_i \otimes \tau_j) = \mu_N(\tilde{\sigma}_i \otimes \tilde{\tau}_j)$ are independent in $H^0(C, K_C)$. This proves the statement. \square

Before proceeding, we need to recall some background on coherent systems (see [BBN08, §2] for an overview and [BGMN03] for more detail). We recall that a *coherent system of type (r, d, k)* is a pair (W, Π) where W is a vector bundle of rank r and degree d over C , and $\Pi \subseteq H^0(C, W)$ is a subspace of dimension k . There is a stability condition for coherent systems depending on a real parameter α , and a moduli space $G(\alpha; r, d, k)$ for equivalence classes of α -semistable coherent systems of type (r, d, k) . If $k \geq r$, then by [BGMN03, Proposition 4.6] there exists $\alpha_L \in \mathbb{R}$ such that $G(\alpha; r, d, k)$ is independent of α for $\alpha > \alpha_L$. This ‘terminal’ moduli space is denoted G_L . Moreover, the locus

$$U(r, d, k) := \{(W, \Pi) \in G_L : W \text{ is a stable vector bundle}\}$$

is an open subset of G_L . For us, $d = r(g - 1)$ and $k = r + 1$. To ease notation, we write $U := U(r, r(g - 1), r + 1)$.

Now let N_1 be a line bundle of degree g with $h^0(C, N_1) \geq r + 2$. Let F be as above, and let $0 \rightarrow F \rightarrow E \rightarrow N_1 \rightarrow 0$ be a general extension.

LEMMA 2.9. *For a general subspace $\Pi \subset H^0(C, E)$ of dimension $r + 1$, the coherent system (E, Π) defines an element of U .*

Proof. Recall the bundle E_0 defined in Lemma 2.7, which clearly is generically generated. Let us describe the subsheaf E'_0 generated by $H^0(C, E_0)$.

Write $p_1 + \dots + p_{r+1} =: D$. Clearly $s \cup [E_0] = 0$ for any $s \in H^0(C, N_1(-D))$. Since the p_i are general points,

$$h^0(C, N_1(-D)) = h^0(C, N_1) - (r + 1) = \dim(\text{Ker}(\cdot \cup [E_0]: H^0(C, N_1) \rightarrow H^1(C, F))).$$

Therefore, the image of $H^0(C, E_0)$ in $H^0(C, N_1)$ is exactly $H^0(C, N_1(-D))$. As the subbundle $F \subset E_0$ is globally generated, E'_0 is an extension $0 \rightarrow F \rightarrow E'_0 \rightarrow N_1(-D) \rightarrow 0$. Dualising and taking global sections, we obtain

$$0 \rightarrow H^0(C, N_1^{-1}(D)) \rightarrow H^0(C, (E'_0)^*) \rightarrow H^0(C, F^*) \rightarrow \dots$$

Since both $N_1^{-1}(D)$ and F^* are semistable of negative degree, $h^0(C, N_1^{-1}(-D)) = h^0(C, F^*) = 0$, so $h^0(C, (E'_0)^*) = 0$.


Now let $\Pi_1 \subset H^0(C, E_0)$ be any subspace of dimension $r + 1$ generically generating E_0 . Since $h^0(C, (E'_0)^*) = 0$, by [BBN08, Theorem 3.1(3)] the coherent system (E_0, Π_1) defines a point of G_L . Since generic generatedness and vanishing of $h^0(C, (E')^*)$ are open conditions on families of bundles with a fixed number of sections, the same is true for a generic (E, Π) where E is an extension $0 \rightarrow F \rightarrow E \rightarrow N_1 \rightarrow 0$. By Lemma 2.6, in fact (E, Π) belongs to U . \square

LEMMA 2.10. *For generic E represented in U , we have $h^0(C, E) = h^0(C, K_C \otimes E^*) = r + 1$.*

Proof. Since C is Petri general, $B_{1,g}^{r+2}$ is irreducible in $\text{Pic}^g(C)$. Thus there exists an irreducible family parametrising extensions of the form $0 \rightarrow F \rightarrow E \rightarrow N_1 \rightarrow 0$ where F is as above and N_1 ranges over $B_{1,g}^{r+2}$. This contains the extension E_0 constructed above. By Lemma 2.7, a general element E_1 of the family satisfies $h^0(C, E_1) = r + 1$. By semicontinuity, the same is true for general E represented in U . \square

Now by [BBN08, Theorem 3.1(4) and Remark 6.2], the locus U is irreducible. Write B for the component of $B_{r,r(g-1)}^{r+1}$ containing the image of U , and B' for the sublocus $\{E \in B : h^0(C, E) = r + 1\}$. Set $U' := U \times_B B'$; clearly $U' \cong B'$.

Let $\tilde{B} \rightarrow B$ be an étale cover such that there is a Poincaré bundle $\mathcal{E} \rightarrow \tilde{B} \times C$. In a natural way we obtain a commutative cube



where all faces are fibre product diagrams. By a standard construction, we can find a complex of bundles $\alpha: K^0 \rightarrow K^1$ over \tilde{B} such that $\text{Ker}(\alpha_b) \cong H^0(C, K_C \otimes \mathcal{E}_b^*)$ for each $b \in \tilde{B}$. Following [ACGH85, ch. IV], we consider the Grassmann bundle $\text{Gr}(r + 1, K^0)$ over \tilde{B} and the sublocus

$$\mathcal{G} := \{\Lambda \in \text{Gr}(r + 1, K^0) : \alpha|_{\Lambda} = 0\}.$$

Write $\mathcal{G}_1 := \mathcal{G} \times_{\tilde{B}} \tilde{U}$. The fibre of \mathcal{G}_1 over $(\mathcal{E}_b, \Pi) \in \tilde{U}$ is then $\text{Gr}(r + 1, H^0(C, K_C \otimes \mathcal{E}_b^*))$.

Now let E_0 be a bundle as constructed in Lemma 2.7 with $h^0(C, E_0) = 2r + 2$, and let Π_0 be a generic choice of $(r + 1)$ -dimensional subspace of $H^0(C, E_0)$. We may assume \tilde{U} is irreducible since U is. Since $\tilde{U} \rightarrow U$ is étale, by Lemma 2.8 in fact U is also smooth at (E_0, Π_0) (cf. [BGMN03, Proposition 3.10]). Therefore, we may choose a one-parameter family $\{(E_t, \Pi_t) : t \in T\}$ in \tilde{U} such that $(E_{t_0}, \Pi_{t_0}) = (E_0, \Pi_0)$ while (E_t, Π_t) belongs to \tilde{U}' for generic $t \in T$. Since the bundles have Euler characteristic zero, for generic $t \in T$ there is exactly one choice of $\Lambda \in \mathcal{G}_1|_{(E_t, \Pi_t)}$. Thus we obtain a section $T \setminus \{0\} \rightarrow \mathcal{G}_1$. As $\dim T = 1$, this section can be extended uniquely to 0. We thus obtain a triple (E_0, Π_0, Λ_0) where $\Lambda \subset H^0(C, K_C \otimes E_0^*)$ has dimension $r + 1$. By Lemma 2.8, this triple is Petri trace injective. Hence

$$(E_t, \Pi_t, \Lambda_t) = (E, H^0(C, E), H^0(C, K_C \otimes E^*))$$

is Petri trace injective for generic $t \in T$. Thus a general bundle E represented in \tilde{U}' is Petri trace injective.

Furthermore, by [BBN08, Theorem 3.1(4)], a general $(E, \Pi) \in U$ is globally generated (not just generically). Thus we obtain the following proposition.

PROPOSITION 2.11. *A general element E of the irreducible component $B \subseteq B_{r, r(g-1)}^{r+1}$ is Petri trace injective and globally generated with $h^0(C, E) = r + 1$.*

We can now prove the theorem.

Proof of Theorem 2.5. Consider the map $a: SU_C(r) \times \text{Pic}^{g-1}(C) \rightarrow U_C(r, r(g - 1))$ given by $(V, M) \mapsto V \otimes M$. This is an étale cover of degree r^{2g} . We write \bar{B} for the inverse image $a^{-1}(B)$. Since a is étale, we have $T_{(V, M)}\bar{B} \cong T_{V \otimes M}B$ for each $(V, M) \in B$. In particular,

$$\dim \bar{B} = \dim B = \dim U_C(r, r(g - 1)) - (r + 1)^2. \tag{6}$$

We write p for the projection $SU_C(r) \times \text{Pic}^{g-1}(C) \rightarrow SU_C(r)$, and p_1 for the restriction $p|_{\bar{B}}: \bar{B} \rightarrow SU_C(r)$.

CLAIM. p_1 is dominant.

To see this: for $(V, M) \in \bar{B}$, the locus $p_1^{-1}(V)$ is identified with an open subset of the twisted Brill–Noether locus

$$B_{1, g-1}^{r+1}(V) = \{M \in \text{Pic}^{g-1}(C) : h^0(C, V \otimes M) \geq r + 1\} \subseteq \text{Pic}^{g-1}(C).$$

Moreover, for each such (V, M) , we have

$$\dim_M(p_1^{-1}(V)) = \dim(T_M(B_{1, g-1}^{r+1}(V))) = \dim \text{Im}(\bar{\mu})^\perp.$$

Since $V \otimes M$ is Petri trace injective, this dimension is $g - (r + 1)^2$. By semicontinuity, a general fibre of p_1 has dimension at most $g - (r + 1)^2$. Therefore, in view of (6), the image of p_1 has dimension at least

$$(\dim U_C(r, r(g - 1)) - (r + 1)^2) - (g - (r + 1)^2) = \dim U_C(r, r(g - 1)) - g = \dim SU_C(r).$$

As $SU_C(r)$ is irreducible, the claim follows.

We can now conclude the proof. Let $V \in SU_C(r)$ be general. By the claim, we can find $(V, M) \in \bar{B}$ such that $h^0(C, V \otimes M) = r + 1$ and $V \otimes M$ is globally generated and Petri trace injective. By Proposition 2.4, the theta divisor Θ_V exists and satisfies $\text{mult}_M \Theta_V = h^0(C, V \otimes M) = r + 1$. Finally, by considering a suitable sum of line bundles, we see that the involution $E \mapsto K_C \otimes E^*$ preserves the component \bar{B} . Since a general element of \bar{B} is globally generated, in general both $V \otimes M$ and $K_C \otimes M^{-1} \otimes V^*$ are globally generated. \square

3. Reconstruction of bundles from tangent cones to theta divisors

3.1 Tangent cones

Let Y be a normal variety and $Z \subset Y$ a divisor. Let p be a smooth point of Y which is a point of multiplicity $n \geq 1$ of Z . A local equation f for Z near p has the form $f_n + f_{n+1} + \dots$, where the f_i are homogeneous polynomials of degree i in local coordinates centred at p . The projectivised tangent cone $\mathcal{T}_p(Z)$ to Z at p is the hypersurface in $\mathbb{P}T_p Y$ defined by the first nonzero component f_n of f . (For a more intrinsic description, see [ACGH85, ch. II.1].)

Now let C be a curve of genus $g \geq (r + 1)^2$. Let E be a Petri trace injective bundle of rank r and degree $r(g - 1)$, with $h^0(C, E) = r + 1$. By Proposition 2.4 (with $h = g - 1$), the theta divisor Θ_E is defined and contains the origin \mathcal{O}_C of $\text{Pic}^0(C)$ with multiplicity $h^0(C, E) = r + 1$.

By [CT11, Theorem 3.4 and Remark 3.8] (see also Kempf [Kem73]), the tangent cone $\mathcal{T}_{\mathcal{O}_C}(\Theta_E)$ to Θ_E at \mathcal{O}_C is given by the determinant of an $(r + 1) \times (r + 1)$ matrix $\Lambda = (l_{ij})$ of linear forms l_{ij} on $H^1(C, \mathcal{O}_C)$, which is related to the multiplication map $\bar{\mu}$ as follows: in appropriate bases (s_i) and (t_j) of $H^0(C, E)$ and $H^0(C, K_C \otimes E^*)$ respectively, Λ is given by

$$(l_{ij}) = (\bar{\mu}(s_i \otimes t_j)).$$

Hence, via Serre duality, Λ coincides with the cup product map

$$\cup: H^0(C, E) \otimes H^1(C, \mathcal{O}_C) \rightarrow H^1(C, E).$$

Thus the matrix $\Lambda = (l_{ij})$ is a matrix of linear forms on the canonical space $|K_C|^*$.

In the following two subsections, we will show on the one hand that one can recover the bundle E from the determinantal representation of the tangent cone $\mathcal{T}_{\mathcal{O}_C}(\Theta_E)$ given by the matrix Λ . On the other hand, up to changing bases in $H^0(C, E)$ and $H^1(C, E)$ there are only two determinantal representations of the tangent cone, namely Λ and Λ^t . Thus the tangent cone determines E up to an involution.

We will denote by $\mathbb{P} = |K_C|^*$ the canonical space and by φ the canonical embedding $C \hookrightarrow \mathbb{P}$.

3.2 Reconstruction of the bundle from the tangent cone

As above, let $\Lambda = (l_{ij})$ be the determinantal representation of the tangent cone given by the cup product mapping. We identify the source of Λ with $H^0(C, E)$ and the target with $H^1(C, E)$:

$$H^0(C, E) \otimes \mathcal{O}_{\mathbb{P}}(-1) \xrightarrow{\Lambda} H^1(C, E) \otimes \mathcal{O}_{\mathbb{P}}.$$

We recall that the Serre duality isomorphism sends $b \in H^1(C, E)$ to the linear form

$$\cdot \cup b: H^0(C, K_C \otimes E^*) \rightarrow H^1(C, K_C) = \mathbb{C}.$$

In the following proofs, we will use principal parts in order to represent cohomology classes of certain bundles. We refer to [Kem83] or [Pau03, § 3.2] for the necessary background. See also Kempf and Schreyer [KS88].

LEMMA 3.1. *Suppose that $h^0(C, E) = r + 1$ and E and $K_C \otimes E^*$ are globally generated. Then the rank of $\Lambda|_C = \varphi^* \Lambda$ is equal to $r = \text{rk } E$ at all points of C . In particular, the canonical curve is contained in $\mathcal{T}_{\mathcal{O}_C}(\Theta_E)$.*

Proof. For each $p \in C$, write β_p for a principal part with a simple pole supported at p . Then (see [KS88]) the cohomology class $[\beta_p]$ is identified with the image of p by φ . Therefore, at $p \in C$, the pullback $\Lambda|_C$ is identified with the cup product map

$$[\beta_p] \otimes s \mapsto [\beta_p] \cup s.$$

The kernel of $[\beta_p] \cup \cdot$ contains the subspace $H^0(C, E(-p))$, which is one-dimensional since E is globally generated and $h^0(C, E) = r + 1$. If $\text{Ker}([\beta_p] \cup \cdot)$ has dimension greater than 1, then there is a section $s' \in H^0(C, E)$ not vanishing at p such that

$$[\beta_p \cdot s'] = [\beta_p] \cup s' = 0 \in H^1(C, E).$$

By Serre duality, this means that

$$[\beta_p \cdot \langle s'(p), t(p) \rangle]$$

is zero in $H^1(C, K_C)$ for all $t \in H^0(C, K_C \otimes E^*)$. Hence the values at p of all global sections of $K_C \otimes E^*$ belong to the hyperplane in $(K_C \otimes E^*)|_p$ defined by contraction with the nonzero vector $s'(p) \in E|_p$. Thus $K_C \otimes E^*$ is not globally generated, contrary to our hypothesis. \square

Remark 3.2. Casalaina-Martin and Teixidor i Bigas in [CT11, §6] prove more generally that if E is a general vector bundle with $h^0(C, E) > kr$, then the k th secant variety of the canonical image $\varphi(C)$ of C is contained in $\mathcal{T}_{\mathcal{O}_C}(\Theta_E)$.

PROPOSITION 3.3. *Let E be a vector bundle with $h^0(C, E) = r + 1$, such that both E and $K_C \otimes E^*$ are globally generated. Then the image of $\Lambda|_C$ is isomorphic to $T_C \otimes E$.*

Proof. As $\varphi^* \mathcal{O}_{\mathbb{P}^{g-1}}(-1) \cong T_C$, the pullback $\varphi^* \Lambda = \Lambda|_C$ is a map

$$\Lambda|_C: T_C \otimes H^0(C, E) \rightarrow \mathcal{O}_C \otimes H^1(C, E).$$

Write $L := \det(E)$, a line bundle of degree $r(g - 1)$. Then $\det(K_C \otimes E^*) = K_C^r \otimes L^{-1}$. As $K_C \otimes E^*$ is globally generated, the evaluation sequence

$$0 \rightarrow K_C^{-r} \otimes L \rightarrow \mathcal{O}_C \otimes H^0(C, K_C \otimes E^*) \rightarrow K_C \otimes E^* \rightarrow 0$$

is exact. For each $p \in C$, the image of $(K_C^{-r} \otimes L)|_p$ in $H^0(C, K_C \otimes E^*)$ is exactly $\mathbb{C} \cdot t_p$, where t_p is the unique section, up to scalar, of $K_C \otimes E^*$ vanishing at p .

Dualising, we obtain a diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & T_C \otimes E & \longrightarrow & \mathcal{O}_C \otimes H^0(C, K_C \otimes E^*)^* & \xrightarrow{e} & K_C^r \otimes L^{-1} \longrightarrow 0 \\ & & & \nearrow & \uparrow \text{Serre} & & \\ & & T_C \otimes H^0(E) & \xrightarrow{\Lambda|_C} & \mathcal{O}_C \otimes H^1(C, E) & & \end{array}$$

Here e_p can be identified up to scalar with the map $f \mapsto f(t_p)$ where t_p is as above.

Now for each $p \in C$, the image

$$[\beta_p] \cup H^0(C, E) \subset H^1(C, E) \cong H^0(C, K_C \otimes E^*)^*$$

annihilates $t_p \in H^0(C, K_C \otimes E^*)$, since the principal part $\beta_p \cdot t_p$ is everywhere regular. Therefore, $\Lambda|_C$ factorises via $\text{Ker}(e) = T_C \otimes E$. Since $\text{rk}(\Lambda|_C) \equiv r$ by Lemma 3.1, we have $\text{Im}(\Lambda|_C) \cong T_C \otimes E$. \square

Remark 3.4. A straightforward computation shows also that

$$\text{Ker}(\Lambda|_C) \cong T_C \otimes L^{-1} \quad \text{and} \quad \text{Coker}(\Lambda|_C) \cong K_C^r \otimes L^{-1}.$$

We will also want to study the transpose Λ^t , which we will consider as a map

$$\Lambda^t: \mathcal{O}_{\mathbb{P}}(-1) \otimes H^0(C, K_C \otimes E^*) \rightarrow \mathcal{O}_{\mathbb{P}} \otimes H^1(C, K_C \otimes E^*).$$

The proof of Proposition 3.3 also shows the following corollary.

COROLLARY 3.5. *Let E and Λ be as above. Then the image of $\Lambda^t|_C$ is isomorphic to E^* .*

Remark 3.6. In order to describe the cokernel of $\Lambda|_C$, it is also enough to know at which points of C a row of $\Lambda|_C$ vanishes. Dualising the sequence

$$0 \rightarrow K_C^r \otimes L^{-1} \rightarrow \mathcal{O}_C \otimes H^0(C, K_C \otimes E^*) \rightarrow K_C \otimes E^* \rightarrow 0,$$

we see that $H^0(C, K_C \otimes E^*)^*$ is canonically identified with a subspace of $H^0(C, K_C^r \otimes L^{-1})$. Using the description of

$$T_C \otimes H^0(C, E) \xrightarrow{\Lambda|_C} \mathcal{O}_C \otimes H^1(C, E) \xrightarrow{\sim} \mathcal{O}_C \otimes H^0(C, K_C \otimes E^*)^*$$

as in the above proof, we see that a row vanishes exactly in a divisor associated to $K_C^r \otimes L^{-1}$. Hence, the cokernel is isomorphic to $K_C^r \otimes L^{-1}$.

3.3 Uniqueness of the linear determinantal representation of the tangent cone

In order to show the desired uniqueness of the determinantal representation of the tangent cone, we use a classical result of Frobenius. See [Fro97] and also for a modern proof [Die49, Wat87] and the references therein. For the sake of completeness we will give a sketch of a proof following Frobenius.

PROPOSITION 3.7. *Suppose $r \geq 1$. Let A and B be $(r+1) \times (r+1)$ matrices of independent linear forms, such that the entries of A are linear combinations of the entries of B and $\det(A) = c \cdot \det(B)$ for a nonzero constant $k \in \mathbb{C}$. Then, there exist invertible matrices $S, T \in \text{Gl}(r+1, \mathbb{C})$, unique up to scalar, such that $A = S \cdot B \cdot T$ or $A = S \cdot B^t \cdot T$.*

Proof by Frobenius [Fro97, pp. 1011–1013]. Note that for $r \geq 1$ only one of the above cases can occur and the matrices S and T are unique up to scalar. Indeed, let $A = SBT = S'BT'$ and set $b_{ii} = 1$ and $b_{ij} = 0$ if $i \neq j$; then $ST = S'T'$. Set $U = T(T'^{-1}) = S(S'^{-1})$; thus $UB = BU$. Since U commutes with every matrix, we have $U = k \cdot E_r$ and hence $S' = k \cdot S$ and $T' = 1/k \cdot T$. Similarly, one can show that B^t is not equivalent to B . Note also that there is no relation between any minors of A or B .

For $l = 0, \dots, r$, let c_{ij}^l be the coefficient of b_{ll} in a_{ij} and let y be a new variable. We substitute b_{ll} with $b_{ll} + y$ in A and B and get new matrices, denoted by $(a_{ij} + y \cdot c_{ij}^l)$ and B^l , respectively. Since $\det B^l$ is linear in y , the coefficient of y^2 in $\det(a_{ij} + y \cdot c_{ij}^l) = \det B^l$ has to vanish. But the coefficient is the sum of products of 2×2 minors of (c_{ij}^l) and $(r-1) \times (r-1)$ minors of A . Since there are no relations between any minors of A , all 2×2 minors of (c_{ij}^l) vanish. Hence, (c_{ij}^l) has rank one for any l and we can write $c_{ij}^l = p_i^l q_j^l$ where p^l and q^l are column and row vectors, respectively.

Let $B_0 = B|_{\{b_{ij}=0, i \neq j\}}$ and $A_0 = A|_{\{b_{ij}=0, i \neq j\}}$. Then

$$A_0 = PB_0Q,$$

where $P = (p_i^l)_{0 \leq i, l \leq r}$ and $Q = (q_j^l)_{0 \leq l, j \leq r}$. Since $\det(A_0) = c \cdot \det(B_0) = c \cdot b_{00} \cdots b_{rr}$, we get $\det(P) \cdot \det(Q) = c$, hence P and Q are invertible.

Let $\widetilde{B} = P^{-1}AQ^{-1}$. By definition $\widetilde{B}|_{\{b_{ij}=0, i \neq j\}} = B_0$. Thus, the entries \widetilde{b}_{ij} for $i \neq j$ and $v_i = \widetilde{b}_{ii} - b_{ii}$ are linear functions in b_{ij} for $i \neq j$. Furthermore, we have

$$\det(\widetilde{B}) = \det(P^{-1}AQ^{-1}) = \det(P^{-1}Q^{-1}) \cdot \det(A) = \frac{1}{c} \cdot c \cdot \det(B) = \det(B).$$

Comparing the coefficients of $b_{11}b_{22} \cdots b_{rr}$ in $\det(\widetilde{B})$ and $\det(B)$, we get $v_0 = 0$. Similarly, $v_i = 0$ for $0 \leq i \leq r$. Comparing the coefficients of $b_{22} \cdots b_{rr}$, we get $b_{12}b_{21} = \widetilde{b}_{12}\widetilde{b}_{21}$ and in general

$$b_{ij}b_{ji} = \widetilde{b}_{ij}\widetilde{b}_{ji}, \quad i \neq j.$$

Comparing the coefficients of $b_{33} \cdots b_{rr}$, we get $b_{12}b_{23}b_{31} + b_{21}b_{13}b_{32} = \widetilde{b}_{12}\widetilde{b}_{23}\widetilde{b}_{31} + \widetilde{b}_{21}\widetilde{b}_{13}\widetilde{b}_{32}$ and in general

$$b_{ij}b_{jk}b_{ki} + b_{ji}b_{ik}b_{kj} = \widetilde{b}_{ij}\widetilde{b}_{jk}\widetilde{b}_{ki} + \widetilde{b}_{ji}\widetilde{b}_{ik}\widetilde{b}_{kj}, \quad i \neq j \neq k \neq i.$$

A careful study of these equations shows that either

$$\widetilde{b}_{ij} = \frac{k_i}{k_j} b_{ij} \quad \text{and} \quad \widetilde{B} = KBK^{-1} \quad \text{or} \quad \widetilde{b}_{ij} = \frac{k_i}{k_j} b_{ji} \quad \text{and} \quad \widetilde{B} = KB^tK^{-1},$$

where $K = (k_i \delta_{ij})_{0 \leq i, j \leq r}$. The claim follows. □

We now assume that E is a Petri trace injective bundle. Let $\Lambda = (l_{ij})$ be a determinantal representation of the tangent cone $\mathcal{T}_{\mathcal{O}_C}(\Theta_E)$ as above. By Petri trace injectivity, the matrix Λ is $(r + 1)$ -generic (see [Eis88] for a definition), that is, there are no relations between the entries l_{ij} or any subminors of Λ .

COROLLARY 3.8. *For a curve of genus $g \geq (r + 1)^2$ and a Petri trace injective bundle E with $r + 1$ global sections of degree $r(g - 1)$, any determinantal representation of the tangent cone $\mathcal{T}_{\mathcal{O}_C}(\Theta_E) \subset |K_C|^*$ is equivalent to Λ or Λ^t .*

Proof. Let α be any determinantal representation of the tangent cone $\mathcal{T}_{\mathcal{O}_C}(\Theta_E)$ in $|K_C|^*$. Then, α is an $(r + 1) \times (r + 1)$ matrix of linear entries, since the degree of the tangent cone is $r + 1$. Furthermore, the entries α_{ij} of α are linear combinations of the entries l_{ij} of Λ . Indeed, assume for some k, l that α_{kl} is not a linear combination of the l_{ij} . Then $\mathcal{T}_{\mathcal{O}_C}(\Theta_E)$ would be the cone over $V(\alpha_{kl}) \cap \mathcal{T}_{\mathcal{O}_C}(\Theta_E)$. Hence, the vertex of $\mathcal{T}_{\mathcal{O}_C}(\Theta_E)$ defined by the entries l_{ij} would have codimension strictly less than $(r + 1)^2$, a contradiction to the independence of the l_{ij} . The corollary follows from Proposition 3.7. □

4. Injectivity of the theta map

THEOREM 4.1. *Suppose $r \geq 2$. Let C be a Petri general curve of genus $g \geq (2r + 2)(2r + 1)$. Then the theta map $\mathcal{D}: SU_C(r) \dashrightarrow |r\Theta|$ is generically injective.*

Proof. Let $V \in SU_C(r)$ be a general stable bundle. By Theorem 2.5, there exists $M \in \Theta_V$ such that $h^0(C, V \otimes M) = r + 1$, the bundle $V \otimes M =: E$ is Petri trace injective, and E and $K_C \otimes E^*$ are globally generated.

Note that tensoring by M^{-1} defines an isomorphism $\Theta_V \xrightarrow{\sim} \Theta_E$ inducing an isomorphism $\mathcal{T}_M(\Theta_V) \xrightarrow{\sim} \mathcal{T}_{\mathcal{O}_C}(\Theta_E)$. In order to use the results of the previous sections, we will work with Θ_E .

Now let

$$\alpha: \mathcal{O}_{\mathbb{P}^{g-1}}(-1) \otimes \mathbb{C}^{r+1} \rightarrow \mathcal{O}_{\mathbb{P}^{g-1}} \otimes \mathbb{C}^{r+1}$$

be a map of bundles of rank $r + 1$ over \mathbb{P}^{g-1} whose determinant defines the tangent cone $\mathcal{T}_{\mathcal{O}_C}(\Theta_E)$. By Corollary 3.8, the map α is equivalent either to Λ or Λ^t , where Λ is the representation given by the cup product mapping as defined in §3. Therefore, by Proposition 3.3 and Corollary 3.5, the image E' of $\alpha|_C$ is isomorphic either to $T_C \otimes E = V \otimes M \otimes T_C$ or to $E^* = V^* \otimes M^{-1}$. Thus V is isomorphic either to

$$E' \otimes K_C \otimes M^{-1} \quad \text{or to} \quad (E')^* \otimes M^{-1}. \tag{7}$$

Now since in particular $g > (r + 1)^2$, the open subset $\{M \in \text{Pic}^{g-1}(C) : h^0(C, V \otimes M) = r + 1\} \subseteq B_{1,g-1}^{r+1}(V)$ has a component of dimension $g - (r + 1)^2 \geq 1$. Therefore, we may assume that $M^{2r} \not\cong K_C^r$. Thus only one of the bundles in (7) can have trivial determinant. Hence there is only one possibility for V .

In summary, the data of the tangent cone $\mathcal{T}_M(\Theta_V)$ and the point M , together with the property $\det(V) = \mathcal{O}_C$, determine the bundle V up to isomorphism. In particular, Θ_V determines V . □

Remark 4.2. The involution $M \mapsto K_C \otimes M^{-1}$ defines an isomorphism of varieties $\Theta_V \xrightarrow{\sim} \Theta_{V^*}$. We observe that the transposed map Λ^t occurs naturally as the cup product map defining the tangent cone $\mathcal{T}_{K_C \otimes M^{-1}}(\Theta_{V^*})$.

Remark 4.3. If C is hyperelliptic, then the canonical map factorises via the hyperelliptic involution ι . Thus the construction in Proposition 3.3 can never give bundles over C which are not ι -invariant. We note that Beauville [Bea88] showed that in rank two, if C is hyperelliptic then the bundles V and ι^*V have the same theta divisor.

ACKNOWLEDGEMENTS

The authors would like to thank Ciro Ciliberto, Peter Newstead, Christian Pauly, Frank-Olaf Schreyer and Edoardo Sernesi for helpful discussions and comments on this work.

REFERENCES

ACGH85 E. Arbarello, M. Cornalba, P. A. Griffiths and J. Harris, *Geometry of algebraic curves*, Vol. I, Grundlehren der Mathematischen Wissenschaften, vol. 267 [Fundamental Principles of Mathematical Sciences] (Springer, New York, 1985).

Bea88 A. Beauville, *Fibrés de rang 2 sur une courbe, fibré déterminant et fonctions thêta. (Rank 2 fiber bundles on a curve, determinant bundle and theta functions)*, Bull. Soc. Math. France **116** (1988), 431–448.

Bea06 A. Beauville, *Vector bundles on curves and theta functions*, in *Moduli spaces and arithmetic geometry. Papers of the 13th International Research Institute of the Mathematical Society of Japan, Kyoto, Japan, September 8–15, 2004* (Mathematical Society of Japan, Tokyo, 2006), 145–156.

- BBN08 U. N. Bhosle, L. Brambila-Paz and P. E. Newstead, *On coherent systems of type $(n, d, n + 1)$ on Petri curves*, Manuscripta Math. **126** (2008), 409–441.
- BBN15 U. N. Bhosle, L. Brambila-Paz and P. E. Newstead, *On linear series and a conjecture of D. C. Butler*, Internat. J. Math. **26** (2015), 1550007, 18.
- BGMN03 S. B. Bradlow, O. García-Prada, V. Muñoz and P. E. Newstead, *Coherent systems and Brill-Noether theory*, Internat. J. Math. **14** (2003), 683–733.
- BV12 S. Brivio and A. Verra, *Plücker forms and the theta map*, Amer. J. Math. **134** (2012), 1247–1273.
- BS11 A. Bruno and E. Sernesi, *A note on the Petri loci*, Manuscripta Math. **136** (2011), 439–443.
- CT11 S. Casalaina-Martin and M. Teixidor i Bigas, *Singularities of Brill-Noether loci for vector bundles on a curve*, Math. Nachr. **284** (2011), 1846–1871.
- CS92 C. Ciliberto and E. Sernesi, *Singularities of the theta divisor and congruences of planes*, J. Algebr. Geom. **1** (1992), 231–250.
- Die49 J. Dieudonné, *Sur une généralisation du groupe orthogonal à quatre variables*, Arch. Math. **1** (1949), 282–287.
- DN89 J.-M. Drezet and M. S. Narasimhan, *Groupe de Picard des variétés de modules de fibrés semi-stables sur les courbes algébriques*, Invent. Math. **97** (1989), 53–94.
- Eis88 D. Eisenbud, *Linear sections of determinantal varieties*, Amer. J. Math. **110** (1988), 541–575.
- Far05 G. Farkas, *Gaussian maps, Gieseker–Petri loci and large theta-characteristics*, J. Reine Angew. Math. **581** (2005), 151–173.
- Fro97 G. F. Frobenius, *Über die Darstellung der endlichen Gruppe durch lineare Substitutionen*, Sitzungsberichte der Königlich Preussischen Akademie der Wissenschaften zu Berlin **2** (1897), 994–1015.
- Gie82 D. Gieseker, *Stable curves and special divisors: Petri’s conjecture*, Invent. Math. **66** (1982), 251–275.
- Gre84 M. L. Green, *Quadrics of rank four in the ideal of a canonical curve*, Invent. Math. **75** (1984), 85–104.
- GT09 I. Grzegorczyk and M. Teixidor i Bigas, *Brill–Noether theory for stable vector bundles*, in *Moduli spaces and vector bundles*, London Mathematical Society Lecture Note Series, vol. 359 (Cambridge University Press, Cambridge, 2009), 29–50.
- Kem73 G. R. Kempf, *On the geometry of a theorem of Riemann*, Ann. of Math. (2) **98** (1973), 178–185.
- Kem83 G. R. Kempf, *Abelian integrals*, Monografías del Instituto de Matemáticas, vol. 13 [Monographs of the Institute of Mathematics] (Universidad Nacional Autónoma de México, México, 1983).
- KS88 G. R. Kempf and F.-O. Schreyer, *A Torelli theorem for osculating cones to the theta divisor*, Compos. Math. **67** (1988), 343–353.
- NR69 M. S. Narasimhan and S. Ramanan, *Moduli of vector bundles on a compact Riemann surface*, Ann. of Math. (2) **89** (1969), 14–51.
- Pau03 C. Pauly, *On cubics and quartics through a canonical curve*, Ann. Sc. Norm. Super. Pisa Cl. Sci. (5) **2** (2003), 803–822.
- Pau10 C. Pauly, *Rank four vector bundles without theta divisor over a curve of genus two*, Adv. Geom. **10** (2010), 647–657.
- Pop99 M. Popa, *On the base locus of the generalized theta divisor*, C. R. Acad. Sci., Paris, Sér. I, Math. **329** (1999), 507–512.
- Ray82 M. Raynaud, *Sections des fibrés vectoriels sur une courbe*, Bull. Soc. Math. France **110** (1982), 103–125.
- TiB88 M. Teixidor i Bigas, *The divisor of curves with a vanishing theta-null*, Compos. Math. **66** (1988), 15–22.

GENERIC INJECTIVITY OF THE THETA MAP

- TiB14 M. Teixidor i Bigas, *Injectivity of the Petri map for twisted Brill–Noether loci*, *Manuscripta Math.* **145** (2014), 389–397.
- vGI01 B. van Geemen and E. Izadi, *The tangent space to the moduli space of vector bundles on a curve and the singular locus of the theta divisor of the Jacobian*, *J. Algebraic Geom.* **10** (2001), 133–177.
- Wat87 W. C. Waterhouse, *Automorphisms of $\det(X_{ij})$: the group scheme approach*, *Adv. Math.* **65** (1987), 171–203.

George H. Hitching george.hitching@hioa.no
Høgskolen i Oslo og Akershus, Postboks 4, St. Olavs plass, 0130 Oslo, Norway

Michael Hoff hahn@math.uni-sb.de
Universität des Saarlandes, Campus E2 4, D-66123 Saarbrücken, Germany