

**A DECOMPOSITION THEOREM ON DIFFERENTIAL  
 POLYNOMIALS OF THETA FUNCTIONS**

HISASI MORIKAWA

Let  $\tau = (\tau_{ij})$  be a symmetric complex  $g \times g$  matrix with the positive definite imaginary part. A theta function of level  $n$  means an entire function  $f(z)$  in  $g$  complex variables  $z = (z_1, \dots, z_g)$  satisfying the difference relations:

$$f(z + \hat{b} + b\tau) = \exp(-\pi n \sqrt{-1}(b\tau^t b + 2z^t b))f(z), \quad ((\hat{b}, b) \in \mathbf{Z}^g \times \mathbf{Z}^g).$$

Denoting by  $\Theta_0^{(n)}$  the vector space of theta functions of level  $n$ , we get the graded algebra of theta functions;

$$\Theta_0 = \sum_{n \geq 1} \Theta_0^{(n)}.$$

Theta series

$$\mathcal{G}^{(n)} \left[ \begin{matrix} a/n \\ 0 \end{matrix} \right] (\tau | z) = \sum_{\ell \in \mathbf{Z}^g} \exp\left(\pi n \sqrt{-1} \left( \left( \ell + \frac{a}{n} \right) \tau^t \left( \ell + \frac{a}{n} \right) + 2z^t \left( \ell + \frac{a}{n} \right) \right)\right),$$

$(a \in \mathbf{Z}^g / n\mathbf{Z}^g)$

form a canonical basis of  $\Theta_0^{(n)}$ , and thus

$$\dim \Theta_0^{(n)} = n^g.$$

In the present article we shall prove the following decomposition theorem:

The algebra of differential polynomials of theta functions has a canonical linear basis

$$\left\{ \left( \frac{\partial}{\partial z} \right)^j \mathcal{G}^{(n)} \left[ \begin{matrix} a/n \\ 0 \end{matrix} \right] (\tau | z) \mid j \in \mathbf{Z}_{\geq 0}^g, a \in \mathbf{Z}^g / n\mathbf{Z}^g, n \geq 1 \right\},$$

i.e. any differential polynomial is uniquely expressed as a linear combination of  $(\partial/\partial z)^j \mathcal{G}^{(n)} \left[ \begin{matrix} a/n \\ 0 \end{matrix} \right] (\tau | z)$ , ( $j \in \mathbf{Z}_{\geq 0}^g$ ,  $a \in \mathbf{Z}^g / n\mathbf{Z}^g$ ,  $n \geq 1$ ) with constant

---

Received November 14, 1983.

coefficients depending on  $\tau$ . More precisely we have the explicit expressions of the components of the decomposition.

The key is a very similar idea as making transvectants in the classical invariant theory, however the Lie algebra is Heisenberg Lie algebra instead of  $sl_2$ . The algebra  $\Theta_0$  of theta functions is embedded in a graded algebra  $\Theta$  of auxiliary theta functions in  $2g$  complex variables  $(u, z) = (u_1, \dots, u_g, z_1, \dots, z_g)$  with the following properties,

1° A realization  $\langle \mathcal{L}, \mathcal{D}_1, \dots, \mathcal{D}_g, \Delta_1, \dots, \Delta_g \rangle$  of Heisenberg Lie algebra acts on  $\Theta$  as derivations,

2°  $\Theta_0$  is the subalgebra consisting of all the elements  $\varphi$  such that  $\mathcal{D}_i \varphi = 0$  ( $1 \leq i \leq g$ ),

3°  $\left\{ \Delta^j \mathcal{G}^{(n)} \begin{bmatrix} a/n \\ 0 \end{bmatrix} (\tau|z) \mid j \in \mathbb{Z}_{\geq 0}^g, a \in \mathbb{Z}^g/n\mathbb{Z}^g, n \geq 1 \right\}$  is a canonical linear basis of  $\Theta$ ,

4° The mapping

$$\Delta^j \mathcal{G}^{(n)} \begin{bmatrix} a/n \\ 0 \end{bmatrix} (\tau|z) \longrightarrow \left( \frac{\partial}{\partial z} \right)^j \mathcal{G}^{(n)} \begin{bmatrix} a/n \\ 0 \end{bmatrix} (\tau|z), \quad (j \in \mathbb{Z}_{\geq 0}^g, a \in \mathbb{Z}^g/n\mathbb{Z}^g, n \geq 1)$$

induces an algebra isomorphism of  $\Theta$  onto the algebra of differential polynomials of theta functions.

We shall also characterize differential polynomials of theta functions which are theta functions.

The associative law for the structure constants of

$$C \left[ \dots, \left( \frac{\partial}{\partial z} \right)^j \mathcal{G}^{(n)} \begin{bmatrix} a/n \\ 0 \end{bmatrix} (\tau|z), \dots \right]$$

with respect to the basis must be very important relations between

$$\left\{ \left( \frac{\partial}{\partial z} \right)^j \mathcal{G}^{(n)} \begin{bmatrix} a/n \\ 0 \end{bmatrix} (\tau| \frac{\hat{a}}{n}) \mid j \in \mathbb{Z}_{\geq 0}^g; a, \hat{a} \in \mathbb{Z}^g/n\mathbb{Z}^g; n \geq 1 \right\}.$$

*Notations.*

$$\mathbb{Z}_{\geq 0} = \{\text{non-negative integers}\}, \mathbb{Z}_{\geq 0}^g = \{j = (j_1, \dots, j_g) \mid j_i \in \mathbb{Z}_{\geq 0}\},$$

$$j \pm \varepsilon_i = (j_1, \dots, j_{i-1}, j_i \pm 1, j_{i+1}, \dots, j_g), j! = j_1! \cdots j_g!,$$

$$\binom{j}{p} = \binom{j_1}{p_1} \cdots \binom{j_g}{p_g}, \binom{j}{k^{(1)}, \dots, k^{(r)}} = \binom{j_1}{k_1^{(1)}, \dots, k_1^{(r)}} \cdots \binom{j_g}{k_g^{(1)}, \dots, k_g^{(r)}},$$

$$|j| = j_1 + \cdots + j_g, u = (u_1, \dots, u_g), z = (z_1, \dots, z_g), u^j = u_1^{j_1} \cdots u_g^{j_g},$$

$$z^j = z_1^{j_1} \cdots z_g^{j_g},$$

$$\left( \frac{\partial}{\partial u} \right)^j = \left( \frac{\partial}{\partial u_1} \right)^{j_1} \cdots \left( \frac{\partial}{\partial u_g} \right)^{j_g}, \left( \frac{\partial}{\partial z} \right)^j = \left( \frac{\partial}{\partial z_1} \right)^{j_1} \cdots \left( \frac{\partial}{\partial z_g} \right)^{j_g},$$

$$\left(2\pi n\sqrt{-1}u + \frac{\partial}{\partial u}\right)^j = \left(2\pi n\sqrt{-1}u_1 + \frac{\partial}{\partial z_1}\right)^{j_1} \cdots \left(2\pi n\sqrt{-1}u_g + \frac{\partial}{\partial z_g}\right)^{j_g}.$$

§1. Auxiliary theta functions

1.1. An auxiliary theta function of level  $n$  means a function  $\varphi(u, z)$  in  $2g$  complex variables  $(u, z) = (u_1, \dots, u_g, z_1, \dots, z_g)$  such that

1°  $\varphi(u, z)$  is a polynomial in  $u = (u_1, \dots, u_g)$  whose coefficients are entire functions in  $z = (z_1, \dots, z_g)$ ,

2°  $\varphi(u + b, z + \hat{b} + b\tau) = \exp(-\pi n\sqrt{-1}(b\tau^t b + 2z^t b))\varphi(u, z)$ ,  $((\hat{b}, b) \in Z^g \times Z^g)$ .

Denoting by  $\Theta^{(n)}$  the vector space of auxiliary theta functions of level  $n$ , we obtain a graded algebra

$$\Theta = \sum_{n \geq 1} \Theta^{(n)}$$

of auxiliary theta functions, which contains the graded algebra  $\Theta_0$  of theta functions as the subalgebra of polynomials of degree zero in  $u$ . Auxiliary theta series are also defined as follows,

$$\begin{aligned} & \mathcal{G}_j^{(n)} \left[ \begin{matrix} a/n \\ 0 \end{matrix} \right] (\tau | u, z) \\ (1.1) \quad &= (2\pi n\sqrt{-1})^{|j|} \sum_{\ell \in Z^g} \left(u + \ell + \frac{a}{n}\right)^j \\ & \cdot \exp \pi n\sqrt{-1} \left( \left(\ell + \frac{a}{n}\right)\tau^t \left(\ell + \frac{a}{n}\right) + 2z^t \left(\ell + \frac{a}{n}\right) \right) \\ & (j \in Z_{\geq 0}^g, a \in Z^g/nZ^g, n \geq 1). \end{aligned}$$

LEMMA 1.1.

$$(1.2) \quad \mathcal{G}_j^{(n)} \left[ \begin{matrix} a/n \\ 0 \end{matrix} \right] (\tau | u, z) = \left(2\pi n\sqrt{-1}u + \frac{\partial}{\partial z}\right)^j \mathcal{G}_j^{(n)} \left[ \begin{matrix} a/n \\ 0 \end{matrix} \right] (\tau | z),$$

$$\begin{aligned} & \mathcal{G}_j^{(n)} \left[ \begin{matrix} a/n \\ 0 \end{matrix} \right] (\tau | u + b, z + \hat{b} + b\tau) \\ (1.3) \quad &= \exp(-\pi n\sqrt{-1}(b\tau^t b + 2z^t b)) \mathcal{G}_j^{(n)} \left[ \begin{matrix} a/n \\ 0 \end{matrix} \right] (\tau | u, z) \\ & ((\hat{b}, b) \in Z^g \times Z^g). \end{aligned}$$

*Proof.* For  $a, b, \hat{b}$  in  $Z^g$  we have

$$\begin{aligned}
 & \left(2\pi n\sqrt{-1}u + \frac{\partial}{\partial z}\right)^j \exp\left(\pi n\sqrt{-1}\left(\left(\ell + \frac{a}{n}\right)\tau^t\left(\ell + \frac{a}{n}\right) + 2z^t\left(\ell + \frac{a}{n}\right)\right)\right) \\
 &= (2\pi n\sqrt{-1})^{|j|} \left(u + \ell + \frac{a}{n}\right)^j \\
 & \quad \exp\left(\pi n\sqrt{-1}\left(\left(\ell + \frac{a}{n}\right)\tau^t\left(\ell + \frac{a}{n}\right) + 2z\left(\ell + \frac{a}{n}\right)\right)\right), \\
 & \left(u + \ell + b + \frac{a}{n}\right)^j \\
 & \quad \cdot \exp\left(\pi n\sqrt{-1}\left(\left(\ell + b + \frac{a}{n}\right)\tau^t\left(\ell + b + \frac{a}{n}\right) + 2z^t\left(\ell + b + \frac{a}{n}\right)\right)\right) \\
 &= \exp\left(\pi n\sqrt{-1}(b\tau^t b + 2z^t b)\right) \left(u + \ell + b + \frac{a}{n}\right)^j \\
 & \quad \cdot \exp\left(\pi n\sqrt{-1}\left(\left(\ell + \frac{a}{n}\right)\tau^t\left(\ell + \frac{a}{n}\right)\tau^t\left(\ell + \frac{a}{n}\right) + 2(z + \hat{b} + b\tau)\left(\ell + \frac{a}{n}\right)\right)\right).
 \end{aligned}$$

Hence, making the sum with respect to  $\ell \in \mathbb{Z}^g$ , we obtain (1.2), (1.3).

**THEOREM 1.1.**  $\left\{\mathcal{G}_j^{(n)}\left[\begin{smallmatrix} a/n \\ 0 \end{smallmatrix}\right](\tau|u, z) \mid j \in \mathbb{Z}_{\geq 0}^g, a \in \mathbb{Z}^g/n\mathbb{Z}^g\right\}$  is a basis of the space  $\Theta^n$  of auxiliary theta functions of level  $n$ .

*Proof.* By virtue of Lemma 1.1  $\mathcal{G}_j^{(n)}\left[\begin{smallmatrix} a/n \\ 0 \end{smallmatrix}\right](\tau|u, z)$  ( $j \in \mathbb{Z}_{\geq 0}^g, a \in \mathbb{Z}^g/n\mathbb{Z}^g$ ) belong to  $\Theta^{(n)}$ , and obviously they are linearly independent. Let  $\varphi(u, z) = \sum_j u^j f_j(z)$  be an element of  $\Theta^{(n)}$ , and let  $u^k f_k(z)$  be one of terms with maximal degree  $k$  in  $u$ . Then, comparing the coefficients of  $u^k$  in the both sides of

$$\sum_j (u + b)^j f_j(z + \hat{b} + b\tau) = \exp(-\pi n\sqrt{-1}(b\tau^t b + 2z^t b)) \sum_j u^j f_j(z),$$

we have

$$f_k(z + \hat{b} + b\tau) = \exp(-\pi n\sqrt{-1}(b\tau^t b + 2z^t b)) f_k(z).$$

This means that there exists a system  $(\alpha_a)_{a \in \mathbb{Z}^g/n\mathbb{Z}^g}$  of constants such that

$$f_k(z) = \sum_a \alpha_a \mathcal{G}_a^{(n)}\left[\begin{smallmatrix} a/n \\ 0 \end{smallmatrix}\right](\tau|z),$$

and thus

$$\varphi(u, z) - \sum_a \alpha_a \mathcal{G}_k^{(n)} \left[ \begin{matrix} a/n \\ 0 \end{matrix} \right] (\tau | u, z)$$

is an element in  $\Theta^{(n)}$  without  $u^k$ -term and all the new terms are of lower degree than  $k$  in  $u$ . Proceeding this process successively, we can express  $\varphi(u, z)$  as a linear sum of  $\mathcal{G}_j^{(n)} \left[ \begin{matrix} a/n \\ 0 \end{matrix} \right] (\tau | u, z)$  ( $j \in \mathbb{Z}_{\geq 0}^g, a \in \mathbb{Z}^g/n\mathbb{Z}^g$ ).

1.2. Denoting the projection operators by

$$\sigma^{(n)}: \Theta \longrightarrow \Theta^{(n)}, \quad (n \geq 1)$$

we define differential operators

$$\begin{aligned} \mathcal{E} &= \sum_{n \geq 1} n \sigma^{(n)}, \\ \mathcal{D}_i &= \sum_{n \geq 1} \frac{1}{2\pi\sqrt{-1}} \frac{\partial}{\partial u_i} \circ \sigma^{(n)}, \\ \Delta_i &= \sum_{n \geq 1} \left( 2\pi n \sqrt{-1} u_i + \frac{\partial}{\partial z_i} \right) \circ \sigma^{(n)}, \\ \mathcal{D}^j &= \mathcal{D}_1^{j_1} \dots \mathcal{D}_g^{j_g}, \quad \Delta_1^{j_1} \dots \Delta_g^{j_g}. \end{aligned}$$

PROPOSITION 1.1.

$$(1.4) \quad \mathcal{D}_i \mathcal{G}_j^{(n)} \left[ \begin{matrix} a/n \\ 0 \end{matrix} \right] (\tau | u, z) = n j_i \mathcal{G}_{j-\varepsilon_i}^{(n)} \left[ \begin{matrix} a/n \\ 0 \end{matrix} \right] (\tau | u, z),$$

$$(1.5) \quad \Delta_i \mathcal{G}_j^{(n)} \left[ \begin{matrix} a/n \\ 0 \end{matrix} \right] (\tau | u, z) = \mathcal{G}_{j+\varepsilon_i}^{(n)} \left[ \begin{matrix} a/n \\ 0 \end{matrix} \right] (\tau | u, z),$$

$$(1.6) \quad \mathcal{G}_j^{(n)} \left[ \begin{matrix} a/n \\ 0 \end{matrix} \right] (\tau | u, z) = \Delta^j \mathcal{G}^{(n)} \left[ \begin{matrix} a/n \\ 0 \end{matrix} \right] (\tau | z),$$

$$(1.7) \quad \frac{1}{p!} \mathcal{D}^p \mathcal{G}_j^{(n)} \left[ \begin{matrix} a/n \\ 0 \end{matrix} \right] (\tau | u, z) = \binom{j}{p} n^{|p|} \mathcal{G}_{j-p}^{(n)} \left[ \begin{matrix} a/n \\ 0 \end{matrix} \right] (\tau | u, z),$$

$$(1.8) \quad \frac{1}{j!} \mathcal{D}^j \mathcal{G}_j^{(n)} \left[ \begin{matrix} a/n \\ 0 \end{matrix} \right] (\tau | u, z) = n^{|j|} \mathcal{G}^{(n)} \left[ \begin{matrix} a/n \\ 0 \end{matrix} \right] (\tau | z)$$

$(j, p \in \mathbb{Z}_{\geq 0}^g, j \geq p, a \in \mathbb{Z}^g/n\mathbb{Z}^g, n \geq 1).$

*Proof.* From the expression

$$\mathcal{G}_j^{(n)} \left[ \begin{matrix} a/n \\ 0 \end{matrix} \right] (\tau | u, z) = \left( 2\pi n \sqrt{-1} u + \frac{\partial}{\partial z} \right)^j \mathcal{G}^{(n)} \left[ \begin{matrix} a/n \\ 0 \end{matrix} \right] (\tau | z)$$

it follows (1.4), (1.5), (1.6). Applying (1.4) and (1.5) successively, we have (1.7), (1.8).

PROPOSITION 1.2.  $\mathcal{E}, \mathcal{D}_1, \dots, \mathcal{D}_g, \Delta_1, \dots, \Delta_g$  are derivations of  $\Theta$  such that

$$(1.9) \quad \begin{aligned} [\mathcal{E}, \mathcal{D}_i] &= [\mathcal{E}, \Delta_i] = [\mathcal{D}_i, \mathcal{D}_j] = [\Delta_i, \Delta_j] = 0, \\ [\mathcal{D}_i, \Delta_{i'}] &= \begin{cases} \mathcal{E} & (i = i') \\ 0 & (i \neq i') \end{cases} \quad (1 \leq i, i', j \leq g). \end{aligned}$$

*Proof.* By virtue of Proposition 1.2  $\mathcal{E}, \mathcal{D}_1, \dots, \mathcal{D}_g, \Delta_1, \dots, \Delta_g$ , map  $\Theta$  into itself. Since  $\Theta = \sum_{n \geq 1} \Theta^{(n)}$  is a graded algebra,  $\mathcal{E}, \mathcal{D}_1, \dots, \mathcal{D}_g, \Delta_1, \dots, \Delta_g$  are derivations of  $\Theta$ . By simple calculation we have (1.9).

Proposition 1.2 states  $\langle \mathcal{E}, \mathcal{D}_1, \dots, \mathcal{D}_g, \Delta_1, \dots, \Delta_g \rangle$  is a realization of Heisenberg Lie algebra acting on  $\Theta$  as derivations.

PROPOSITION 1.3. *The graded algebra of theta functions is the sub-algebra consisting of all the elements  $\phi$  such that  $\mathcal{D}_i \phi = 0$  ( $1 \leq i \leq g$ ).*

*Proof.* Each  $\phi$  in  $\Theta_0$  contains no  $u_i$  and

$$\mathcal{D}_i = \sum_{n \geq 1} \frac{1}{2\pi\sqrt{-1}} \frac{\partial}{\partial u_i} \circ \sigma^{(n)} \quad (1 \leq i \leq g),$$

hence we have  $\mathcal{D}_i \phi = 0$  ( $1 \leq i \leq g$ ). Conversely, assume

$$\mathcal{D}_i \left( \sum \alpha_{j, a/n, n} \mathcal{D}_j^{(n)} \begin{bmatrix} a/n \\ 0 \end{bmatrix} (\tau | u, z) \right) = 0 \quad (1 \leq i \leq g).$$

Then it follows

$$\sum nj_i \alpha_{j, a/n, n} \mathcal{D}_{j-\varepsilon_i}^{(n)} \begin{bmatrix} a/n \\ 0 \end{bmatrix} (\tau | u, z) = 0 \quad (1 \leq i \leq g).$$

This means  $\alpha_{j, a/n, n} = 0$  for  $j \neq 0$ .

## §2. Projection operators

2.1. In order to express the projection operators

$$\sigma_j^{(n)}: \Theta \longrightarrow \Delta^j \Theta_0^{(n)} \quad (j \in \mathbf{Z}_{\geq 0}^g, n \geq 1),$$

we need a lemma.

LEMMA 2.1.

$$(2.1) \quad \left( \sum_{p \leq k} \frac{(-1)^{|p|}}{p!} n^{-|p|} \Delta^p \mathcal{D}^p \right) \mathcal{D}_k^{(n)} \begin{bmatrix} a/n \\ 0 \end{bmatrix} (\tau | u, z) = \begin{cases} \mathcal{D}^{(n)} \begin{bmatrix} a/n \\ 0 \end{bmatrix} (\tau | z) & (k = 0) \\ 0 & (k \neq 0) \end{cases},$$

$$\begin{aligned}
 (2.2) \quad & \left( \Delta^j \left( \sum_p \frac{(-1)^{|p|}}{p!} n^{-|p|} \Delta^p \mathcal{D}^p \right) \frac{1}{j!} n^{-|j|} \mathcal{D}^j \right) \mathcal{G}_k^{(n)} \begin{bmatrix} a/n \\ 0 \end{bmatrix} (\tau | u, z) \\
 & = \begin{cases} \mathcal{G}_j^{(n)} \begin{bmatrix} a/n \\ 0 \end{bmatrix} (\tau | u, z) & (k = j) \\ 0 & (k \neq j) \end{cases}, \\
 & (j, k \in \mathbb{Z}_{\geq 0}^g, a \in \mathbb{Z}^g / n\mathbb{Z}^g, n \geq 1).
 \end{aligned}$$

*Proof.* From (1.4), (1.5), (1.6), (1.7) it follows

$$\begin{aligned}
 & \left( \sum_p \frac{(-1)^{|p|}}{p!} n^{-|p|} \Delta^p \mathcal{D}^p \right) \mathcal{G}_k^{(n)} \begin{bmatrix} a/n \\ 0 \end{bmatrix} (\tau | u, z) \\
 & = \sum_{p \leq k} (-1)^{|p|} \binom{k}{p} \Delta^p \mathcal{G}_{k-p}^{(n)} \begin{bmatrix} a/n \\ 0 \end{bmatrix} (\tau | u, z) \\
 & = \left( \sum_{p \leq k} (-1)^{|p|} \binom{k}{p} \right) \cdot \mathcal{G}_k^{(n)} \begin{bmatrix} a/n \\ 0 \end{bmatrix} (\tau | u, z) \\
 & = \begin{cases} \mathcal{G}^{(n)} \begin{bmatrix} a/n \\ 0 \end{bmatrix} (\tau | z) & (k = 0) \\ 0 & (k \neq 0) \end{cases}, \\
 & \left( \Delta^j \left( \sum_p \frac{(-1)^{|p|}}{p!} n^{-|p|} \Delta^p \mathcal{D}^p \right) \frac{1}{j!} n^{-|j|} \mathcal{D}^j \right) \mathcal{G}_k^{(n)} \begin{bmatrix} a/n \\ 0 \end{bmatrix} (\tau | u, z) \\
 & = \Delta^j \left( \sum_p \frac{(-1)^{|p|}}{p!} n^{-|p|} \Delta^p \mathcal{D}^p \right) \binom{k}{j} \mathcal{G}_{k-j}^{(n)} \begin{bmatrix} a/n \\ 0 \end{bmatrix} (\tau | u, z) \\
 & = \binom{k}{j} \Delta^j \left( \sum_p \frac{(-1)^{|p|}}{p!} n^{-|p|} \Delta^p \mathcal{D}^p \right) \mathcal{G}_{k-j}^{(n)} \begin{bmatrix} a/n \\ 0 \end{bmatrix} (\tau | u, z) \\
 & = \begin{cases} \Delta^j \mathcal{G}^{(n)} \begin{bmatrix} a/n \\ 0 \end{bmatrix} (\tau | z) = \mathcal{G}_j^{(n)} \begin{bmatrix} a/n \\ 0 \end{bmatrix} (\tau | u, z) & (j = k) \\ 0 & (j \neq k) \end{cases}
 \end{aligned}$$

**THEOREM 2.1.**  $\Theta$  has the direct sum decomposition

$$(2.3) \quad \Theta = \sum_{i \in \mathbb{Z}_{\geq 0}^g} \Delta^i \Theta_0 = \sum_{n \geq 1} \sum_{j \in \mathbb{Z}_{\geq 0}^g} \Delta^j \Theta_0^{(n)}$$

such that  $\Delta^j$  induces a vector space isomorphism of  $\Theta_0^{(n)}$  onto  $\Delta^j \Theta_0^{(n)}$ . The projection operators

$$\sigma_j^{(n)}: \Theta \longrightarrow \Delta^j \Theta_0^{(n)}$$

are given by

$$\begin{aligned}
 (2.4) \quad & \sigma_j^{(n)} = \Delta^j \left( \sum_p \frac{(-1)^{|p|}}{p!} n^{-|p|} \Delta^p \mathcal{D}^p \right) \frac{1}{j!} n^{-|j|} \mathcal{D}^j \circ \sigma^{(n)} \\
 & (j \in \mathbb{Z}_{\geq 0}^g, n \geq 1).
 \end{aligned}$$

*Proof.* The first part of the assertion is a direct consequence of the fact:  $\left\{ \mathcal{D}_j^{(n)} \begin{bmatrix} a/n \\ 0 \end{bmatrix} (\tau | u, z) \mid j \in \mathbb{Z}_{\geq 0}, a \in \mathbb{Z}^g / n\mathbb{Z}^g, n \geq 1 \right\}$ ,  $\left\{ \mathcal{D}_j^{(n)} \begin{bmatrix} a/n \\ 0 \end{bmatrix} (\tau | u, z) \mid a \in \mathbb{Z}^g / n\mathbb{Z}^g \right\}$  and  $\left\{ \mathcal{D}^{(n)} \begin{bmatrix} a/n \\ 0 \end{bmatrix} (\tau | z) \mid a \in \mathbb{Z}^g / n\mathbb{Z}^g \right\}$  are the basis of  $\Theta$ ,  $\Delta^j \Theta_0^{(n)}$  and  $\Theta_0^{(n)}$ , respectively. The expression (2.4) is a direct consequence of (2.2).

**COROLLARY.** *The inverse mapping of  $\Delta^j: \Theta_0^{(n)} \rightarrow \Delta^j \Theta_0^{(n)}$  is given by*

$$(2.5) \quad \left( \sum_p \frac{(-1)^{p|j}}{p!} n^{-|p|} \Delta^p \mathcal{D}^p \right) \frac{1}{j!} n^{-|j|} \mathcal{D}^j \quad (j \in \mathbb{Z}_{\geq 0}, n \geq 1).$$

*Proof.* Since the mapping

$$\mathcal{D}^{(n)} \begin{bmatrix} a/n \\ 0 \end{bmatrix} (\tau | z) \longrightarrow \Delta^j \mathcal{D}^{(n)} \begin{bmatrix} a/n \\ 0 \end{bmatrix} (\tau | z) = \mathcal{D}_j^{(n)} \begin{bmatrix} a/n \\ 0 \end{bmatrix} (\tau | u, z)$$

is a bijection, (2.4) implies (2.5).

### §3. Decomposition theorem on differential polynomials of theta functions

**3.1.** First let us prove the algebra isomorphic theorem:

**THEOREM 3.1.** *The replacement*

$$\Delta^j \varphi(z) \longrightarrow \left( \frac{\partial}{\partial z} \right)^j \varphi(z) \quad (j \in \mathbb{Z}_{\geq 0}, \varphi \in \Theta_0)$$

*induces a  $\Theta_0$ -algebra isomorphism of  $\Theta$  onto the algebra*

$$\mathbb{C} \left[ \dots, \left( \frac{\partial}{\partial z} \right)^j \mathcal{D}^{(n)} \begin{bmatrix} a/n \\ 0 \end{bmatrix} (\tau | z), \dots \right]$$

*of differential polynomials of theta functions, namely*

- 1°  $G \left( \dots, \Delta^j \mathcal{D}^{(n)} \begin{bmatrix} a/n \\ 0 \end{bmatrix} (\tau | z), \dots \right) = 0,$   
*if and only if*  $G \left( \dots, \left( \frac{\partial}{\partial z} \right)^j \mathcal{D}^{(n)} \begin{bmatrix} a/n \\ 0 \end{bmatrix} (\tau | z), \dots \right) = 0,$
- 2°  $G \left( \dots, \Delta^j \mathcal{D}^{(n)} \begin{bmatrix} a/n \\ 0 \end{bmatrix} (\tau | z), \dots \right) = G \left( \dots, \left( \frac{\partial}{\partial z} \right)^j \mathcal{D}^{(n)} \begin{bmatrix} a/n \\ 0 \end{bmatrix} (\tau | z), \dots \right)$   
*if and only if*  $G \left( \dots, \left( \frac{\partial}{\partial z} \right)^j \mathcal{D}^{(n)} \begin{bmatrix} a/n \\ 0 \end{bmatrix} (\tau | z), \dots \right) \in \Theta_0.$

*Proof.* It is enough to assume  $G \left( \dots, \Delta^j \mathcal{D}^{(n)} \begin{bmatrix} a/n \\ 0 \end{bmatrix} (\tau | u, z), \dots \right)$  belongs



to  $\Theta^{(m)}$  with some  $m$ . If  $G\left(\dots, \Delta^j \mathcal{G}^{(n)} \begin{bmatrix} a/n \\ 0 \end{bmatrix}(\tau|z), \dots\right) = 0$ , then putting  $u = 0$ , we obtain  $G\left(\dots, \left(\frac{\partial}{\partial z}\right)^j \mathcal{G}^{(n)} \begin{bmatrix} a/n \\ 0 \end{bmatrix}(\tau|z), \dots\right) = 0$ . By virtue of the direct decomposition theorem we may put

$$G\left(\dots, \Delta^j \mathcal{G}^{(n)} \begin{bmatrix} a/n \\ 0 \end{bmatrix}(\tau|z), \dots\right) = \sum_h \Delta^h \phi_h(z)$$

with  $\phi_h \in \Theta_0^{(m)}$ . If we assume  $G\left(\dots, \left(\frac{\partial}{\partial z}\right)^j \mathcal{G}^{(n)} \begin{bmatrix} a/n \\ 0 \end{bmatrix}(\tau|z), \dots\right) = 0$ , then we have

$$\begin{aligned} \sum_h \left(\frac{\partial}{\partial z}\right)^h \phi_h(z) &= G\left(\dots, \Delta^j \mathcal{G}^{(n)} \begin{bmatrix} a/n \\ 0 \end{bmatrix}(\tau|z), \dots\right)_{|u=0} \\ &= G\left(\dots, \left(\frac{\partial}{\partial z}\right)^j \mathcal{G}^{(n)} \begin{bmatrix} a/n \\ 0 \end{bmatrix}(\tau|z), \dots\right) = 0. \end{aligned}$$

Therefore it is enough to show  $\phi_h(z) = 0$  under the condition

$$\sum_h \left(\frac{\partial}{\partial z}\right)^h \phi_h(z) = 0 \quad \text{and} \quad \phi_h(z) \in \Theta_0^{(m)}.$$

For each  $b \in \mathbb{Z}^g$  it follows

$$\begin{aligned} \phi_h(z + b\tau) &= \exp(-\pi m \sqrt{-1}(b\tau^t b + 2z^t b)) \phi_h(z), \\ \sum_h \left(\frac{\partial}{\partial z}\right)^h \phi_h(z + b\tau) &= \sum_h \left(\frac{\partial}{\partial z}\right)^h (\exp(-\pi m \sqrt{-1}(b\tau^t b + 2z^t b)) \phi_h(z)) \\ &= \exp(-\pi m \sqrt{-1}(b\tau^t b + 2z^t b)) \sum_h \sum_p \binom{h}{p} \\ &\quad \cdot (-2\pi m \sqrt{-1}b)^p \left(\frac{\partial}{\partial z}\right)^{h-p} \phi_h(z) \quad (b \in \mathbb{Z}^g), \end{aligned}$$

and thus

$$(*) \quad \sum_h \sum_p \binom{h}{p} (-2\pi m \sqrt{-1}b)^p \left(\frac{\partial}{\partial z}\right)^{h-p} \phi_h(z) = 0 \quad (b \in \mathbb{Z}^g).$$

Let  $h_0$  be one of maximal  $h$  in the above sum. Then, the coefficients of  $b^{h_0}$  in the polynomial relation (\*) in  $b$  is given by  $(-2\pi m \sqrt{-1})^{|h_0|} \phi_{h_0}(z)$ , hence we may conclude  $\phi_{h_0}(z) = 0$ . Proceeding this process successively we have  $\phi_h(z) = 0$ , i.e.  $G\left(\dots, \Delta^j \mathcal{G}^{(n)} \begin{bmatrix} a/n \\ 0 \end{bmatrix}(\tau|z), \dots\right) = 0$ . Since  $G\left(\dots, \Delta^j \mathcal{G}^{(n)} \begin{bmatrix} a/n \\ 0 \end{bmatrix}(\tau|z), \dots\right)$  belongs to  $\Theta^{(m)}$ , assuming

$$G\left(\dots, \Delta^j \mathcal{G}^{(n)} \begin{bmatrix} a/n \\ 0 \end{bmatrix}(\tau|z), \dots\right) = G\left(\dots, \left(\frac{\partial}{\partial z}\right)^j \mathcal{G}^{(n)} \begin{bmatrix} a/n \\ 0 \end{bmatrix}(\tau|z), \dots\right),$$

we have

$$\begin{aligned} & G\left(\dots, \left(\frac{\partial}{\partial z}\right)^j \mathcal{G}^{(n)} \begin{bmatrix} a/n \\ 0 \end{bmatrix}(\tau|z), \dots\right)_{|z \rightarrow z + \hat{b} + \hat{b}\tau} \\ &= G\left(\dots, \Delta^j \mathcal{G}^{(n)} \begin{bmatrix} a/n \\ 0 \end{bmatrix}(\tau|z), \dots\right)_{|(u, z) \rightarrow (u + b, z + \hat{b} + \hat{b}\tau)} \\ &= \exp(-\pi m \sqrt{-1}(b\tau' b + 2z' b)) G\left(\dots, \Delta^j \mathcal{G}^{(n)} \begin{bmatrix} a/n \\ 0 \end{bmatrix}(\tau|z), \dots\right) \\ &= \exp(-\pi m \sqrt{-1}(b\tau' b + 2z' b)) G\left(\dots, \left(\frac{\partial}{\partial z}\right)^j \mathcal{G}^{(n)} \begin{bmatrix} a/n \\ 0 \end{bmatrix}(\tau|z), \dots\right) \end{aligned}$$

i.e.

$$G\left(\dots, \left(\frac{\partial}{\partial z}\right)^j \mathcal{G}^{(n)} \begin{bmatrix} a/n \\ 0 \end{bmatrix}(\tau|z), \dots\right) \in \Theta_0^{(m)}.$$

Conversely, if

$$G\left(\dots, \left(\frac{\partial}{\partial z}\right)^j \mathcal{G}^{(n)} \begin{bmatrix} a/n \\ 0 \end{bmatrix}(\tau|z), \dots\right) \in \Theta_0^{(m)},$$

then applying  $1^\circ$  for

$$\begin{aligned} & F\left(\dots, \Delta^j \mathcal{G}^{(n)} \begin{bmatrix} a/n \\ 0 \end{bmatrix}(\tau|z), \dots\right) \\ &= G\left(\dots, \Delta^j \mathcal{G}^{(n)} \begin{bmatrix} a/n \\ 0 \end{bmatrix}(\tau|z), \dots\right) - G\left(\dots, \left(\frac{\partial}{\partial z}\right)^j \mathcal{G}^{(n)} \begin{bmatrix} a/n \\ 0 \end{bmatrix}(\tau|z), \dots\right) \end{aligned}$$

we obtain

$$F\left(\dots, \Delta^j \mathcal{G}^{(n)} \begin{bmatrix} a/n \\ 0 \end{bmatrix}(\tau|z), \dots\right) = 0,$$

i.e.

$$G\left(\dots, \Delta^j \mathcal{G}^{(n)} \begin{bmatrix} a/n \\ 0 \end{bmatrix}(\tau|z), \dots\right) = G\left(\dots, \left(\frac{\partial}{\partial z}\right)^j \mathcal{G}^{(n)} \begin{bmatrix} a/n \\ n \end{bmatrix}(\tau|z), \dots\right).$$

Combining Theorem 2.1 and Theorem 3.1 we obtain the decomposition theorem.

**THEOREM 3.2.** *The algebra  $C\left[\dots, \left(\frac{\partial}{\partial z}\right)^j \mathcal{G}^{(n)} \begin{bmatrix} a/n \\ 0 \end{bmatrix}(\tau|z), \dots\right]$  of differential polynomials of theta functions has a canonical linear basis*

$$(3.1) \quad \left\{ \left( \frac{\partial}{\partial z} \right)^j \mathcal{G}^{(n)} \left[ \begin{matrix} a/n \\ 0 \end{matrix} \right] (\tau | z) \mid j \in \mathbb{Z}_{\geq 0}^g, a \in \mathbb{Z}^g / n\mathbb{Z}^g, n \geq 1 \right\},$$

namely differential polynomials of theta functions are uniquely expressed as linear combinations of (3.1) with constant coefficients depending on  $\tau$ .

**3.2.** In order to express the decomposition of differential polynomials of theta functions explicitly, we introduce differential polynomials in  $Y_1, \dots, Y_r$

$$(3.2) \quad \begin{aligned} & F_{j^{(1)}, \dots, j^{(r)}; h}^{(n_1, \dots, n_r)}(Y_1, \dots, Y_r | z) \\ &= \frac{1}{h!(n_1 + \dots + n_r)^{|h|}} \sum_p \frac{(-1)^{|p|}}{p!} \left( \frac{1}{n_1 + \dots + n_r} \frac{\partial}{\partial z} \right)^p \\ & \cdot \left\{ \sum_{\substack{k^{(1)} + \dots + k^{(r)} = p+h \\ k^{(\alpha)} \leq j^{(\alpha)}}} \binom{p+h}{k^{(1)}, \dots, k^{(r)}} \frac{1}{(j^{(1)} - k^{(1)})!} \dots \frac{1}{(j^{(r)} - k^{(r)})!} \right. \\ & \cdot \left. \left( \frac{1}{n_1} \frac{\partial}{\partial z} \right)^{j^{(1)} - k^{(1)}} Y_1 \dots \left( \frac{1}{n_r} \frac{\partial}{\partial z} \right)^{j^{(r)} - k^{(r)}} Y_r \right\} \\ & (j^{(1)}, \dots, j^{(r)}, h \in \mathbb{Z}_{\geq 0}^g, n_1, \dots, n_r \geq 1). \end{aligned}$$

**THEOREM 3.3.** For theta functions  $\varphi_\alpha(z) \in \Theta_0^{(n_\alpha)}$  ( $1 \leq \alpha \leq r$ )  $F_{j^{(1)}, \dots, j^{(r)}; h}^{(n_1, \dots, n_r)} \times (\varphi_1, \dots, \varphi_r | z)$ , ( $j^{(1)}, \dots, j^{(r)}, h \in \mathbb{Z}_{\geq 0}^g$ ) are theta functions of level  $n_1 + \dots + n_r$  such that

$$(3.3) \quad \begin{aligned} & \frac{1}{j^{(1)}! \dots j^{(r)}!} \left( \frac{1}{n_1} \frac{\partial}{\partial z} \right)^{j^{(1)}} \varphi_1(z) \dots \left( \frac{1}{n_r} \frac{\partial}{\partial z} \right)^{j^{(r)}} \varphi_r(z) \\ &= \sum_{h \leq j^{(1)} + \dots + j^{(r)}} \left( \frac{\partial}{\partial z} \right)^h F_{j^{(1)}, \dots, j^{(r)}; h}^{(n_1, \dots, n_r)}(\varphi_1, \dots, \varphi_r | z) \\ &= \sum_c \lambda_{(j^{(1)}, \dots, j^{(r)}; h), c/(n_1 + \dots + n_r)}(\varphi_1, \dots, \varphi_r) \left( \frac{\partial}{\partial z} \right)^h \mathcal{G}^{(n_1 + \dots + n_r)} \\ & \cdot \left[ \begin{matrix} c/(n_1 + \dots + n_r) \\ 0 \end{matrix} \right] (\tau | z), \end{aligned}$$

where

$$(3.4) \quad \begin{aligned} & \lambda_{(j^{(1)}, \dots, j^{(r)}; h), c/(n_1 + \dots + n_r)}(\varphi_1, \dots, \varphi_r) \\ &= \frac{1}{(n_1 + \dots + n_r)^{|c|}} \sum_{\tilde{c} \in \mathbb{Z}^g / (n_1 + \dots + n_r)\mathbb{Z}^g} \exp \frac{2\pi \sqrt{-1} \tilde{c}^t c}{n_1 + \dots + n_r} \\ & \cdot \mathcal{G}^{(n_1 + \dots + n_r)} \left[ \begin{matrix} c/(n_1 + \dots + n_r) \\ 0 \end{matrix} \right] (\tau | 0)^{-1} F_{j^{(1)}, \dots, j^{(r)}; h}^{(n_1, \dots, n_r)} \\ & \cdot \left( \varphi_1, \dots, \varphi_r \mid \frac{\hat{c}}{n_1 + \dots + n_r} \right). \end{aligned}$$

*Proof.* Putting

$$\begin{aligned} & \frac{1}{j^{(1)}! \cdots j^{(r)}!} \left(\frac{1}{n_1} \Delta\right)^{j^{(1)}} \varphi_1(z) \cdots \left(\frac{1}{n_r} \Delta\right)^{j^{(r)}} \varphi_r(z) \\ &= \sum \Delta^h \psi_h(z) \end{aligned}$$

with  $\psi_h(z) \in \Theta_0^{(n_1, \dots, n_r)}$ , by virtue of Corollary of Theorem 2.1 (1.6) and (1.7) we have

$$\begin{aligned} \psi_h(z) &= \frac{1}{h!(n_1 + \cdots + n_r)^{|h|}} \sum_p \frac{(-1)^{|p|}}{p!} \left(\frac{1}{n_1 + \cdots + n_r} \Delta\right)^p \mathcal{D}^{p+h} \\ & \quad \cdot \frac{1}{j^{(1)}! \cdots j^{(r)}!} \left(\frac{1}{n_1} \Delta\right)^{j^{(1)}} \varphi_1(z) \cdots \left(\frac{1}{n_r} \Delta\right)^{j^{(r)}} \varphi_r(z) \\ &= \frac{1}{h!(n_1 + \cdots + n_r)^{|h|}} \sum_p \frac{(-1)^{|p|}}{p!} \left(\frac{1}{n_1 + \cdots + n_r} \Delta\right)^p \\ & \quad \left\{ \sum_{k^{(1)} + \cdots + k^{(r)} = p+h} \left[ \begin{matrix} p+h \\ k^{(1)}, \dots, k^{(r)} \end{matrix} \right] \frac{n_1^{|k^{(1)}|} \cdots n_r^{|k^{(r)}|}}{j^{(1)}! \cdots j^{(r)}! n_1^{|j^{(1)}|} \cdots n_r^{|j^{(r)}|}} \right. \\ & \quad \left. \cdot n^{-|k^{(1)}|} \mathcal{D}^{k^{(1)}} \Delta^{j^{(1)}} \varphi_1(z) \cdots n^{-|k^{(r)}|} \mathcal{D}^{k^{(r)}} \Delta^{j^{(r)}} \varphi_r(z) \right\} \\ &= \frac{1}{h!(n_1 + \cdots + n_r)^{|h|}} \sum_p \frac{(-1)^{|p|}}{p!} \left(\frac{1}{n_1 + \cdots + n_r} \Delta\right)^p \\ & \quad \left\{ \sum_{\substack{k^{(1)} + \cdots + k^{(r)} = p+h \\ k^{(a)} \leq j^{(a)}}} \left[ \begin{matrix} p+h \\ k^{(1)}, \dots, k^{(r)} \end{matrix} \right] \frac{1}{(j^{(1)} - k^{(1)})! \cdots (j^{(r)} - k^{(r)})!} \right. \\ & \quad \left. \cdot \left(\frac{1}{n_1} \Delta\right)_{\varphi_1(z)}^{j^{(1)} - k^{(1)}} \cdots \left(\frac{1}{n_r} \Delta\right)_{\varphi_r(z)}^{j^{(r)} - k^{(r)}} \right\} \\ &= F_{j^{(1)}, \dots, j^{(r)}; h}^{(n_1, \dots, n_r)}(\varphi_1, \dots, \varphi_r | z). \end{aligned}$$

Hence, replacing  $\Delta_i$  by  $\partial/\partial z_i$  ( $1 \leq i \leq g$ ), we prove the first assertion of Theorem 3.3. Putting

$$\begin{aligned} & F_{j^{(1)}, \dots, j^{(r)}; h}^{(n_1, \dots, n_r)}(\varphi_1, \dots, \varphi_r | z) \\ &= \sum_{c \in \mathbb{Z}^g / (n_1 + \cdots + n_r) \mathbb{Z}^g} \lambda_{h,c} \mathcal{D}^{(n_1 + \cdots + n_r)} \left[ \begin{matrix} c / (n_1 + \cdots + n_r) \\ 0 \end{matrix} \right] (\tau | z), \end{aligned}$$

we have

$$\begin{aligned} & F_{j^{(1)}, \dots, j^{(r)}; h}^{(n_1, \dots, n_r)} \left( \varphi_1, \dots, \varphi_r \left| \frac{\hat{c}}{n_1 + \cdots + n_r} \right. \right) \\ &= \sum_c \lambda_{h,c} \mathcal{D}^{(n_1 + \cdots + n_r)} \left[ \begin{matrix} c / (n_1 + \cdots + n_r) \\ 0 \end{matrix} \right] \left( \tau \left| \frac{\hat{c}}{n_1 + \cdots + n_r} \right. \right) \\ &= \sum_c \lambda_{h,c} \exp \left( \frac{2\pi \sqrt{-1} \hat{c}^t c}{n_1 + \cdots + n_r} \right) \mathcal{D}^{(n_1 + \cdots + n_r)} \left[ \begin{matrix} c / (n_1 + \cdots + n_r) \\ 0 \end{matrix} \right] (\tau | 0) \\ & \quad (c \in \mathbb{Z}^g / (n_1 + \cdots + n_r) \mathbb{Z}^g). \end{aligned}$$

Hence, by virtue of the orthogonal relation for characters

$$\sum_c \exp\left(\frac{2\pi\sqrt{-1}\hat{c}^t c}{n_1 + \dots + n_r}\right) = \begin{cases} (n_1 + \dots + n_r)^g & \hat{c} \equiv 0 \pmod{n_1 + \dots + n_r} \\ 0 & \hat{c} \not\equiv 0 \pmod{n_1 + \dots + n_r}, \end{cases}$$

it follows

$$\lambda_{h,c} = \frac{1}{(n_1 + \dots + n_r)^g} \sum_c \exp\left(\frac{-2\pi\sqrt{-1}\hat{c}^t c}{n_1 + \dots + n_r}\right) \mathcal{G}^{(n_1 + \dots + n_r)} \cdot \left[ \frac{c/(n_1 + \dots + n_r)}{0} \right] (\tau|0)^{-1} F_{j^{(1)}, \dots, j^{(r)}; h}^{(n_1, \dots, n_r)} \left( \varphi_1, \dots, \varphi_r \middle| \frac{\hat{c}}{n_1 + \dots + n_r} \right).$$

Specializing

$$(\varphi_1(z), \varphi_2(z)) \quad \text{to} \quad \left( \mathcal{G}^{(n_1)} \left[ \frac{a_1/n_1}{0} \right] (\tau|z), \mathcal{G}^{(n_r)} \left[ \frac{a_r/n_r}{0} \right] (\tau|z) \right),$$

we obtain the explicit expression of structure constants of

$$C \left[ \dots, \left( \frac{\partial}{\partial z} \right)^j \mathcal{G}^{(n)} \left[ \frac{a/n}{0} \right] (\tau|z), \dots \right]$$

with respect to the basis

$$\left\{ \left( \frac{\partial}{\partial z} \right)^j \mathcal{G}^{(n)} \left[ \frac{a/n}{0} \right] (\tau|z) \right\}.$$

**THEOREM 3.4.** *The structure constants of*

$$C \left[ \dots, \left( \frac{\partial}{\partial z} \right)^j \mathcal{G}^{(n)} \left[ \frac{a/n}{0} \right] (\tau|z), \dots \right]$$

are given by

$$\begin{aligned} & \left( \frac{\partial}{\partial z} \right)^{j^{(1)}} \mathcal{G}^{(n_1)} \left[ \frac{a_1/n_1}{0} \right] (\tau|z) \left( \frac{\partial}{\partial z} \right)^{j^{(2)}} \mathcal{G}^{(n_2)} \left[ \frac{a_2/n_2}{0} \right] (\tau|z) \\ &= \sum_h \sum_c \gamma_{(j^{(1)}, a_1/n_1, n_1), (j^{(2)}, a_2/n_2, n_2)}^{(h, c/(n_1+n_2), n_1+n_2)}(\tau) \left( \frac{\partial}{\partial z} \right)^h \mathcal{G}^{(n_1+n_2)} \left[ \frac{c/(n_1+n_2)}{0} \right] (\tau|z), \\ (3.5) \quad & \gamma_{(j^{(1)}, a_1/n_1, n_1), (j^{(2)}, a_2/n_2, n_2)}^{(h, c/(n_1+n_2), n_1+n_2)}(\tau) \\ &= \frac{j^{(1)}! j^{(2)}! n_1^{j^{(1)}} n_2^{j^{(2)}}}{h! (n_1 + n_2)^{g+|h|}} \sum_{\hat{c} \in \mathbb{Z}^g / (n_1+n_2)\mathbb{Z}^g} \exp\left(\frac{-2\pi\sqrt{-1}\hat{c}^t c}{n_1 + n_2}\right) \mathcal{G}^{(n_1+n_2)} \\ & \cdot \left[ \frac{c/(n_1+n_2)}{0} \right] (\tau|0)^{-1} \left[ \sum_p \frac{(-1)^{|p|}}{p!} \left( \frac{1}{n_1 + n_2} \frac{\partial}{\partial z} \right)^p \sum_{\substack{k^{(1)}+k^{(2)}=p+h \\ k^{(1)} \leq j^{(1)}, k^{(2)} \leq j^{(2)}}} \right. \\ & \cdot \left. \frac{[p+h]}{[k^{(1)}, k^{(2)}]} \frac{1}{(j^{(1)} - k^{(1)})! (j^{(2)} - k^{(2)})!} \left( \frac{1}{n_1} \frac{\partial}{\partial z} \right)^{j^{(1)} - k^{(1)}} \mathcal{G}^{(n_1)} \left[ \frac{a_1/n_1}{0} \right] \right] \end{aligned}$$

$$\cdot (\tau | z) \left( \frac{1}{n_2} \frac{\partial}{\partial z} \right)^{j^{(2)} - k^{(2)}} \mathcal{G}^{(n_2)} \begin{bmatrix} a_2/n_2 \\ 0 \end{bmatrix} (\tau | z) \Big|_{z=\hat{z}(n_1+n_2)} \cdot$$

For theta functions  $\varphi_\alpha(z)$  ( $1 \leq \alpha \leq z$ ), if a differential polynomial  $G(\dots, (\partial/\partial z)^j \varphi_\alpha(z), \dots)$  is a theta function, then by virtue of Theorems 2.1 and 3.1  $G(\dots, (\partial/\partial z)^j \varphi_\alpha(z), \dots)$  is itself the  $\theta_0$ -component of the decomposition. Hence, Theorem 3.4 implies the following characterization of differential polynomials of  $\varphi_\alpha(z)$  ( $1 \leq \alpha \leq r$ ) which are also theta functions.

**THEOREM 3.5.** *For theta functions  $\varphi_\alpha(z) \in \theta_0^{(n_\alpha)}$  the space*

$$\mathbb{C} \left[ \dots, \left( \frac{\partial}{\partial z} \right)^j \varphi_\alpha(z), \dots \right] \cap \theta_0^{(m)}$$

*is linearly spanned by*

$$F_{\binom{(n_1, \dots, n_1, \dots, n_r, \dots, n_r)}{(j^{(1,1)}, \dots, j^{(1, \epsilon_1)}, \dots, j^{(r,1)}, \dots, j^{(r, \epsilon_r)}}; 0} \underbrace{\varphi_1, \dots, \varphi_1}_{\epsilon_1}, \dots, \underbrace{\varphi_r, \dots, \varphi_r}_{\epsilon_r} | z) \\ (\sum_\alpha e_\alpha n_\alpha = m; j^{(1,1)}, \dots, j^{(1, \epsilon_1)}, \dots, j^{(r,1)}, \dots, j^{(r, \epsilon_r)} \in \mathbf{Z}_{\geq 0}^{\epsilon}).$$

REFERENCES

- [ 1 ] R. Hirota, A direct method of finding exact solution of nonlinear evolution equations, *Lecture Notes in Mathematics*, No. 515, 40–68 (1976).
- [ 2 ] H. Morikawa, Some analytic and geometric applications of the invariant theoretic method, *Nagoya Math. J.*, **30** (1980), 1–47.
- [ 3 ] ———, On Poisson brackets of semi-invariants, *Manifolds and Lie groups*, 267–281, *Progress in Math.* Birkhäuser (1981).

*Department of Mathematics  
Faculty of Science  
Nagoya University  
Chikusa-ku 464  
Japan*