

A GENERAL LIMIT THEORY FOR NONLINEAR FUNCTIONALS OF NONSTATIONARY TIME SERIES

QIYING WANG
University of Sydney

PETER C. B. PHILLIPS 
*Yale University, University of Auckland, and Singapore Management
University*

New limit theory is provided for a wide class of sample variance and covariance functionals involving both nonstationary and stationary time series. Sample functionals of this type commonly appear in regression applications and the asymptotics are particularly relevant to estimation and inference in nonlinear nonstationary regressions that involve unit root, local unit root, or fractional processes. The limit theory is unusually general in that it covers both parametric and nonparametric regressions. Self-normalized versions of these statistics are considered that are useful in inference. Numerical evidence reveals interesting strong bimodality in the finite sample distributions of conventional self-normalized statistics similar to the bimodality that can arise in t -ratio statistics based on heavy tailed data. Bimodal behavior in these statistics is due to the presence of long memory innovations and is shown to persist for very large sample sizes even though the limit theory is Gaussian when the long memory innovations are stationary. Bimodality is shown to occur even in the limit theory when the long memory innovations are nonstationary. To address these complications, new self-normalized versions of the test statistics are introduced that deliver improved approximations that can be used for inference.

1. INTRODUCTION

Parametric and nonparametric regressions with nonstationary data have attracted considerable recent attention because of the prevalence of nonstationary time series in applied work across many different disciplines and the need for asymptotic theory to support methods of estimation and inference in the presence of nonstationarity. Much of this work has focused on cointegrating regression where linkages between nonstationary processes and stationary innovations play an integral role in the notion of cointegration and its various extensions to fractional

Wang acknowledges research support from the Australian Research Council (Grant No. DP170104385). Phillips acknowledges research support from the NSF (Grant No. SES 18-50860) and a Kelly Fellowship at the University of Auckland. Address correspondence to Qiyang Wang, University of Sydney, Sydney, NSW, Australia; e-mail: qiyang.wang@sydney.edu.au.

© The Author(s), 2024. Published by Cambridge University Press. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted re-use, distribution, and reproduction in any medium, provided the original work is properly cited.

processes involving long memory time series. The literature in this area is now voluminous, as discussed in recent papers (e.g., Duffy and Kasparis, 2021; Wang, Phillips, and Kasparis, 2021). Readers are referred to Park (2014) and Tjøstheim (2020) for partial overviews of the field of nonlinear cointegration studies that cover many of the relevant contributions and empirical applications. In almost all of this literature, a key role in the asymptotic development is played by sample covariance functionals that involve (possibly nonlinear functions of) nonstationary processes and stationary time series. Sample covariances of this type take similar but subtly different forms in parametric and nonparametric regressions. They typically appear in signal functions and score functions whose asymptotic behavior is critical in determining the limit theory needed for estimation, inference, and specification testing in such regressions. Prototypical forms of these functionals for nonparametric and parametric cases are shown below in (1.3) and (1.4) by R_{2n} and $R_{2n}(\theta^0)$. The goal of the present article is to extend existing results on such functionals, accommodate these two forms in a general limit theory, and develop self-normalized statistics that will be useful for inference in regression. We open the discussion with three illustrative examples.

In the nonparametric case, simple nonlinear nonstationary regressions typically have the form

$$y_k = g(x_k) + u_k, \quad k = 1, \dots, n, \tag{1.1}$$

with an $I(1)$ regressor generated by the partial sum model $x_k = x_{k-1} + \xi_k$ with weakly dependent and possibly correlated innovations $\{u_k, \xi_k\}$, thereby allowing for endogeneity. In the nonparametric case, the nonlinear cointegrating function $g(x_k)$ may be estimated at some point x by local-level kernel regression in the usual manner via the criterion

$$\mathcal{Q}_{n,h}(g) = \sum_{k=1}^n K_h(x_k - x)(y_k - g(x_k))^2, \tag{1.2}$$

giving $\hat{g}(x) = \operatorname{argmin}_g \mathcal{Q}_{n,h}(g) = \frac{\sum_{k=1}^n y_k K_h(x_k - x)}{\sum_{k=1}^n K_h(x_k - x)}$, where $K_h(s) = \frac{1}{h} K(\frac{s}{h})$, $K(\cdot)$ is a nonnegative real kernel function and the bandwidth parameter $h = h_n \rightarrow 0$ as $n \rightarrow \infty$. The limit theory of $\hat{g}(x)$ then depends on the behavior of suitably normalized forms of the two sample functionals

$$R_{1n} = \sum_{k=1}^n K_h(x_k - x) \text{ and } R_{2n} = \sum_{k=1}^n K_h(x_k - x) u_k, \tag{1.3}$$

where R_{1n} is a sample signal process and R_{2n} is a sample score process, both of which are nonlinear in the nonstationary regressor x_k . Test statistics typically also require estimation of the innovations using the regression residuals $\hat{u}_k = y_k - \hat{g}(x_k)$ and a sample functional such as $R_{3n} = \sum_{k=1}^n K_h^2(x_k - x) \hat{u}_k^2$. Full development of a limit theory for estimation and inference concerning the function $g(\cdot)$ in (1.1) requires joint convergence results for suitably normalized forms of sample functionals such as (R_{1n}, R_{2n}, R_{3n}) . In applications, allowance is typically made for

endogeneity of the regressor x_k in the regression (1.1). Importantly, as shown in the nonlinear cointegration study of Wang and Phillips (2009b), such nonparametric nonstationary regressions do not require the use of instrumental variables and do not suffer from ill-posedness, in contrast to stationary regressions and there is, in contrast therefore, no need for regularization.

In the parametric case, the nonlinear cointegrating function has a specific functional form $g(x_k) = g(x_k; \theta)$ that depends on some unknown parameter vector $\theta \in \Theta \subset \mathbb{R}^p$, where Θ is a compact subspace of \mathbb{R}^p for some finite p . The nonlinear least squares estimator is then $\hat{g}(x) = g(x; \hat{\theta})$ with $\hat{\theta} = \operatorname{argmin}_{\theta \in \Theta} Q_n(\theta)$ where $Q_n(\theta) = \sum_{k=1}^n (y_k - g(x_k; \theta))^2$. In this case, the limit theory for $\hat{\theta}$ depends on normalized versions of the sample functionals

$$R_{1n}(\theta^0) = \sum_{k=1}^n G_k^0 G_k^{0'} \text{ and } R_{2n}(\theta^0) = \sum_{k=1}^n G_k^0 u_k, \tag{1.4}$$

where $G_k^0 = \partial g(x_k; \theta^0) / \partial \theta$ and θ^0 is the true value of θ . As in the nonparametric case, test statistics usually depend on regression residuals $\hat{u}_k = y_k - g(x_k; \hat{\theta})$, leading to sample functionals such as $R_{3n}(\hat{\theta}) = \sum_{k=1}^n G_k^{\hat{\theta}} G_k^{\hat{\theta}'} \hat{u}_k^2$.

The sample variance and covariance functionals in (1.3) and (1.4) are closely related but differ because of the critical role played by the presence of the bandwidth sequence h in the functions of (1.3), making a general theory difficult. Asymptotics for regression estimation and inference in such cases have therefore been studied in past research separately and often in special cases.¹ More complex models that include spurious nonlinear regression (Phillips, 2009; Tu and Wang, 2022) and functional coefficient (FC) nonstationary regressions involve similar sample functionals for which asymptotic theory is also needed to facilitate empirical work.

FC regressions are of particular interest in applications because covariate dependence or time variation in the regression coefficients is often of interest in applications. Such models with nonstationary regressors were originally considered by Xiao (2009).² It was later shown in Phillips and Wang (2023) that important subtleties arise in such FC regressions that affect the limit theory in material ways because nonstationarity in the regressors amplifies the impact of bias in nonparametric FC regression. Models of this type are typically linear in (possibly multivariate) regressors x_k and take the form

$$y_k = \theta(z_k)' x_k + u_k, \quad k = 1, \dots, n, \tag{1.5}$$

with coefficients $\theta(z_k)$ that are smooth functions of a covariate z_k that may be stationary or nonstationary. In FC models of this type, estimation of the coefficient

¹See, for instance, Phillips and Park (1998), Park and Phillips (1999, 2000, 2001), Karlsen and Tjøstheim (2001), Wang and Phillips (2009a, 2009b), Gao and Phillips (2013), Li, Tjøstheim, and Gao (2016), Wang and Phillips (2016), and Wang et al. (2021)

²See also Cai, Li, and Park (2009), Sun and Li (2011), Sun, Cai, and Li (2016), and Liang, Shen, and Wang (2023).

functions $\theta(\cdot)$ at some point z in the domain of z_t necessarily involves the three sample functionals

$$R_{4n} = \sum_{k=1}^n x_k x'_k K_h(z_k - z),$$

$$R_{5n} = \sum_{k=1}^n K_h(z_k - z) u_k, \quad R_{6n} = \sum_{k=1}^n x_k x'_k [\theta(z_k) - \theta(z)] K_h(z_k - z), \tag{1.6}$$

where R_{6n} is an additional sample covariance bias functional that depends on the regressors, the kernel function, and bias effects that need further decomposition to fully resolve the asymptotic theory.³

These examples motivate a general formulation that is relevant in many different applications. To fix ideas, suppose an observable time series x_t is a scalar nonstationary process, either integrated $I(1)$, near $I(1)$, or a similar time series with fractional process innovations, as detailed in what follows, and $w_k = (w_{1k}, \dots, w_{dk})$ is a sequence of stationary random vectors. The article is concerned with sample quantities S_n of x_k and w_k defined by sample sums of nonlinear functions of x_k and w_k that take the general form

$$S_n = \sum_{k=1}^n f(x_k/h, w_k),$$

where $h \equiv h_n > 0$ is a sequence of positive constants indexed by the sample size n and $f(x, y)$ is a real function on R^{1+d} . The partial sum S_n is a scalar nonlinear functional of multivariate arguments that involve both stationary and nonstationary processes. Such functionals play a dominant role in the development of the theory of estimation and inference in nonlinear cointegrating regression, where the regressor is usually a nonstationary time series, including those with autoregressive unit roots and local unit root properties. In such regression contexts, a prominent example of S_n has the form of a sample covariance function that involves both the nonstationary regressor and the equation innovations. In this case, two covariance functions are most typical, one of the form $S_{1n} = \sum_{k=1}^n f(x_k, w_{2k}, \dots, w_{dk}) w_{1k}$ and the other of the form $S_{2n} = \sum_{k=1}^n f(x_k/h) w_{1k}$, where an auxiliary sequence $h = h_n$ may be present that depends on the sample size, as in nonparametric kernel regression discussed above.

As is now well known in the literature (see, for instance, Karlsen and Tjøstheim, 2001; Park and Phillips, 2001; Wang and Phillips, 2009a, 2009b; Chan and Wang, 2015; Dong and Linton, 2018; Duffy, 2020; Hu, Phillips, and Wang, 2021 and the references therein), covariance expressions such as S_{1n} occur in nonlinear parametric cointegrating regression and expressions such as S_{2n} , with the auxiliary

³As explained in Phillips and Wang (2023), the bias effect R_{6n} has both a “deterministic” component ($\sum_{k=1}^n x_k x'_k$) $\mathbb{E} \xi_{\beta k}$ and a “random” component ($\sum_{k=1}^n x_k x'_k$) $\eta_{\beta k}$ where $\xi_{\beta k} = [\beta(z_k) - \beta(z)] K_h(z_k - z)$ and $\eta_{\beta k} = \xi_{\beta k} - \mathbb{E} \xi_{\beta k}$. The presence of these two components influences the limit theory, rates of convergence, and bandwidth choice in important ways. Readers are referred to Phillips and Wang (2023) for details.

sequence h , arise naturally in Nadaraya–Watson estimation where $f(x)$ is a kernel function and $h \rightarrow 0$ is a bandwidth used in the nonparametric regression.

It transpires that the limit behavior of S_n depends on the value of the integral $\int_{-\infty}^{\infty} g(s) ds$, where $g(x) = \mathbb{E}f(x, w_1)$. When $\int_{-\infty}^{\infty} g(s) ds \neq 0$, it was shown in Wang et al. (2021) that upon suitable normalization S_n satisfies

$$\frac{d_n}{nh} S_n \rightarrow_D \int_{-\infty}^{\infty} g(x) dx L_G(1, 0), \tag{1.7}$$

provided $d_n/nh \rightarrow 0$ and $d_n/h \rightarrow \infty$, with $d_n^2 = \text{var}(x_n)$ and where $L_G(t, s)$ is the local time of a stochastic process $\mathcal{G}(t)$ at the spatial point s , as defined in the following section. Result (1.7) was established in quite general settings, generalizing and improving previous related work on convergence to local time given by Akonom (1993), Borodin and Ibragimov (1995), Phillips and Park (1998), Jeganathan (2004), Wang and Phillips (2009a, 2016), and Duffy (2016). This fundamental limit result enabled the investigation of asymptotic theory for latent variable nonparametric cointegrating regression in which some variables were observed with measurement error.

The present work is concerned with developing a limit theory for the sample function S_n in the case where $\int_{-\infty}^{\infty} g(s) ds = 0$, which is commonly known as the zero energy case. Towards this end, in some specialized cases such as $f(x, y) = m(x)$ or $f(x, y) = m(x)y$ where $m(x)$ is bounded and integrable, the asymptotic behavior of S_n is known and has been considered in Wang and Phillips (2009b, 2011), with the attendant requirement that $h \rightarrow 0$, and in an unpublished manuscript by Jeganathan (2008) (with $h = 1$). This article provides a unified extension of these existing results that encompasses the two cases where $h = 1$ and $h \rightarrow 0$, together with the setting of general functionals $f(x, y)$ rather than the specialized forms $f(x, y) = m(x)y$ or $m(x)$.

In unifying, the two standard limit cases where $h = 1$ and $h \rightarrow 0$, our work might be compared with Gozalo and Linton (2000) who showed how to nonparametrically encompass a parametric model by using a local nonlinear least squares criterion that allows for recentering a nonparametric regression on a specific parametric model. In the present context, that approach would involve replacing (1.2) with the criterion $\mathcal{Q}_{n,h}(x, \alpha) = \sum_{k=1}^n K_h(x_k - x)(y_k - m(x_k, \alpha))^2$ for some parametric function $m(x_k, \alpha)$, leading to the estimate $\hat{g}(x) = m(x, \hat{\alpha})$, where $\hat{\alpha} = \text{argmin}_{\alpha} \mathcal{Q}_{n,h}(x, \alpha)$. When the parametric form $m(x; \alpha)$ is correct or nearly correct around the point x , there is an advantage to using a wider bandwidth h in such a regression; and, if the parametric model $m(x; \alpha)$ were correct almost everywhere, there would be an advantage in letting $h \rightarrow \infty$ rather than $h \rightarrow 0$. The limit theory for this approach in Gozalo and Linton (2000) relies on an independent and identically distributed (i.i.d.) setup. Extending that approach to the present setting and exploring possible advantages of parametric information in local nonparametric nonlinear regression with nonstationary data are interesting lines of future research.

It should be mentioned that the zero energy case where the functional $\int_{-\infty}^{\infty} g(s) ds = 0$, in which $g(x) = \mathbb{E}f(x, w_1)$, arises naturally in regression

applications. For instance, in nonparametric cointegrating regression, the development of a limit theory for normalized versions of functionals such as the sample covariance S_{2n} is vital for both estimation and inference. Thus, when x_k is an $I(1)$ regressor and w_{1k} is an error process, use of the natural centralizing condition $\mathbb{E}w_{11} = 0$ in turn implies that $\int_{-\infty}^{\infty} g(s) ds = \int_{-\infty}^{\infty} f(x) dx \mathbb{E}w_{11} = 0$. Such situations arise even in complex settings where endogeneity is present (see Wang and Phillips, 2009b, 2011, 2016 for details and econometric applications). Similarly, in regression with nonstationary nonlinear heteroskedasticity when nonstationary volatility is present in the errors [with $u_t = f(x_t, w_t)$, say], the zero energy condition $\int_{-\infty}^{\infty} g(s) ds = 0$ where again $g(x) = \mathbb{E}f(x, w_1)$ is usually required for the development of an asymptotic theory. In this case, the use of general functionals such as $f(x, y)$ in the sample covariance limit theory enables a full representation of nonstationary nonlinear volatility in the regression errors.

The remainder of the article is organized as follows: Section 2 provides the main limit theory for nonlinear functionals of nonstationary time series and a series of remarks that analyze the findings and connect to later discussion. Section 3 provides numerical evidence which reveals an intriguing bimodality for self-normalized statistics that arises in finite samples and that can persist in extremely large samples even though the limit theory is Gaussian. Section 4 discusses these findings, explains the slow convergence, and shows how bimodal limit theory does arise in the presence of nonstationary long memory innovations. Alternative self-normalized statistics are considered that substantially improve finite sample performance. Concluding remarks are in Section 5. Proofs of the main results are given in Section 6 and supporting propositions and lemmas that play key roles in proving the main results are in Section 7. Proofs of the lemmas are in the Appendix.

Throughout the article, \Rightarrow denotes weak convergence of probability measures with respect to the uniform topology (see, for instance, Billingsley, 1968) and \rightarrow_D is distributional convergence in euclidean space. For a vector $A = (A_1, \dots, A_d)$, we define $\|A\| = |A_1| + \dots + |A_d|$. Constants are represented by C, C_1, C_2, \dots , which may differ in different locations.

2. MAIN RESULTS

2.1. Assumptions and Preliminaries

Let $\lambda_i = (\epsilon_i, e_i)'$, $i \in \mathbb{Z}$ be a sequence of iid random vector innovations with $\mathbb{E}\|\lambda_0\|^2 < \infty$. Let $\xi_k = \sum_{j=0}^{\infty} \phi_j \epsilon_{k-j}$ be a linear process where the coefficients $\phi_k, k \geq 0$, satisfy $\phi_0 \neq 0$ and one of the following conditions:

LM: $\phi_k \sim k^{-\mu} \rho(k)$, $1/2 < \mu < 1$ and $\rho(x)$ is a function that is slowly varying at ∞ .⁴

SM: $\sum_{k=0}^{\infty} |\phi_k| < \infty$ and $\phi \equiv \sum_{k=0}^{\infty} \phi_k \neq 0$.

⁴That is, $\rho(x)$ is a measurable function from $(0, \infty)$ to $(0, \infty)$ so that, for all $a > 0$, $\rho(ax)/\rho(x) \rightarrow 1$ as $x \rightarrow \infty$, for example, a positive constant, $\log(x)$ or $\log^b(x)$ for any real $b > 0$.

In the following development, observable nonstationary time series x_k are generated by the linear process innovations ξ_k as detailed in the near unit root process given in **A1(i)**. The inclusion of additional innovations e_i in λ_i is useful for specifying (possibly correlated) model disturbances, as in the generating mechanisms used in simulations later in the article in Sections 3 and 4. For the development of the asymptotic theory in our main results, the following assumptions are made about the components of $S_n = \sum_{k=1}^n f(x_k/h, w_k)$.

- A1** (i) $x_k = \rho_n x_{k-1} + \xi_k$, where $x_0 = 0$, $\rho_n = 1 - \gamma n^{-1}$ for some constant $\gamma \geq 0$.
- (ii) $\mathbb{E}e_1 = 0$ and $\int_{-\infty}^{\infty} |\mathbb{E}e^{it\epsilon_1}| dt < \infty$.
- A2** (a) $w_k = (w_{1k}, \dots, w_{dk})$, where $w_{ik} = \Gamma_i(\lambda_k, \dots, \lambda_{k-m_0})$ for some fixed $m_0 \geq 0$ and $\Gamma_i(\cdot), i = 1, 2, \dots, d$, are real measurable functions of their respective components.
- (b) $\mathbb{E} \|w_1\|^{\max\{2, 4\beta\}} < \infty$, where β is given in **A3(I)**.
- A3** (I) A bounded function $T(x)$ exists such that, for some $\beta > 0$,

$$|f(x, y)| \leq T(x)(1 + |y|^\beta) \quad \text{and} \quad \int_{-\infty}^{\infty} (1 + |x|)T(x)dx < \infty.$$

(II) $\int_{-\infty}^{\infty} g(x)dx = 0$, where $g(x) = \mathbb{E}f(x, w_1)$.

(III) $\int_{-\infty}^{\infty} \mathbb{E}|\hat{f}(x, w_1)|dx < \infty$, where $\hat{f}(x, y) = \int_{-\infty}^{\infty} e^{itx}f(t, y)dt$.

Assumption **A1(i)** accommodates near integrated time series x_k that are derived from either short memory (*SM*) or long memory (*LM*) innovations, thereby covering a large class of nonstationary time series. The extra distributional assumption **A1(ii)** is a smoothness condition requiring integrability of the characteristic function $\mathbb{E}e^{it\epsilon_1}$ that is often useful in establishing convergence to a local time process. The condition can be relaxed to $\limsup_{|t| \rightarrow \infty} |t|^a \mathbb{E}e^{it\epsilon_1} < \infty$ for some $a > 0$, but is generally difficult to eliminate completely in the development of limit theory for nonlinear cointegrating regression. The zero initialization $x_0 = 0$ is assumed for convenience to avoid notational clutter and can be considerably relaxed, as is well known from earlier research. In particular, all the main results still hold if instead $x_0 = o_P(d_n)$, where $d_n^2 = \text{var}(\sum_{k=1}^n \xi_k)$. It is also well-known (see, for instance, Wang, Lin, and Gulati, 2003) that

$$d_n^2 \sim \mathbb{E}\epsilon_0^2 \begin{cases} c_\mu n^{3-2\mu} \rho^2(n), & \text{under } LM, \\ \phi^2 n, & \text{under } SM, \end{cases}$$

and $x_{\lfloor nt \rfloor} / d_n \Rightarrow Z_t$ on $D[0, 1]$, where $c_\mu = \frac{1}{(1-\mu)(3-2\mu)} \int_0^\infty x^{-\mu} (x+1)^{-\mu} dx$ and

$$Z_t = W(t) + \gamma \int_0^t e^{-\gamma(t-s)} W(s) ds, \quad t \geq 0,$$

$$W(t) = \begin{cases} B_{3/2-\mu}(t), & \text{under } LM, \\ B_{1/2}(t), & \text{under } SM, \end{cases}$$

and $B_H(t)$ is fractional Brownian motion with Hurst exponent H and $B_{1/2}(t)$ is standard Brownian motion. In this event, Z_t is a fractional Ornstein–Uhlenbeck

process, having a continuous local time process which we denote by $L_Z(t, x)$. As in Geman and Horowitz (1980), the local time process $L_Z(t, x)$ is defined as

$$L_Z(t, x) = \lim_{\epsilon \rightarrow 0} \frac{1}{2\epsilon} \int_0^t I(|Z_r - x| \leq \epsilon) dr. \tag{2.1}$$

These notations will be used subsequently without further explanation.

Assumption **A2** ensures that $w_k, k \geq 1$, is a sequence of stationary random vectors. No restriction is imposed on the relationship between ϵ_k and e_k of $\lambda_k = (\epsilon_k, e_k)'$, which enables the results established here to be widely applicable in non-linear cointegrating regression models with endogeneity, where the components ϵ_k and e_k drive regressor time series and regressor errors, respectively. The extension of **A2** to include linear process formulations is possible if the functional $f(x, y)$ has a certain structure still allowing for endogeneity. We refer to Corollary 2.1 for further details on this extension.

Finally, Assumption **A3** provides conditions on the function $f(x, y)$. These, together with **A2**(b), ensure that,

$$\int_{-\infty}^{\infty} [\mathbb{E}f^2(x, w_1) + \mathbb{E}f^4(x, w_1)] dx \leq C \mathbb{E} \|w_1\|^{\max\{2, 4\beta\}} \int_{-\infty}^{\infty} T(x) dx < \infty, \tag{2.2}$$

the Fourier transform $\hat{f}(t, y) = \int_{-\infty}^{\infty} e^{itx} f(x, y) dx$ is well defined, $\sup_x g(x) < \infty$, $\int |g(x)| dx \leq \int \mathbb{E} |f(x, w_1)| dx < \infty$, and $\int_{-\infty}^{\infty} (1 + |x|) \mathbb{E} |f(x, w_1)| dx < \infty$. Furthermore, it follows from $\mathbb{E}\hat{f}(0, w_1) = \int_{-\infty}^{\infty} \mathbb{E}f(x, w_1) dx = 0$ that

$$|\mathbb{E}\hat{f}(t, w_1)| \leq \int_{-\infty}^{\infty} |(e^{itx} - 1) \mathbb{E}f(x, w_1)| dx \leq C \min\{1, |t|\}. \tag{2.3}$$

On the other hand, using the inverse Fourier transformation, **A3**(III) ensures the representation of $f(x, w_k)$, almost surely,

$$f(x, w_k) = \frac{1}{2\pi} \int_{-\infty}^{\infty} e^{-itx} \hat{f}(t, w_k) dt. \tag{2.4}$$

These properties will be used in the main results that follow without further reference.

2.2. Asymptotic Theory

Our main result is as follows.

THEOREM 2.1. *Suppose **A1–A3** hold. For any $h \equiv h_n \rightarrow 0$ satisfying $nh/d_n \rightarrow \infty$, we have*

$$\begin{aligned} & \left(\frac{d_n}{nh} \sum_{k=1}^{\lfloor nt \rfloor} f^2(x_k/h, w_k), \left(\frac{d_n}{nh} \right)^{1/2} \sum_{k=1}^{\lfloor nt \rfloor} f(x_k/h, w_k) \right) \\ & \Rightarrow (\tau^2 L_Z(t, 0), \tau \mathbb{N} L_Z^{1/2}(t, 0)), \end{aligned} \tag{2.5}$$

on $D_{R^2}[0, 1]$, where $\tau^2 = \int_{-\infty}^{\infty} \mathbb{E}f^2(s, w_1)ds$, and \mathbb{N} is a standard normal variate independent of $L_Z(t, 0)$ for $0 \leq t \leq 1$.

If in addition $\gamma = 0$, where γ is used in **A1(i)**, and $\int_{-\infty}^{\infty} \mathbb{E}\{|\hat{f}(t, w_0)|(1 + \|w_r\|^\beta)\}dt < \infty$ for any $r \geq 0$, then

$$\left(\frac{d_n}{n} \sum_{k=1}^{\lfloor nt \rfloor} f^2(x_k, w_k), \left(\frac{d_n}{n}\right)^{1/2} \sum_{k=1}^{\lfloor nt \rfloor} f(x_k, w_k)\right) \Rightarrow (\tau^2 L_Z(t, 0), \tau_1 \mathbb{N}L_Z^{1/2}(t, 0)), \tag{2.6}$$

on $D_{R^2}[0, 1]$ (recall $Z_t = W(t)$ when $\gamma = 0$), where $\tau_1^2 = G_0 + 2 \sum_{r=1}^{\infty} G_r$ with

$$G_r = \frac{1}{2\pi} \int_{-\infty}^{\infty} \mathbb{E}\{\hat{f}(s, w_0)\hat{f}(s, w_r)e^{-isx_r}\}ds = \int_{-\infty}^{\infty} \mathbb{E}\{f(y, w_0)f(y + x_r, w_r)\}dy. \tag{2.7}$$

Remark 2.1. Different constants τ and τ_1 appear in the second components of results (2.5) and (2.6). In fact, as $h \rightarrow 0$, we have

$$\frac{d_n}{nh} \sum_{k=1}^n \sum_{j=k+1}^n \mathbb{E}\{f(x_k/h, w_k)f(x_{k+j}/h, w_{k+j})\} = o(1),$$

(see the proof of (7.2) in Proposition 7.3); but when $h = 1$ and $\gamma = 0$

$$\frac{d_n}{n} \sum_{k=1}^n f(x_k, w_k)f(x_{k+j}, w_{k+j}) \rightarrow_D G_j L_Z(1, 0), \tag{2.8}$$

for any $j \geq 1$ (see (7.5) of Proposition 7.4). These facts indicate that the influence of cross product terms such as $f(x_k/h, w_k)f(x_{k+j}/h, w_{k+j})$ on the variance of $(\frac{d_n}{nh})^{1/2} \sum_{k=1}^{\lfloor nt \rfloor} f(x_k/h, w_k)$ is eliminated as $h \rightarrow 0$, but this is not the case when $h = 1$. In consequence, different constants appear in the two results (2.5) and (2.6). In addition to (2.6), the following joint convergence holds in which, for any $q > 0$,

$$\left(\frac{d_n}{n} \sum_{k=1}^{\lfloor nt \rfloor} f^2(x_k, w_k), \frac{d_n}{n} \sum_{k=1}^{\lfloor nt \rfloor} f(x_k, w_k)f(x_{k+1}, w_{k+1}), \dots, \frac{d_n}{n} \sum_{k=1}^{\lfloor nt \rfloor} f(x_k, w_k)f(x_{k+q}, w_{k+q}), \left(\frac{d_n}{n}\right)^{1/2} \sum_{k=1}^{\lfloor nt \rfloor} f(x_k, w_k)\right) \Rightarrow (\tau^2 L_Z(t, 0), G_1 L_Z(t, 0), \dots, G_q L_Z(t, 0), \tau_1 \mathbb{N}L_Z^{1/2}(t, 0)), \tag{2.9}$$

on $D_{R^{q+1}}[0, 1]$. The proof of (2.9) involves only minor additions to that of (2.6) and the details are omitted.

Remark 2.2. In special cases where $f(x, y) = K(x)y$ (with $K(x)$ bounded and integrable) and $f(x, y) = K(x)$ (with $\int K(x)dx = 0$ and $K(x)$ bounded and integrable), a similar result to (2.5) has been considered in Wang and Phillips (2009b) and Wang and Phillips (2011), respectively, and a similar result to (2.6) can be found in Jeganathan (2008). Theorem 2.1 provides a unified generalization of these existing results to functional limit theorems. Our proof makes use of the methodology developed in Wang and Phillips (2009b), which seems simpler than that used in Jeganathan (2008).

Remark 2.3. The quantity m_0 given in A2 (a) is set to be a fixed constant, but it can be chosen as large as required in applications. Further, careful examination the proof reveals that the result continues to hold when $m_0 = m_n \rightarrow \infty$ provided the expansion rate is slow enough. Moreover, when $f(x, y) = K(x)y$, the stationary component w_k in Theorem 2.1 can be extended to include linear processes and endogeneity, as the following corollary shows, thereby covering regression models with errors u_t and regressors x_t that allow for endogeneity.

COROLLARY 2.1. *In addition to A1, suppose that:*

- (a) $K(x)$ is a bounded continuous function satisfying $\int K(x)dx < \infty$ and $\int |\hat{K}(x)|dx < \infty$, where $\hat{K}(x) = \int e^{bxs}K(s)ds$.
- (b) $u_k = \sum_{j=0}^{\infty} \psi_j \lambda_{k-j}$, where $\mathbb{E} \lambda_1 = 0$, $\mathbb{E} \|\lambda_1\|^4 < \infty$ and the coefficient vector $\psi_k = (\psi_{k1}, \psi_{k2})$ satisfies $\sum_{k=0}^{\infty} k(|\psi_{1k}| + |\psi_{2k}|) < \infty$ and $\sum_{k=0}^{\infty} \psi_k \neq 0$.

For any $h \equiv h_n \rightarrow 0$ satisfying $nh/d_n \rightarrow \infty$, we have

$$\left(\frac{d_n}{nh} \sum_{k=1}^n K^2(x_k/h) u_k^2, \left(\frac{d_n}{nh} \right)^{1/2} \sum_{k=1}^n K(x_k/h) u_k \right) \rightarrow_D (\tilde{\tau}^2 L_Z(1, 0), \tilde{\tau} \mathbb{N} L_Z^{1/2}(1, 0)), \tag{2.10}$$

where $\tilde{\tau}^2 = \int_{-\infty}^{\infty} K^2(s)ds \mathbb{E} u_1^2$ and \mathbb{N} is a standard normal variate independent of $L_Z(1, 0)$.

If $h = 1$ and in addition $\gamma = 0$, where γ is used in A1(i), then

$$\left(\frac{d_n}{n} \sum_{k=1}^n K^2(x_k) u_k^2, \frac{d_n}{n} \mathcal{J}_n, \left(\frac{d_n}{n} \right)^{1/2} \sum_{k=1}^n K(x_k) u_k \right) \rightarrow_D (\tilde{\tau}^2 L_Z(1, 0), \tilde{\tau}_1^2 L_Z(1, 0), \tilde{\tau}_1 \mathbb{N} L_Z^{1/2}(1, 0)), \tag{2.11}$$

where, for some $M = M_n \rightarrow \infty$,

$$\mathcal{J}_n = \sum_{k=1}^n K^2(x_k) u_k^2 + 2 \sum_{j=1}^M \ell \left(\frac{j}{M} \right) \sum_{k=1}^{n-j} K(x_k) K(x_{k+j}) u_k u_{k+j}, \tag{2.12}$$

takes the form of a heteroskedastic and autocorrelation consistent (HAC) estimator in which $\ell(\frac{j}{M})$ is a lag kernel weight function such as the Bartlett triangular kernel $\ell(\frac{j}{M}) = 1 - \frac{|j|}{M}$, and where $\tilde{\tau}_1^2 = \tilde{G}_0 + 2 \sum_{r=1}^{\infty} \tilde{G}_r$ with

$$\tilde{G}_r = \frac{1}{2\pi} \int_{-\infty}^{\infty} |\hat{K}(s)|^2 \mathbb{E} \{ u_0 u_r e^{-isx_r} \} ds = \int_{-\infty}^{\infty} K(y) \mathbb{E} \{ u_0 u_r K(y + x_r) \} dy.$$

2.3. Self-Normalized Statistics and Discussion

Result (2.10) coincides with (7.4) of Proposition 7.1 in Wang and Phillips (2016) but with less restrictions on h (the requirement $h \log n \rightarrow 0$ used there is removed here), indicating the following self-normalized result: as $h \rightarrow 0$ and $nh/d_n \rightarrow \infty$,

$$J_n(h) := \frac{\sum_{k=1}^n K(x_k/h) u_k}{\sqrt{\sum_{k=1}^n K^2(x_k/h) u_k^2}} \rightarrow_D \mathcal{N}(0, 1). \tag{2.13}$$

In view of the standard normal asymptotics, this result is convenient and useful for purposes of estimation and inference in nonparametric regression models involving nonstationary time series and kernel estimation with a shrinking bandwidth parameter $h \rightarrow 0$, as explained in Section 1.

Result (2.11) with fixed $h = 1$ is similar to that of Theorem 5 in Jeganathan (2008). In this case, a suitable self-normalized version of the sample covariance statistic can be constructed from the elements of (2.11) and (2.12) as

$$J_n^*(1) := \mathcal{J}_n^{-1/2} \sum_{k=1}^n K(x_k) u_k \rightarrow_D \mathcal{N}(0, 1), \tag{2.14}$$

which again has standard normal asymptotics making the formulation convenient in applications that involve nonlinear parametric regressions with nonstationary time series. We mention that the result that $\mathcal{J}_n^2 \rightarrow_D \tilde{\tau}_1^2 L_Z(1, 0)$ holds for any continuous function $\ell(x)$ satisfying $\ell(0) = 1$, although in the present case, it is assumed that $\ell(\frac{j}{M})$ is a lag kernel weight function which ensures the positivity of \mathcal{J}_n in finite samples. Furthermore, we prove (2.11) for some $M_n \rightarrow \infty$. The existence of such an M_n is clear from (6.14) and (6.15) in the proof of Corollary 2.1.

While these naturally constructed self-normalized statistics have elegant Gaussian limit results, numerical work (reported below in Section 3) reveals that neither (2.13) nor (2.14) perform well in finite sample simulations. In particular, when x_r is generated with long memory innovations in ξ_t and the memory parameter is large (μ close to 0.5), bimodality appears in the finite sample densities even when the sample size is as large as $n = 5,000$. Such bimodality is known to arise with self-normalized statistics and t ratios in other contexts, especially in the presence of heavy tailed data where individual large draws can dominate both the numerator and the denominator in the ratio (see Logan et al., 1973; Fiorio,

Hajivassiliou, and Phillips, 2010). The explanation of the phenomena in the present setting is unrelated to heavy tails but is instead related to strong dependence in the data. Heuristically, strong memory when μ is close to 0.5 ensures that the weight function $K(x_k)$ is generally so small that only a limited number of terms dominate the numerator and denominator summations $\sum_{k=1}^n K(x_k)u_k$ and $\sum_{k=1}^n K^2(x_k)u_k^2$ (see Figure 4 for illustrative trajectories), thereby inducing bimodality in the finite sample densities of $\mathcal{J}_n^*(1)$ around the modes ± 1 . To control this behavior, a modification of (2.14) such as the following:

$$\widehat{\mathcal{J}}_n^*(1) := \widehat{\mathcal{J}}_n^{-1/2} \sum_{k=1}^n K(x_k)u_k \rightarrow_D \mathcal{N}(0, 1), \tag{2.15}$$

might be considered where \mathcal{J}_n in (2.12) is replaced by

$$\widehat{\mathcal{J}}_n = \widehat{\sigma}_n^2 \sum_{k=1}^n K^2(x_k) + 2 \sum_{j=1}^M \ell\left(\frac{j}{M}\right) \sum_{k=1}^{n-j} K(x_k)K(x_{k+j})u_k u_{k+j}, \tag{2.16}$$

for some consistent estimator $\widehat{\sigma}_n^2$ of $\sigma^2 = \mathbb{E}u_1^2$ and with $M = M_n \rightarrow \infty$ as $n \rightarrow \infty$. The advantage of $\widehat{\mathcal{J}}_n$ is that the use of $\widehat{\sigma}_n^2 \sum_{k=1}^n K^2(x_k)$ in the first term, rather than $\sum_{k=1}^n K^2(x_k)u_k^2$, attenuates the bimodality induced by the numerator and denominator summations $\sum_{k=1}^n K(x_k)u_k$ and $\sum_{k=1}^n K^2(x_k)u_k^2$ discussed above and in the heuristic analysis of (3.4). However, the estimate $\widehat{\mathcal{J}}_n$ in (2.16) is not necessarily positive. For instance, in 40,000 replications when $n = 100$ around 15 cases of negative values occur with $d = 0.1$, rising to 60 cases with $d = 0.55$. To address this difficulty, the following adjustment to (2.16) is employed

$$\widehat{\mathcal{J}}_{nM^*} = \widehat{\sigma}_n^2 \sum_{k=1}^n K^2(x_k) + 2 \sum_{j=1}^{M^*} \ell\left(\frac{j}{M}\right) \sum_{k=1}^{n-j} K(x_k)K(x_{k+j})u_k u_{k+j}, \tag{2.17}$$

where

$$M^* := M \times \mathbb{I}(\widehat{\mathcal{J}}_n \geq 0) + M^* \times \mathbb{I}(\widehat{\mathcal{J}}_n < 0) \mathbb{I}(\widehat{\mathcal{J}}_{nM^*} > 0), \tag{2.18}$$

in which the truncation lag number M is reduced by one lag at a time when $\widehat{\mathcal{J}}_n < 0$ to the first value M^* for which $\widehat{\mathcal{J}}_{nM^*} > 0$. In 50,000 replications with $n = 100$ and $n = 1,000$, the modification (2.17), with the simple rule (2.18), was found to work well. Using $\widehat{\mathcal{J}}_{nM^*}$ in place of $\widehat{\mathcal{J}}_n$ leads to the same standard normal asymptotics as (2.15) for the statistic

$$\widetilde{\mathcal{J}}_n(1) := \widehat{\mathcal{J}}_{nM^*}^{-1/2} \sum_{k=1}^n K(x_k)u_k \rightarrow_D \mathcal{N}(0, 1), \tag{2.19}$$

provided $M^* \rightarrow \infty$ as $n \rightarrow \infty$. Simulation results for the statistic $\tilde{J}_n(1)$ are shown in Figure 3 in the following numerical section and confirm that the statistic removes bimodality in finite samples and has distributions considerably closer to the standard normal limit than the statistic $J_n^*(1)$ in (2.14) for various values of the long memory parameter d and samples as small as $n = 100$.

Similarly, we may use the following result instead of (2.13): as $h \rightarrow 0$ and $nh/d_n \rightarrow \infty$,

$$\hat{J}_n(h) := \frac{\sum_{k=1}^n K(x_k/h)u_k}{\sqrt{\hat{\sigma}_n^2 \sum_{k=1}^n K^2(x_k/h)}} \rightarrow_D \mathcal{N}(0, 1). \tag{2.20}$$

The proofs of (2.15) and (2.20) follow easily from (2.14), (2.13) and the following fact by using (4.8) of Wang et al. (2021) [see also (7.42) in the proof of Proposition 7.4 with $f(x, y) = K(x)y$]: for any $h > 0$,

$$\frac{d_n}{nh} \sum_{k=1}^n K^2(x_k/h) (\mathbb{E}u_k^2 - u_k^2) = o_P(1). \tag{2.21}$$

The details are omitted.

3. NUMERICAL EVIDENCE

We explore the finite sample properties of the self-normalized statistics J_n and $J_n^*(1)$ defined as in (2.13) and (2.14). Since earlier research has considered models with shrinking bandwidths $h \rightarrow 0$, the model employed here focuses mainly on the case $h = 1$ for which the general limit theory is given in (2.9). As indicated above, the key difference in this case is that the cross product term (2.8) is not eliminated when $h \not\rightarrow 0$. The statistic $J_n^*(1)$ takes this into account by estimating the appropriate self-normalizing quantity. As is apparent from (2.9) and (2.11), the limiting form of the denominator of $J_n^*(1)$ has the form of a long run self-normalization, with the major difference that in the present case, this quantity has a random limit since $\mathcal{J}_n \rightarrow \tilde{\tau}_1^2 L_Z(1, 0)$ as $n \rightarrow \infty$ in place of the usual nonrandom quantity that arises in standard problems with stationary short memory time series.

In the simulations here, x_t is generated according to **A1** with autoregressive coefficient $\rho_n = 1$. The linear process $\xi_t = \sum_{j=0}^{\infty} \phi_j \epsilon_{t-j}$ in LM is generated using the fractional integration mechanism $\xi_t = (1 - L)^{-d} \epsilon_t = \sum_{j=0}^{\infty} \frac{(d)_j}{j!} \epsilon_{t-j}$, where $(d)_j = \frac{\Gamma(d+j)}{\Gamma(1+j)}$, so that $\phi_j \sim \frac{1}{\Gamma(d)j^{1-d}}$, where $\Gamma(\cdot)$ is the gamma function and the memory parameter $d = 1 - \mu \in (0, 0.5)$. Endogeneity in x_t is introduced by defining the innovations in the linear process ξ_t by $\epsilon_t = (1 - \rho^2)^{1/2} \epsilon_{xt} + \rho u_t$ where u_t is the short memory autoregressive process $u_t = \theta u_{t-1} + e_{ut}$, $|\theta| < 1$, with $e_{ut} \sim_{iid} \mathcal{N}(0, 1)$ and independent of $\epsilon_{xt} \sim_{iid} \mathcal{N}(0, 1)$. With this specification of u_t , we have

$$\begin{aligned}
 \xi_t &= \sum_{j=0}^{\infty} \phi_j \epsilon_{t-j} = (1 - \rho^2)^{1/2} \sum_{k=0}^{\infty} \phi_k \epsilon_{xt-k} + \rho \sum_{j=0}^{\infty} \phi_j \sum_{\ell=0}^{\infty} \theta^\ell \epsilon_{ut-j-\ell} \\
 &= (1 - \rho^2)^{1/2} \sum_{k=0}^{\infty} \phi_k \epsilon_{xt-k} + \rho \sum_{k=0}^{\infty} \left(\sum_{\ell=0}^k \phi_{k-\ell} \theta^\ell \right) \epsilon_{ut-k} \\
 &= \sum_{k=0}^{\infty} \left[\bar{\psi}_{1k} \epsilon_{xt-k} + \bar{\psi}_{2k} \epsilon_{ut-k} \right]
 \end{aligned} \tag{3.1}$$

with $\bar{\psi}_{1k} = (1 - \rho^2)^{1/2} \phi_k$ and $\bar{\psi}_{2k} = \rho \sum_{\ell=0}^k \phi_{k-\ell}$. The innovation ξ_t has long memory parameter d and endogeneity measured through the correlation coefficient ρ .

The self-normalized statistics $J_n(h)$, $J_n(1)$, and $J_n^*(1)$ defined in (2.13) and (2.14) are computed for $f(x_t/h, w_t) = K(x_t/h)u_t$ with $h = 2/n^{0.2}$ or $h = 1$. In the following computations, we used $K(x) = (1/\sqrt{2\pi})e^{-x^2/2}$, $\theta = 0.5$, $\rho = 5.0$, and $d \in \{0.1, 0.25, 0.4, 0.55\}$, where $d = 0.55$ lies in the nonstationary long memory region and is included for comparison. Kernel estimates of the densities of $J_n(h)$ were computed using

$$J_n(h) = \frac{\sum_{k=1}^n K(x_k/h) u_k}{\sqrt{\sum_{k=1}^n K^2(x_k/h) u_k^2}}, \tag{3.2}$$

for $h = 2/n^{0.2}$ and $h = 1$ and are shown in Figures 1(a) and 1(b). The self-normalized statistic $J_n^*(1)$ was computed by the explicit formula

$$J_n^*(1) = \frac{\sum_{k=1}^n K(x_k) u_k}{\left[\sum_{k=1}^n K^2(x_k) u_k^2 + 2 \sum_{j=1}^M \ell\left(\frac{j}{M}\right) \sum_{k=1}^{n-j} K(x_k) K(x_{k+j}) u_k u_{k+j} \right]^{1/2}} \tag{3.3}$$

with lag truncation parameter $M = \lfloor 2n^{1/6} \rfloor$ and its densities are shown in Figures 2(b) and 2(c). The number of replications employed was 40,000, with sample size $n = 100$ in Figure 1 and $n = 1,000$ in Figure 2.

The densities in Figure 1 where $n = 100$ are all non-normal. Bimodality with modes around ± 1 are clearly evident in all cases and all values of d . For $J_n(1)$, the dual modes are evident but somewhat less pronounced than for $J_n(h)$ with $h = 2/n^{0.2}$. The bimodality is clearly stronger in the presence of nonstationary long memory innovations ξ_t with $d = 0.55$ (shown by dashed green lines). Bimodality is most prominent and with greatest concentration for the statistic $J_n^*(1)$. Bimodality is evidently weaker for the lower memory parameters, particularly cases where $d = 0.10$ (shown by black unbroken lines).

In Figure 2, the densities are computed for $n = 1,000$. In Figure 2(a), bimodality is clearly evident for $J_n(h)$, applies for all values of d and is again stronger in the

nonstationary case. The densities of $J_n(1)$ and $J_n^*(1)$ in Figures 2(b) and 2(c), where $n = 1,000$, are closer to normal than when $n = 100$ except for the nonstationary innovation case ($d = 0.55$); and bimodality is still more pronounced for $J_n^*(1)$ than for $J_n(1)$. When $d = 0.1$, there are no apparent modes in the density of $J_n(1)$ and only minor modes in the density of $J_n^*(1)$. Nonetheless, convergence to normality when $0 < d < 0.5$ appears slow and shape differences in the densities persist between the stationary and nonstationary error cases. The tendency to bimodality continues to be more marked in the nonstationary case.

As discussed in Section 2.3, when the innovations ξ_k have strong dependence with memory parameter d close to the nonstationary boundary 0.5, the weight function $K(x_t)$ is negligible except for a very small number of terms in which $x_t = \sum_{k=1}^t \xi_k \approx 0$. Suppose x_t is closest to zero for $t = \tau$, then $K(x_\tau) \approx 1$ and so $J_n(1) \approx \pm 1$, thereby inducing a tendency to bimodality in the finite sample densities of $J_n(1)$ around modes at ± 1 . When $h \rightarrow 0$, this facet of the weight function is accentuated for $K(x_t/h)$ and we may therefore expect greater evidence of bimodality in finite samples for $J_n(h)$, which is corroborated by the results in Figures 1(a) and 2(b).

Further, in Figures 1 and 2, it is evident that $J_n^*(1)$ shows more evidence of bimodality than $J_n(1)$. This may be explained by the following heuristic. Suppose x_t is closest to zero in the sample at $t = \tau$ and next closest to zero at $t = \tau + 1$, so that $K(x_\tau) \approx K(0) \approx 1/\sqrt{2\pi}$ and then $K(x_{\tau+1}) \approx K(\xi_{\tau+1}) = e^{-\xi_{\tau+1}^2/2}/\sqrt{2\pi}$ (Figure 4 shows an illustrative case). With a Bartlett kernel $\ell(\cdot)$, we then have

$$\begin{aligned}
 J_n^*(1) &\approx \frac{K(x_\tau)u_\tau + K(x_{\tau+1})u_{\tau+1}}{[K(x_\tau)^2u_\tau^2 + K(x_{\tau+1})^2u_{\tau+1}^2 + 2(1 - \frac{1}{M})K(x_\tau)K(x_{\tau+1})u_\tau u_{\tau+1}]^{1/2}} \\
 &= \frac{K(x_\tau)u_\tau + K(x_{\tau+1})u_{\tau+1}}{|K(x_\tau)u_\tau + K(x_{\tau+1})u_{\tau+1}| + O_p(\frac{1}{M})} = \pm 1 + O_p\left(\frac{1}{M}\right),
 \end{aligned}
 \tag{3.4}$$

showing a clear tendency to bimodality.




FIGURE 1. Empirical densities of $J_n(h)$ with $h = \frac{2}{n^{0.2}}$, $J_n(1)$, and $J_n^*(1)$ for sample size $n = 100$ and $d \in \{0.10, 0.25, 0.40, 0.55\}$.




FIGURE 2. Empirical densities of $J_n(h)$, $J_n(1)$, and $J_n^*(1)$ for sample size $n = 1,000$ and $d \in \{0.10, 0.25, 0.40, 0.55\}$.




FIGURE 3. Empirical densities of $\tilde{J}_n(1)$ for sample sizes $n = 100$ and for $n = 1,000$ and $n = 5,000$ and $d \in \{0.10, 0.25, 0.40, 0.55\}$.

Next, note that $\xi_t = (1 - L)^{-d}\epsilon_t$ has variance $\sigma_\xi^2 = \sigma_\epsilon^2 \frac{\Gamma(1-2d)}{\Gamma(1-d)^2} \sim_a \frac{\sigma_\epsilon^2/\pi}{1-2d} \rightarrow \infty$ as $d \rightarrow 0.5$. Let $\xi_t = \sigma_\xi \tilde{\xi}_t$, where $\tilde{\xi}_t$ has unit variance. Then $K(x_{\tau+1}) \approx K(\xi_{\tau+1}) = e^{-\sigma_\xi^2 \tilde{\xi}_{\tau+1}^2 / \sqrt{2\pi}}$ and

$$\begin{aligned}
 J_n(1) &\approx \frac{K(x_\tau)u_\tau + K(x_{\tau+1})u_{\tau+1}}{[K(x_\tau)^2u_\tau^2 + K(x_{\tau+1})^2u_{\tau+1}^2]^{1/2}} \approx \frac{u_\tau + e^{-\sigma_\xi^2 \tilde{\xi}_{\tau+1}^2} u_{\tau+1}}{[u_\tau^2 + e^{-\sigma_\xi^2 \tilde{\xi}_{\tau+1}^2}]^{1/2}} \approx \frac{u_\tau}{|u_\tau|} + O_p\left(e^{-\sigma_\xi^2}\right) \\
 &\approx \pm 1 + O_p\left(\frac{1}{1-2d}\right), \tag{3.5}
 \end{aligned}$$

showing a tendency to bimodality as the memory parameter $d \rightarrow 0.5$. The same tendency to bimodality is also present in the approximation of $J_n^*(1)$ in addition to that given in (3.4), thereby implying that $J_n^*(1)$ is more likely to manifest bimodal behavior in finite samples than $J_n(1)$, corroborating the simulation findings.

Figure 3 shows finite sample densities of the statistic $\tilde{J}_n(1)$ in (2.17) using the same simulation design with the same set of long memory parameters, endogeneity correlation $\rho = 0.5$, and for sample sizes increasing from $n = 100$ to $n = 5,000$ based on 40,000 replications. As evident in the graphics, the statistic removes bimodality in finite samples although there are extended shoulders on either side

of the origin to around ± 1 , particularly when $n = 100$. The distributions are far closer to the standard normal limit than those of the statistic $J_n^*(1)$ in (2.14) at every sample size with evident convergence in shape to normal for all values of the long memory parameter and clearest for $d = 0.1$, as would be expected. These findings support the heuristic analysis leading to (3.4) and (3.5). For when the variance estimate $\widehat{\mathcal{J}}_{nM^*}$ is employed, the scaling-out effect that leads to bimodality is removed, thereby explaining the finite sample distributions being closer to the standard normal.

4. FURTHER ANALYSIS: FINITE SAMPLE AND ASYMPTOTIC BIMODALITY

As noted in Section 2.3, natural self-normalization of sample covariance statistics does not perform well in finite samples relative to the asymptotic theory when strong effects of long memory are present in the data. This result in nonlinear nonparametric regression is new to the literature. But the observed finite sample bimodality has a subtle connection in its origins with earlier findings on bimodal t ratios where behavior is dominated by a few observations when there is heavy tailed data. In the present case, behavior is dominated by the few neighboring observations whose impact is not diminished by the kernel weights under strong dependence. Figure 4 illustrates with a single shot picture of typical data trajectories generated for x_t and u_t with $d = 0.1$ and $n = 1,000$.

Some additional analysis and computations are now provided to shed light on the finite sample properties of self-normalized sample covariance statistics in which nonstationarity originates in partial sums of long memory processes. The following




FIGURE 4. Single shot trajectories of x_t and u_t generated with $d = 0.10$ and $n = 1,000$ according to the simulation design given below.

simple framework with no endogeneity is used for the following discussion and data generation.

Simulation design

- Both ϵ_k and u_k are iid $\mathcal{N}(0, 1)$ and the ϵ_k are independent of the u_k .
- $x_k = \sum_{j=1}^k \xi_j$, where $(1 - L)^d \xi_j = \epsilon_j$, with $0 < d < 1/2$ and $1/2 < \mu = 1 - d < 1$, so that $\xi_j = (1 - L)^{-d} \epsilon_j = \sum_{i=0}^{\infty} \phi_i \epsilon_{j-i}$ with $\phi_i \sim \frac{1}{\Gamma(d)} i^{-(1-d)}$.
- $K(x) = e^{-x^2/2} / \sqrt{2\pi}$.

For $j = 1$ and 2 , define

$$S_{jn} = \mathcal{J}_{jn}^{-1/2} \sum_{k=1}^n K(x_k) u_k,$$

where $\mathcal{J}_{1n} = \sum_{k=1}^n K^2(x_k)$ and $\mathcal{J}_{2n} = \sum_{k=1}^n K^2(x_k) u_k^2$. Under these conditions, ξ_k is a long memory process with memory parameter $0 < d = 1 - \mu < 1/2$ and x_k is nonstationary with memory parameter $1 + d$. S_{2n} is a natural self-normalized sample covariance statistic, matching $J_n^*(1)$ in (2.14).⁵

Recall that $d_n^2 = \text{var}(x_n) \sim A_d n^{1+2d}$, where A_d is a positive constant depending only on d . It is readily seen from (2.11) and (2.21) that

$$\begin{aligned} \frac{1}{n^{1/2-d}} \mathcal{J}_{1n}, \quad \frac{1}{n^{1/2-d}} \mathcal{J}_{2n} &\rightarrow_D \left(\frac{A_d}{2}\right)^{1/2} L_{B_{(1+2d)/2}}(1, 0), \\ \frac{\mathcal{J}_{2n} - \mathcal{J}_{1n}}{\mathcal{J}_{1n}} &\rightarrow_P 0, \end{aligned} \tag{4.1}$$

where $B_H(t)$ is fractional Brownian motion with Hurst exponent H and $L_{B_H}(t, s)$ is the local time process of $\{B_H(t)\}_{t \geq 0}$. In view of the independence of x_k and u_k and since $u_k \sim_{iid} \mathcal{N}(0, 1)$, we have $S_{1n} \sim_d \mathcal{N}(0, 1)$ for all $n \geq 1$ and

$$S_{2n} = \left(\frac{\mathcal{J}_{1n}}{\mathcal{J}_{2n}}\right)^{1/2} S_{1n} \rightarrow_D \mathcal{N}(0, 1), \tag{4.2}$$

so that S_{2n} has a standard normal limit distribution. Now consider the finite sample performance of the statistics S_{1n} and S_{2n} .

A. Simulation results for S_{1n} : Kernel density estimates of the finite sample distributions of S_{1n} are shown in Figure 5(a) for sample size $n = 100$ with $d \in \{0.1, 0.25, 0.4, 0.55\}$ from 40,000 replications. The graphs confirm the exact finite sample $\mathcal{N}(0, 1)$ distribution for all values of the memory parameter d , including the nonstationary case $d = 0.55$.

B. Simulation results for S_{2n} : Figure 5(b) shows the finite sample densities of S_{2n} for $n = 100$ and same memory parameter values $d \in \{0.1, 0.25, 0.4, 0.55\}$ again from 40,000 replications. Bimodality in these distributions around the points ± 1 is clearly evident for all $d > 0.10$ and strong in the nonstationary case $d = 0.55$;

⁵When ϵ_k and x_k are independent of u_k the term $2 \sum_{j=1}^M \ell(\frac{j}{M}) \sum_{k=1}^{n-j} K(x_k) K(x_{k+j}) u_k u_{k+j}$ that is included in \mathcal{J}_n is unnecessary since the terms \tilde{G}_r appearing in Corollary 2.1 are zero for all $r \geq 1$.




FIGURE 5. Empirical densities of S_{1n} and S_{2n} for $n = 100$, $d \in \{0.1, 0.25, 0.4, 0.55\}$.




FIGURE 6. Empirical densities of S_{2n} for sample sizes $n = 1,000$ and $n = 5,000$ and $d \in \{0.1, 0.25, 0.4, 0.55\}$.

for $d = 0.10$ the density has shoulders at the same points ± 1 . Figures 6(a) and 6(b) show the corresponding densities for $n = 1,000$ and $n = 5,000$. The slow convergence of these distributions to normality in the presence of stationary long memory is evident, especially for $d = 0.4$ where shoulders in the density around ± 1 are evident even when $n = 5,000$. In the nonstationary $d = 0.55$ case, bimodality remains evident, although it is not as strong as it is for smaller sample sizes.

Although S_{2n} has a normal limit distribution for all memory parameters $d \in (0, 0.5)$, the finite sample performance of S_{2n} depends on the value of d , in contrast to S_{1n} . Bimodality is strongest for stationary values of d closest to the boundary $d = 0.5$ and remains present even for very large sample sizes. This anomalous behavior can be explained in terms of relative convergence rates as follows. Recalling (4.1), when $d = 0.4$, we have

$$\left(\frac{\mathcal{J}_{1n}}{\mathcal{J}_{2n}}\right)^{1/2} - 1 = \frac{\mathcal{J}_{1n} - \mathcal{J}_{2n}}{\mathcal{J}_{2n}^{1/2}(\mathcal{J}_{1n}^{1/2} + \mathcal{J}_{2n}^{1/2})} = O_p(n^{-0.05}),$$




FIGURE 7. Empirical densities of S_{2n} for sample sizes $n = 100, 1,000$ and $n = 5,000$ and $d \in \{0.75, 1.00\}$.

whence $\mathcal{J}_{2n}/\mathcal{J}_{1n} \rightarrow_P 1$ as $n \rightarrow \infty$; but the convergence rate is seen to be very slow. With such a slow convergence rate, even for $n = 5,000$ (where $n^{-0.05} \approx 0.65$) and with $S_{1n} \sim_d \mathcal{N}(0, 1)$ for all $n \geq 1$, the value of $S_{2n} = (\frac{\mathcal{J}_{1n}}{\mathcal{J}_{2n}})^{1/2} S_{1n}$ can be substantially impacted by the factor $(\frac{\mathcal{J}_{1n}}{\mathcal{J}_{2n}})^{1/2}$, leading to departures from the normality of S_{1n} and the presence of bimodality in the distribution.

When $x_k = \sum_{j=1}^k \xi_j$ with $(1 - L)^d \xi_j = \epsilon_j$ and $d > 1/2$, the input ξ_j is a nonstationary long memory process and the limit distribution S_{2n} is not normal. In fact, bimodality must appear in this case and we have

$$\mathcal{J}_{1n} \rightarrow_P A := \sum_{k=1}^{\infty} K^2(x_k), \quad \mathcal{J}_{2n} \rightarrow_P B := \sum_{k=1}^{\infty} K^2(x_k) u_k^2, \tag{4.3}$$

where A and B ($A \neq B$) are well-defined positive random variables. Hence, as $n \rightarrow \infty$,

$$S_{2n} = \left(\frac{\mathcal{J}_{1n}}{\mathcal{J}_{2n}}\right)^{1/2} S_{1n} \rightarrow_D \left(\frac{A}{B}\right)^{1/2} \mathcal{N}(0, 1), \tag{4.4}$$

since $S_{1n} \sim \mathcal{N}(0, 1)$ for all $n \geq 1$. The presence of the ratio A/B of the random variables (A, B) assures bimodality in the limit distribution (4.4).

The proof of (4.3) and (4.4) is straightforward. Let $A_{m,n} = \sum_{k=m}^n K^2(x_k)$ and recall that $x_n \sim_d \mathcal{N}(0, d_n)$, where $d_n^2 = \text{var}(x_n) \sim_a A_d n^{1+2d}$ as $n \rightarrow \infty$, it is readily seen that, whenever $d > 1/2$ and $m, n \rightarrow \infty$,

$$\begin{aligned} \mathbb{E}A_{m,n} &= \sum_{k=m}^n \mathbb{E}K^2(x_k) = \sum_{k=m}^n \int K^2(d_k y) e^{-y^2/2} dy \\ &\leq C \sum_{k=m}^n d_k^{-1} = C_1 \sum_{k=m}^n k^{-(1+2d)/2} \rightarrow 0. \end{aligned}$$

Hence, $A := \sum_{k=1}^{\infty} K^2(x_k)$ is a well-defined random variable and $\mathcal{J}_{1n} \rightarrow_P A$. Similarly, we have $\mathbb{E}B_{m,n} \rightarrow 0$ where $B_{m,n} = \sum_{k=m}^n K^2(x_k) \eta_k^2$, and hence $\mathcal{J}_{2n} \rightarrow_P B$.

Figure 7 gives simulation results for S_{2n} in the nonstationary innovation cases $d = 0.75$ and $d = 1$ for $n = 100, 1,000,$ and $5,000$ based on 25,000 replications. Bimodality appears a prominent feature of the densities of S_{2n} for both $d = 0.75$ and $d = 1$, showing little tendency to diminish even in very large sample sizes, corroborating the non-Gaussian limit theory in the nonstationary case. The bimodality is stronger when $d = 1$ than when $d = 0.75$ for all sample sizes.

5. CONCLUDING REMARKS

Sample covariance functionals of regressors and innovations play a key role in nonlinear nonstationary regression models and self-normalized versions of these statistics are a foundation for inference. The limit theory given here covers a wide class of such functionals and reveals important differences between stationary and nonstationary long memory innovations. Methods involving bandwidths $h = h_n \rightarrow 0$ in nonparametric models and fixed $h = 1$ suited for parametric applications are jointly included in the present findings. Numerical work shows strong bimodality in the finite sample distributions, slow convergence to the Gaussian limit theory under stationary long memory innovations and non-Gaussian limit theory when the innovations have nonstationary long memory. New forms of self-normalization are shown to provide the same limit theory but improved finite sample performance suitable for practical work in these difficult cases.

It is of interest to explore the performance of this modified form of self-normalization in regression test applications. Bimodality, when induced by self-normalization as in the cases considered here, typically leads to the presence of modes around ± 1 (Logan et al., 1973; Fiorio et al., 2010). The general impact of such bimodality is to transfer extreme tail probability in the distribution towards the modes, which in turn, typically makes testing somewhat conservative in applications and this is inclined to reduce power in testing under local alternatives when using nominal asymptotic critical values. We might therefore expect some such impact in the present examples with long memory innovations. The new form of self-normalization introduced here is designed to attenuate such effects and an investigation of the size/power implications of this modification in regression applications is a topic for future research.

The present framework, in conjunction with earlier findings in the nonstationary nonlinear regression literature, can be extended to cover a wider class of models than already discussed. One such model is a nonlinear distributed lag cointegrating regression of the following additive nonparametric type $y_k = g(x_k) + \sum_{j=1}^J g_j(\Delta x_{k-j}) + u_k$, where the $I(1)$ regressor x_k is nonlinearly related to y_k with additive and nonlinear distributed lag effects from the regressors $\{\Delta x_{k-j} : j = 1, \dots, J\}$. In such models, the cointegrating function $g(x_k)$ is usually of primary interest. If the additive component $\sum_{j=1}^J g_j(\Delta x_{k-j})$ were ignored and instead absorbed into the primary component, the equation $y_k = \mu + g(x_k) + v_k = g_\mu(x_k) + v_k$ may be consistently estimated by kernel methods. Indeed, with some modification, the results and limit theory of Wang and Phillips (2009b) would

continue to hold in such cases because they cover regressions with an endogenous regressor x_k correlated with a stationary error such as v_k . If the g_j are measurable, integrable functions and Δx_k is stationary, then setting $\mu = \sum_{j=1}^J \mathbb{E}g_j(\Delta x_{k-j})$ and $v_k = u_k + \sum_{j=1}^J (g_j(\Delta x_{k-j}) - \mathbb{E}g_j(\Delta x_{k-j}))$, estimation and inference concerning $g_\mu(x_k)$ in the system $y_k = g_\mu(x_k) + v_k$ can be justified as in Wang and Phillips (2009b) under some extension of the underlying conditions to accommodate the properties of the induced error process v_k . Full exploration of this and related extensions is left for future research.

6. PROOFS OF THE MAIN RESULTS

Proof of Theorem 2.1. First note that, for any bounded $h > 0$ and $nh/d_n \rightarrow \infty$,

$$\left(\frac{d_n}{nh}\right)^{1/2} \max_{1 \leq k \leq n} |f(x_k/h, w_k)| = o_p(1), \tag{6.1}$$

by a similar argument as in Proposition 7.4.⁶ Due to (6.1), without loss of generality, we assume

$$f(x_k/h, w_k) = 0 \quad \text{for } k = 1, \dots, A_0, \tag{6.3}$$

where A_0 is a fixed constant that can be chosen large enough. This convention will reduce notational complexity in the proofs of propositions that are given in Section 7 and the lemmas in the Appendix.

We adopt the methodology employed in Wang and Phillips (2009b), starting with an outline of the proof of (2.6), where some useful propositions will be given in Section 7. Define, for $0 \leq t \leq 1$,

$$S_n(t) = \left(\frac{d_n}{n}\right)^{1/2} \sum_{k=1}^{\lfloor nt \rfloor} f(x_k, w_k),$$

$$Y_{nq}(t) = \psi_{n0}(t) + 2 \sum_{j=1}^q \psi_{nj}(t),$$

⁶Indeed, as in (7.4) of Proposition 7.4, it follows from $nh/d_n \rightarrow \infty$ that, for any $A > 0$,

$$\begin{aligned} & \left(\frac{d_n}{nh}\right)^{1/2} \max_{1 \leq k \leq n} |f(x_k/h, w_k)| \\ & \leq \left[\frac{d_n}{nh} \sum_{k=1}^n f^2(x_k/h, w_k) I(|f(x_k/h, w_k)| \geq A) \right]^{1/2} + A \left(\frac{d_n}{nh}\right)^{1/2} \\ & \rightarrow_D \left[\int_{-\infty}^{\infty} \mathbb{E}f^2(x, w_1) I(|f(x, w_1)| \geq A) dx L_Z(1, 0) \right]^{1/2}, \quad \text{as } n \rightarrow \infty. \end{aligned} \tag{6.2}$$

This implies (6.1) since $\int_{-\infty}^{\infty} \mathbb{E}f^2(x, w_1) I(|f(x, w_1)| \geq A) dx \leq A^{-2} \int_{-\infty}^{\infty} \mathbb{E}f^4(x, w_1) dx \rightarrow 0$ by (2.2), as $A \rightarrow \infty$.

where for $j = 0, 1, \dots, q$,

$$\psi_{nj}(t) = \frac{d_n}{n} \sum_{k=1}^{\lfloor nt \rfloor} f(x_k, w_k) f(x_{k+j}, w_{k+j}),$$

and for all $\alpha_i, \beta_j \in \mathbb{R}$, $0 \leq s_0 < s_1 < \dots < s_m < \infty$ and $0 \leq t_0 < t_1 < \dots < t_l < \infty$,

$$Z_{n2} = \sum_{i=1}^l \alpha_i [\zeta_{n1}(t_i) - \zeta_{n1}(t_{i-1})] + \sum_{i=1}^m \beta_i [\zeta_{n2}(s_i) - \zeta_{n2}(s_{i-1})],$$

where $\zeta_{n1}(t) = \frac{1}{\sqrt{n}} \sum_{j=1}^{\lfloor nt \rfloor} \epsilon_j$ and $\zeta_{n2}(t) = \frac{1}{\sqrt{n}} \sum_{j=1}^{\lfloor nt \rfloor} \epsilon_{-j}$. An application of Proposition 7.4 implies that, for any $q \geq 1$,

$$(\psi_{n0}, \psi_{n1}, \dots, \psi_{nq}, Y_{nq}(t)) \Rightarrow (G_0, G_1, \dots, G_q, \Lambda_q) L_Z(t, 0), \tag{6.4}$$

on $D_{R^{q+2} \times 0, 1}$, where $\Lambda_q = G_0 + 2 \sum_{r=1}^q G_r$. This, together with the tightness of $\{S_n(t)\}_{n \geq 1}$ (see Proposition 7.2 with $h = 1$), yields

$$\{S_n(t), Y_{nq}(t), Z_{n2}\}_{n \geq 1} \text{ is tight on } D_{\mathbb{R}^3}[0, 1]. \tag{6.5}$$

Hence, for each $\{n'\} \subseteq \{n\}$, there exists a subsequence $\{n''\} \subseteq \{n'\}$ such that

$$\{S_{n''}(t), Y_{n''q}(t), Z_{n''2}\} \Rightarrow \{\eta(t), \Lambda_q L_Z(t, 0), Z_2\}, \tag{6.6}$$

on $D_{\mathbb{R}^3}[0, 1]$, where

$$Z_2 = \sum_{i=1}^l \alpha_i (B_{1t_i} - B_{1t_{i-1}}) + \sum_{i=1}^m \beta_i (B_{2s_i} - B_{2s_{i-1}}),$$

and $\eta(t)$ is a process continuous with probability one due to (6.1).

Let $Z_{n3} = \sum_{i=1}^v \gamma_i [S_n(t_i) - S_n(t_{i-1})]$ and $Z_3 = \sum_{i=1}^v \gamma_i [\eta(t_i) - \eta(t_{i-1})]$, where $\gamma_j \in \mathbb{R}$ and $0 \leq t_0 < t_1 < \dots < t_v \leq s$. Since, for each $0 \leq t \leq 1$, $S_n(t)$ is uniformly integrable (see Proposition 7.1 with $h = 1$), it follows from Proposition 7.3(i) with $h = 1$ that, for any $s < t$,

$$\begin{aligned} & \mathbb{E} e^{i(Z_3 + Z_2)} [\eta(t) - \eta(s)] \\ &= \lim_{n'' \rightarrow \infty} \mathbb{E} e^{i(Z_{n''3} + Z_{n''2})} [S_{n''}(t) - S_{n''}(s)] = 0 \end{aligned} \tag{6.7}$$

(see, e.g., Billingsley, 1968, Thm. 5.4). Similarly, by Propositions 7.1 with $h = 1$ and 7.3(iii) with $h = 1$, we have

$$\mathbb{E} e^{i(Z_3 + Z_2)} \{[\eta(t) - \eta(s)]^2 - [Y(t) - Y(s)]\} = 0, \tag{6.8}$$

where $Y(t) = \tau_1^2 L_Z(t, 0)$. Indeed, by letting $Y_q(t) = \Lambda_q L_Z(t, 0)$ and noting

$$\sup_{0 \leq t \leq 1} \mathbb{E} |Y_q(t) - Y(t)| \leq 2 |\Lambda_q - \tau_1^2| E \sup_{0 \leq t \leq 1} L_Z(t, 0) \leq C \sum_{r=q+1}^{\infty} |G_r| \rightarrow 0,$$

due to Proposition 7.5, it follows from Propositions 7.1 with $h = 1$ and 7.3(iii) with $h = 1$ that, for any $\epsilon > 0$,

$$\begin{aligned}
 & \left| \mathbb{E} e^{i(Z_3+Z_2)} \{[\eta(t) - \eta(s)]^2 - [Y(t) - Y(s)]\} \right| \\
 & \leq \left| \mathbb{E} e^{i(Z_3+Z_2)} \{[\eta(t) - \eta(s)]^2 - [Y_q(t) - Y_q(s)]\} \right| \\
 & \quad + \mathbb{E} \left| [Y_q(t) - Y(t)] \right| + \mathbb{E} \left| [Y_q(s) - Y(s)] \right| \\
 & \leq \lim_{n'' \rightarrow \infty} \left| \mathbb{E} e^{i(Z_{n''3}+Z_{n''2})} \{[S_{n''}(t) - S_{n''}(s)]^2 - [Y_{n''q}(t) - Y_{n''q}(s)]\} \right| + 2\epsilon \\
 & \leq 3\epsilon,
 \end{aligned} \tag{6.9}$$

by letting $q \rightarrow \infty$. This yields (6.8) as the left-hand side of (6.9) does not depend on ϵ .

Let $\mathcal{F}_s = \sigma\{B_{1t}, 0 \leq t \leq 1; B_{2t}, 0 \leq t < \infty, \eta(t), 0 \leq t \leq s\}$. Results (6.7) and (6.8) imply that, for any $0 \leq s < t \leq 1$,

$$\begin{aligned}
 & \mathbb{E} \left([\eta(t) - \eta(s)] \mid \mathcal{F}_s \right) = 0, \quad a.s., \\
 & \mathbb{E} \left(\{[\eta(t) - \eta(s)]^2 - [Y(t) - Y(s)]\} \mid \mathcal{F}_s \right) = 0, \quad a.s.
 \end{aligned}$$

Note that $\mathcal{F}_s \uparrow$, $\eta(s)$ is \mathcal{F}_s -measurable for each $0 \leq s \leq 1$ and $Y(t) = \tau_1^2 L_Z(t, 0)$ (for any fixed $t \in [0, 1]$) is \mathcal{F}_s -measurable for each $0 \leq s \leq 1$. It follows from Wang (2015, Lem. 3.4) that the finite-dimensional distributions of $(\eta(t), Y(t))$ coincide with those of $\{\mathbb{N} Y^{1/2}(t), Y(t)\}$, where \mathbb{N} is a normal variate independent of $Y(t)$. Since $\eta(t)$ does not depend on the choice of the subsequence $\{n''\}$, it follows from (6.5) and (6.6) that

$$\{S_n(t), Y_{nq}(t)\} \Rightarrow \{[\tau_1 L_Z(t, 0)]^{1/2} \mathbb{N}, \Lambda_q L_Z(t, 0)\}, \tag{6.10}$$

on $D_{R^2}[0, 1]$, where \mathbb{N} is a normal variate independent of $L_Z(t, 0)$. This, together with (6.4) and the continuous mapping theorem, yields (2.6).

The proof of (2.5) is similar. Set, for $0 \leq t \leq 1$ and $h > 0$,

$$S_{n,h}(t) = \left(\frac{d_n}{nh}\right)^{1/2} \sum_{k=1}^{\lfloor nt \rfloor} f(x_k/h, w_k), \quad Z_{n,h}(t) = \frac{d_n}{nh} \sum_{k=1}^{\lfloor nt \rfloor} f^2(x_k/h, w_k).$$

As $h \rightarrow 0$ and $nh/d_n \rightarrow \infty$, $Z_{n,h}(t) \Rightarrow Z(t) := \tau^2 L_Z(t, 0)$ by (7.4) in Proposition 7.4. The same arguments as those leading to (2.6) can be used to establish (2.5) except that $S_n(t), Y_{nq}(t)$, and $Y(t)$ are replaced by $S_{n,h}(t), Z_{n,h}(t)$, and $Z(t)$, respectively. The corresponding propositions with $h \rightarrow 0$ are given in Section 7. □

Proof of Corollary 2.1. We only prove (2.11). The proof of the other result is similar. Let $u_{1k} = \sum_{j=0}^{m_0} \psi_j \lambda'_{k-j}$, $u_{2k} = u_k - u_{1k} = \sum_{j=m_0+1}^{\infty} \psi_j \lambda'_{k-j}$ and, for $r = 0, 1, 2, \dots$,

$$\tilde{G}_{r,m_0} = \int_{-\infty}^{\infty} K(y) \mathbb{E} \{ u_{10} u_{1r} K(y + x_r) \} dy.$$

Using (2.9), for any $m_0 > 0$ and $q \geq 0$, we have

$$\begin{aligned} & \left(\frac{d_n}{n} \sum_{k=1}^n K^2(x_k) u_{1k}^2, \frac{d_n}{n} \sum_{k=1}^n K(x_k) u_{1k} K(x_{k+1}) u_{1,k+1}, \dots, \right. \\ & \quad \left. \frac{d_n}{n} \sum_{k=1}^n K(x_k) u_{1k} K(x_{k+q}) u_{1,k+q}, \left(\frac{d_n}{n} \right)^{1/2} \sum_{k=1}^n K(x_k) u_{1k} \right) \\ \Rightarrow & (\tilde{G}_{0,m_0} L_Z(1, 0), \tilde{G}_{1,m_0} L_Z(1, 0), \dots, \tilde{G}_{q,m_0} L_Z(1, 0), \tilde{\tau}_{1,m_0} \mathbb{N} L_Z^{1/2}(1, 0)), \end{aligned}$$

where $\tilde{\tau}_{1,m_0} = \tilde{G}_{0,m_0} + 2 \sum_{r=1}^{\infty} \tilde{G}_{r,m_0}$. This implies that, for any $m_0 > 0, q \geq 0$ and any continuous function with $l(0) = 1$,

$$\begin{aligned} & \left(\frac{d_n}{n} \sum_{k=1}^n K^2(x_k) u_{1k}^2, \widetilde{\mathcal{J}}_{n,q}, \left(\frac{d_n}{n} \right)^{1/2} \sum_{k=1}^n K(x_k) u_{1k} \right) \\ \rightarrow_D & (\tilde{G}_{0,m_0} L_Z(1, 0), \tilde{\tau}_{1,q}^2 L_Z(1, 0), \tilde{\tau}_{1,m_0} \mathbb{N} L_Z^{1/2}(1, 0)), \end{aligned}$$

where $\tilde{\tau}_{1,q}^2 = \tilde{G}_{0,m_0} + 2 \sum_{r=1}^q \tilde{G}_{r,m_0}$ and

$$\widetilde{\mathcal{J}}_{n,q} = \frac{d_n}{n} \sum_{k=1}^n K^2(x_k) u_{1k}^2 + \frac{2d_n}{n} \sum_{j=1}^q \ell \left(\frac{j}{M} \right) \sum_{k=1}^{n-j} K(x_k) K(x_{k+j}) u_{1k} u_{1,k+j}.$$

Consequently, to prove Corollary 2.1, it suffices to show the following:

(a) As $m_0 \rightarrow \infty$,

$$|\tilde{G}_0 - \tilde{G}_{0,m_0}| + \sum_{r=1}^{\infty} |\tilde{G}_r - \tilde{G}_{r,m_0}| \rightarrow 0. \tag{6.11}$$

(b) For any $m_0 \geq 1$,

$$\mathbb{E} \left| \sum_{k=1}^n u_{2k} K(x_k) \right|^2 \leq C(n/d_n) \left[\sum_{j=m_0}^{\infty} j^{1/4} (|\psi_{1j}| + |\psi_{2j}|) \right]^2. \tag{6.12}$$

(c) For any $r \geq 0$, as $n \rightarrow \infty$ first and then $m_0 \rightarrow \infty$,

$$\frac{d_n}{n} \sum_{k=1}^{n-r} K(x_k) K(x_{k+r}) (u_{1k} u_{1,k+r} - u_k u_{k+r}) = o_P(1). \tag{6.13}$$

Further, if $m_0 = m_0(n) \rightarrow \infty$, that is, m_0 depends on n , it also follows that there exists $M_1 \equiv M_{1n}$ depending on m_0 such that, as $n \rightarrow \infty$,

$$R_n := \frac{d_n}{n} \sum_{r=1}^{M_1} \left| \sum_{k=1}^{n-r} K(x_k)K(x_{k+r}) (u_{1k} u_{1,k+r} - u_k u_{k+r}) \right| = o_P(1). \tag{6.14}$$

(d) There exists $M \equiv M_n \rightarrow \infty$ so that, as $n \rightarrow \infty$ first and then $q \rightarrow \infty$,

$$\frac{d_n}{n} \sum_{r=q+1}^M \ell\left(\frac{r}{M}\right) \sum_{k=1}^{n-r} K(x_k)K(x_{k+r}) u_k u_{k+r} = o_P(1). \tag{6.15}$$

For the proofs of (6.11), (6.12)–(6.14), and (6.15), we refer to Propositions 7.5, 7.6, and 7.7, respectively. □

7. SUBSIDIARY PROPOSITIONS

This section proves the following propositions which are required in the proofs of Theorem 2.1 and Corollary 2.1. The notation is the same as in the previous section except where explicitly mentioned.

PROPOSITION 7.1. *For any fixed $0 \leq t \leq 1$, $r \geq 0$ and any bounded $h > 0$ satisfying $nh/d_n \rightarrow \infty$, $\psi_{nr}(t)$, $Z_{n,h}(t)$, and $S_{n,h}^2(t)$, $n \geq 1$, are uniformly integrable.*

PROPOSITION 7.2. *For any bounded $h > 0$ satisfying $nh/d_n \rightarrow \infty$, $\{Z_{n,h}(t)\}_{n \geq 1}$ and $\{S_{n,h}(t)\}_{n \geq 1}$ are tight on $D[0, 1]$.*

PROPOSITION 7.3. *For any $0 \leq s < t \leq 1$, we have that:*

(i) *if $h > 0$ is bounded satisfying $nh/d_n \rightarrow \infty$, then*

$$\lim_{n \rightarrow \infty} \mathbb{E} e^{i(Z_{n3}+Z_{n2})} [S_{n,h}(t) - S_{n,h}(s)] = 0; \tag{7.1}$$

(ii) *if $h \rightarrow 0$ satisfying $nh/d_n \rightarrow \infty$, then*

$$\lim_{n \rightarrow \infty} \mathbb{E} e^{i(Z_{n3}+Z_{n2})} \{ [S_{n,h}(t) - S_{n,h}(s)]^2 - [Z_{n,h}(t) - Z_{n,h}(s)] \} = 0; \tag{7.2}$$

(iii) *for any $\epsilon > 0$, there exists a $q_0 > 0$ such that*

$$\lim_{n \rightarrow \infty} \left| \mathbb{E} e^{i(Z_{n3}+Z_{n2})} \{ [S_n(t) - S_n(s)]^2 - [Y_{nq}(t) - Y_{nq}(s)] \} \right| \leq \epsilon, \tag{7.3}$$

for all $q \geq q_0$.

PROPOSITION 7.4. *For any bounded $h > 0$ satisfying $nh/d_n \rightarrow \infty$, we have*

$$Z_{n,h}(t) \Rightarrow \tau^2 L_Z(t, 0), \tag{7.4}$$

on $D_R[0, 1]$. If, in addition, $\gamma = 0$ and $\int \mathbb{E} \{ |\hat{f}(t, w_0)(1 + \|w_r\|^\beta) \} dt < \infty, 0 \leq r \leq m$, then

$$\{ \psi_{n0}(t), \psi_{n1}(t), \dots, \psi_{nm}(t) \} \Rightarrow \{ G_0, G_1, \dots, G_m \} L_Z(t, 0), \tag{7.5}$$

on $D_{Rm+1}[0, 1]$.

PROPOSITION 7.5. If $\gamma = 0$, we have $\sum_{r=1}^\infty |G_r| < \infty$ and $\sum_{r=1}^\infty |\tilde{G}_r| < \infty$, and (6.11) also holds.

PROPOSITION 7.6. Results (6.12)–(6.14) hold and, for any bounded $h > 0$ satisfying $nh/d_n \rightarrow \infty$, we have

$$\mathbb{E} \left| \sum_{k=1}^n u_{2k} K(x_k/h) \right|^2 \leq C(nh/d_n) \left[\sum_{j=m_0}^\infty j^{1/4} (|\psi_{1j}| + |\psi_{2j}|) \right]^2. \tag{7.6}$$

PROPOSITION 7.7. Result (6.15) holds.

7.1. Preliminary Lemmas

Except where explicitly mentioned, the proofs of all lemmas are given in the Appendix. Throughout this section, we let $\mathcal{F}_k = \sigma(\lambda_k, \lambda_{k-1}, \dots)$.

LEMMA 7.1. Let $p(s, s_1, \dots, s_m)$ be a real function of its components and $t_1, \dots, t_m \in \mathbb{Z}$, where $m \geq 0$. There exists an $A_0 > 0$ such that the following results hold.

(i) For any $h > 0$ and $k \geq 2m + A_0$, we have

$$\mathbb{E} |p(x_k/h, \lambda_{t_1}, \dots, \lambda_{t_m})| \leq \frac{Ch}{d_k} \int_{-\infty}^\infty \mathbb{E} |p(t, \lambda_1, \dots, \lambda_m)| dt. \tag{7.7}$$

(ii) For any $h > 0$, $k - j \geq 2m + A_0$ and $j + 1 \leq t_1, \dots, t_m \leq k$, we have

$$\mathbb{E} [|p(x_k/h, \lambda_{t_1}, \dots, \lambda_{t_m})| \mid \mathcal{F}_j] \leq \frac{Ch}{d_{k-j}} \int_{-\infty}^\infty \mathbb{E} |p(t, \lambda_1, \dots, \lambda_m)| dt. \tag{7.8}$$

(iii) For any $h > 0$ and $k - j \geq 1$, we have

$$\mathbb{E} [|p(x_k/h)| \mid \mathcal{F}_j] \leq \frac{Ch}{d_{k-j}} \int_{-\infty}^\infty |p(x)| dx. \tag{7.9}$$

Proof. For the proofs of (7.7) and (7.8), we refer to Lemma A.1 of Wang et al. (2021). As $\phi_0 \neq 0$, the proof of (7.9) is simple (see, for instance, Lemma 2.1(iii) of Wang, 2015). □

Recalling (6.3), $f(x, y) \leq T(x)(1 + \|y\|^\beta)$ and $\mathbb{E}\|w_1\|^{\max\{2, 4\beta\}} < \infty$, where $T(x)$ is bounded and integrable, a simple application of Lemma 7.1(i) and (ii) yields that, for any $h > 0$,

$$\sum_{k=1}^n \mathbb{E}f^2(x_k/h, w_k) \leq Cnh/d_n, \quad \mathbb{E}\left[\sum_{k=1}^n f^2(x_k/h, w_k)\right]^2 \leq C(nh/d_n)^2, \tag{7.10}$$

and (7.10) still holds if $f^2(x_k/h, w_k)$ is replaced by Y_{kj}^2 defined by

$$Y_{kj} = \mathbb{E}[f(x_k/h, w_k)|\mathcal{F}_{k-j}] - \mathbb{E}[f(x_k/h, w_k)|\mathcal{F}_{k-j-1}],$$

where $j \geq 0$ is a fixed integer. Furthermore, it follows from Lemma 7.1(iii) that, for any $r \geq 1$,

$$\mathbb{E}[|f(x_{k+r}/h, w_{k+r})||\mathcal{F}_k] \leq \{\mathbb{E}[T^2(x_{k+r}/h) | \mathcal{F}_k]\}^{1/2} \{\mathbb{E}[(1 + \|w_{k+r}\|^{2\beta})|\mathcal{F}_k]\}^{1/2} \leq Ch^{1/2} R_k,$$

where $R_k = \{\mathbb{E}[(1 + \|w_{k+r}\|^{2\beta})|\mathcal{F}_k]\}^{1/2}$ depending only on $\lambda_k, \dots, \lambda_{k-m_0}$. Hence, for any $r \geq 1, h > 0$, and $0 \leq s < t \leq 1$, we also have

$$\begin{aligned} & \sum_{k=[ns]+B_0}^{[nt]} \mathbb{E}\left[|f(x_k/h, w_k)| |f(x_{k+r}/h, w_{k+r})| \mid \mathcal{F}_{[ns]}\right] \\ & \leq \sum_{k=[ns]+B_0}^{[nt]} \mathbb{E}\left[|f(x_k/h, w_k)| \mathbb{E}\{|f(x_{k+r}/h, w_{k+r})| \mid \mathcal{F}_k\} \mid \mathcal{F}_{[ns]}\right] \\ & \leq Ch^{1/2} \sum_{k=[ns]+B_0}^{[nt]} \mathbb{E}\{|f(x_k/h, w_k)| R_k \mid \mathcal{F}_{[ns]}\} \\ & \leq Cnh^{3/2}(t-s)^\alpha/d_n, \end{aligned} \tag{7.11}$$

for some $\alpha > 0$, whenever B_0 is sufficiently large so that (7.8) is applicable. We remark that (7.11) holds for $r = 0$ if $h^{3/2}$ is replaced by h . These results will be used later.

In the next lemma, Ω_1 is set to be a subset of $\Omega = \{1, 2, \dots, k\}$, $\Omega_2 = \Omega - \Omega_1$ and

$$z_k(t) = \sum_{v=1}^k \epsilon_v(t\alpha_v + \beta_v).$$

LEMMA 7.2. *Suppose that $\sum_{v=1}^k \alpha_v^2 \leq C\tau_k^2$ and, for any Ω_1 satisfying $\#\Omega_1 \leq \sqrt{k}$,*

$$B_{1k} := \sum_{v \in \Omega_2} \alpha_v^2 \geq \tau_k^2, \tag{7.12}$$

for some sequence of constants τ_k . Then, for any $\delta \geq 0$ and $s_1, s_2 \in \mathbb{R}^+$, we have

$$\int \min\{1, s_1 |t|^\delta + s_2\} |\mathbb{E} e^{iz_k(t)}| dt \leq C(k^{-3} + s_1 \tau_k^{-1-\delta} [1 + (\sum_{v=1}^k \beta_v^2)^{\delta/2}] + s_2 \tau_k^{-1}); \tag{7.13}$$

$$\int \min\{1, s_1 |t|\} \min\{1, |t|\} |\mathbb{E} e^{iz_k(t)}| dt \leq C(k^{-3} + s_1 \tau_k^{-3} [1 + \sum_{v=1}^k \beta_v^2]). \tag{7.14}$$

If, in addition, $\sum_{v=1}^k \beta_v^2 \leq a < \infty$, then

$$\int_{|t| \geq B/\tau_k} |\mathbb{E} e^{iz_k(t)}| dt \leq C(k^{-3} + \tau_k^{-1} B^{-1}), \tag{7.15}$$

for any $B \geq 2a^{1/2}$.

Proof. The proof of Lemma 7.2 is similar to that of Wang and Phillips (2011, pp. 246–247) and is therefore omitted. But an outline of the proof is given in Appendix A.1 for completeness. \square

Since Lemma 7.2 still holds when $z_k(t)$ is replaced by $z_{k-m_0}(t)$ when $k \geq m_0^2$ and since w_k depends only on $\lambda_k, \dots, \lambda_{k-m_0}$, the following lemma is a direct consequence of Lemma 7.2.

LEMMA 7.3. Let $g(x, y)$ be a real function satisfying

- $|\mathbb{E} g(t, w_1)| \leq C \min\{1, |t|\}$ and $\sup_t \mathbb{E} \{(1 + |\epsilon_0|) |g(t, w_1)|\} < \infty$.

For any bounded $h > 0$ and $\tau_k \leq Ck^2$, we have

$$\int_{-\infty}^{\infty} |\mathbb{E} e^{iz_k(t/h)} g(t, w_k)| dt \leq Ch \tau_k^{-1}, \tag{7.16}$$

for all $k \geq m_0^2$. Instead of (7.16), we also have

$$\int_{-\infty}^{\infty} |\mathbb{E} e^{iz_k(t/h)} g(t, w_k)| dt \leq Ch \{(1 + \alpha_{k0}) \tau_k^{-2} [1 + (\sum_{v=1}^k \beta_v^2)^{1/2}] + \beta_{k0} \tau_k^{-1}\}, \tag{7.17}$$

where $\alpha_{k0} = \max_{0 \leq i \leq m_0 \vee (k-1)} |\alpha_{k-i}|$ and $\beta_{k0} = \max_{0 \leq i \leq m_0 \vee (k-1)} |\beta_{k-i}|$. Similarly, when $\sup_k \alpha_{k0} = O(1)$, we have

$$\int_{-\infty}^{\infty} \min\{1, |t|/h\} |\mathbb{E} e^{iz_k(t/h)} g(t, w_k)| dt \leq Ch \{k^{-3} + [\beta_{k0}(\tau_k^{-2} + k^{-3}) + \tau_k^{-3}](1 + \sum_{\nu=1}^k \beta_{\nu}^2)\}. \tag{7.18}$$

Proof. See Appendix A.2. □

Let $I_k(m) = \int \mathbb{E} (e^{isx_k/h+i \sum_{j=m+1}^l \gamma_j \epsilon_j} g(s, w_k) | \mathcal{F}_m) ds$ and

$$I_{k,l}(m) = \int \int \mathbb{E} (e^{isx_k/h+itx_l/h+i \sum_{j=m+1}^l \gamma_j \epsilon_j} g(s, w_k)g(t, w_l) | \mathcal{F}_m) ds dt,$$

where $g(x,y)$ is a real function given in Lemma 7.3, and let

$$\Pi_{k,l}(B) = \int_{|s| \geq B/d_k} \int_{|t| \geq B/d_l} g_1(t)g_2(t) \mathbb{E} (e^{isx_k/h+itx_l/h+i \sum_{j=1}^l \gamma_j \epsilon_j} | \mathcal{F}_0) ds dt,$$

where $g_1(t)$ and $g_2(t)$ are bounded real functions. The next lemma is an application of Lemma 7.3.

LEMMA 7.4. *Let $m \geq 0$, $l - k \geq A_0^2 + 1$ and $k - m \geq A_0^2 + 1$, where $A_0 \geq m_0$ and m_0 is given as in Lemma 7.3. Suppose $a := \sum_{j=1}^l \gamma_j^2 < \infty$.*

(i) *For any $h > 0$, we have*

$$|I_k(m)| \leq Ch [d_{k-m}^{-2}(1 + a^{1/2}) + \beta_{l0} d_{k-m}^{-1}], \tag{7.19}$$

$$|I_{k,l}(m)| \leq Ch^2 d_{k-m}^{-1} [d_{l-k}^{-2}(1 + a^{1/2}) + \beta_{l0} d_{l-k}^{-1}], \tag{7.20}$$

where $\beta_{l0} = \max_{0 \leq j \leq m_0} |\gamma_{l-j}|$.

(ii) *Under SM, if $|\gamma_j| \leq C/\sqrt{n}$ where $m \leq j \leq l$, for any $h > 0$, we have*

$$|I_k(m)| \leq Ch ((k - m)^{-1} + \sqrt{k - m}/\sqrt{n}), \tag{7.21}$$

$$|I_{k,l}(m)| \leq Ch^2 [(l - k)^{-1}(k - m)^{-1} + (l - k)^{-3/2}(k - m)^{-1/2}]. \tag{7.22}$$

(iii) *For any $h > 0$ and $B \geq 2a^{1/2}$, we have*

$$|\Pi_{k,l}(B)| \leq Ch^2 [(l - k)^{-2} + B^{-1} d_{l-k}^{-1}] d_k^{-1}. \tag{7.23}$$

Proof. See Appendix A.3. □

Let $I_k(h) = f(x_k/h, w_k) \exp \{i \sum_{j=m+1}^n \mu_j \epsilon_j / \sqrt{n}\}$ and

$$I_{lk}(h) = f(x_k/h, w_k) f(x_l/h, w_l) \exp \{i \sum_{j=m+1}^n \mu_j \epsilon_j / \sqrt{n}\},$$

where μ_l are constants satisfying $|\mu_l| \leq C$. Using Lemma 7.4, we have the following results.

LEMMA 7.5. *There exists a $B_0 \geq m_0$ such that, for all $m \geq 0, l - k \geq B_0, k - m \geq B_0$, and bounded $h > 0$,*

(i) *under LM,*

$$|\mathbb{E}[I_k(h) | \mathcal{F}_m]| \leq Ch(d_{k-m}^{-2} + d_{k-m}/\sqrt{n}), \tag{7.24}$$

$$|\mathbb{E}[II_k(h) | \mathcal{F}_m]| \leq Ch^2 d_{k-m}^{-1} (d_{l-k}^{-2} + d_{l-k}/\sqrt{n}), \tag{7.25}$$

(ii) *under SM,*

$$|\mathbb{E}[I_k(h) | \mathcal{F}_m]| \leq Ch((k - m)^{-1} + \sqrt{k - m}/\sqrt{n}), \tag{7.26}$$

$$|\mathbb{E}[II_k(h) | \mathcal{F}_m]| \leq Ch^2 [(l - k)^{-1} (k - m)^{-1} + (l - k)^{-3/2} (k - m)^{-1/2}]. \tag{7.27}$$

LEMMA 7.6. *There exists a $B_0 \geq m_0$ such that, for all $m \geq 0, l - k \geq B_0, k - m \geq B_0$ and bounded $h > 0$,*

(i) *under LM,*

$$|\mathbb{E}\{f(x_l/h, w_l)\mathbb{E}[f(x_k/h, w_k) | \mathcal{F}_{k-m}]\}| \leq Ch^2 d_k^{-1} d_{l-k}^{-2}, \tag{7.28}$$

(ii) *under SM,*

$$\begin{aligned} &|\mathbb{E}\{f(x_l/h, w_l)\mathbb{E}[f(x_k/h, w_k) | \mathcal{F}_{k-m}]\}| \\ &\leq Ch^2 [(l - k)^{-1} k^{-1} + (l - k)^{-3/2} k^{-1/2}]. \end{aligned} \tag{7.29}$$

The proofs of Lemmas 7.5 and 7.6 are given in Appendices A.4 and A.5.

LEMMA 7.7. *Let $\Gamma(\cdot)$ be a measurable function with $\Gamma(\lambda_1) = 0$ and $\mathbb{E}\Gamma^2(\lambda_1) < \infty$. There exists an A_0 such that:*

(a) *for all $k \geq A_0$ and $|l - k| \leq A_0$,*

$$|\mathbb{E}\{\Gamma(\lambda_{k-j})\Gamma(\lambda_{l-j})K(x_k/h)K(x_l/h)\}| \leq Chd_k^{-1}; \tag{7.30}$$

(b) *for all $k \geq A_0, l - k \geq A_0$, and $l - j \leq k$,*

$$|\mathbb{E}\{\Gamma(\lambda_{k-j})\Gamma(\lambda_{l-j})K(x_k/h)K(x_l/h)\}| \leq Ch^2 d_k^{-1} d_{l-k}^{-1}; \tag{7.31}$$

(c) *for all $k \geq A_0, l - k \geq A_0$, and $l - j > k$,*

$$\begin{aligned} &|\mathbb{E}\{\Gamma(\lambda_{k-j})\Gamma(\lambda_{l-j})K(x_k/h)K(x_l/h)\}| \\ &\leq Ch^2 \begin{cases} \sum_{k=0}^j |\phi_k| d_k^{-1} d_{l-k}^{-2}, & \text{under LM,} \\ k^{-1} (l - k)^{-1} + k^{-1/2} (l - k)^{-3/2}, & \text{under SM.} \end{cases} \end{aligned} \tag{7.32}$$

Similarly, uniformly for $y \in R$, we have

$$\begin{aligned} & \left| \mathbb{E} \left\{ K(y + x_l/h) \Gamma(\lambda_{l-j}) \Gamma(\lambda_{-k}) \right\} \right| \\ & \leq Ch \begin{cases} d_l^{-1}, & \text{if } |l-j+k| \leq A_0, \\ \sum_{s=0}^j |\phi_s| \sum_{s=k}^{l+k} |\phi_s| (d_l^{-3} + l^{-3}), & \text{if } |l-j+k| > A_0, \end{cases} \end{aligned} \tag{7.33}$$

for any $A_0 \geq 1$ and $j, k \geq 0$.

Proof. See Appendix A.6. □

Our final lemma gives a useful tightness criterion for a class of stochastic processes on $D[0, 1]$.

LEMMA 7.8. *Let X_{nk} be a sequence of random variables and $X_n(t) = \sum_{k=1}^{[nt]} X_{nk}$. The sequence $\{X_n(t)\}$ is tight in $D[0, 1]$ if $\max_{1 \leq k \leq n} |X_{nk}| = o_P(1)$ and there exist an integer $A_0 \geq 0$ and a number $\alpha_n(\epsilon, \delta)$ such that*

$$P\left(\left| \sum_{k=[nt_m]+A_0}^{[ns]} X_{nk} \right| \geq \epsilon \mid X_n(t_1), \dots, X_n(t_m) \right) \leq \alpha_n(\epsilon, \delta),$$

and

$$\lim_{\delta \rightarrow 0} \limsup_{n \rightarrow \infty} \alpha_n(\epsilon, \delta) = 0,$$

for each positive $\epsilon > 0$, where $0 \leq t_1 \leq t_2 \leq \dots \leq t_m \leq s \leq 1$ and $s - t_m \leq \delta$.

Proof. If $A_0 = 0$, Lemma 7.8 is a special case of Billingsley (1974, Thm. 4). Extension to integer $A_0 \geq 1$ is trivial under the condition that $\max_{1 \leq k \leq n} |X_{nk}| = o_P(1)$. The details are omitted. □

7.2. Proofs of Propositions

Propositions 7.4 and 7.7 are treated separately due to their complexity and their proofs are given later in Sections 7.3 and 7.4, respectively.

Proof of Proposition 7.1. We only prove uniformity of $S_{n,h}^2(1)$ for bounded $h > 0$ satisfying $nh/d_n \rightarrow \infty$. The other results are similar and simpler. Let $m \geq m_0$ be a constant that will be specified later. Let

$$\begin{aligned} S_{1n} &= \left(\frac{d_n}{nh} \right)^{1/2} \sum_{k=1}^n \mathbb{E} [f(x_k/h, w_k) \mid \mathcal{F}_{k-m}], \\ S_{2n} &= \left(\frac{d_n}{nh} \right)^{1/2} \sum_{k=1}^n \{f(x_k/h, w_k) - \mathbb{E} [f(x_k/h, w_k) \mid \mathcal{F}_{k-m}]\}. \end{aligned}$$

Note that, for any $A \geq 2$,

$$\begin{aligned} \mathbb{E}S_{n,h}^2(1)I(S_{n,h}^2(1) \geq A) &\leq 2\mathbb{E}S_{1n}^2 + 2\mathbb{E}S_{2n}^2I(S_{1n}^2 + S_{2n}^2 \geq A/2) \\ &\leq 2\mathbb{E}S_{1n}^2 + 8A^{-1}\mathbb{E}S_{2n}^4 + 2\mathbb{E}S_{2n}^2I(S_{1n}^2 \geq A/4) \\ &\leq 4\mathbb{E}S_{1n}^2 + 16A^{-1}\mathbb{E}S_{2n}^4. \end{aligned}$$

It suffices to show that, for some $c_0 > 0$,

- (a) $\mathbb{E}S_{2n}^4 \leq c_0m^4$;
- (b) under LM , $\mathbb{E}S_{1n}^2 \leq c_0d_m^{1/2-\mu}$;
- (c) under SM , $\mathbb{E}S_{1n}^2 \leq c_0(d_m^{-1/2} + \log^2 n/\sqrt{n})$.

Indeed, for any $\epsilon > 0$, by taking A, n sufficiently large and $m = A^{1/8}$, it follows from (a) to (c) that

$$\mathbb{E}S_{n,h}^2(1)I(S_{n,h}^2(1) \geq A) \leq 4c_0(d_m^{-1/2} + d_m^{1/2-\mu}) + 16c_1A^{-1/2} + c_0\log^2 n/\sqrt{n} \leq \epsilon,$$

under both LM and SM , due to $d_m \rightarrow 0$ and $\mu > 1/2$.

To prove (a), let $Y_{kj} = \mathbb{E}[f(x_k/h, w_k)|\mathcal{F}_{k-j}] - \mathbb{E}[f(x_k/h, w_k)|\mathcal{F}_{k-j-1}]$, $0 \leq j \leq m-1$. We may write

$$S_{2n} = \left(\frac{d_n}{nh}\right)^{1/2} \sum_{j=0}^{m-1} \sum_{k=1}^n Y_{kj}.$$

Note that Y_{kj} forms a martingale difference. Hölder’s and Burkholder’s inequalities imply that

$$\mathbb{E}S_{2n}^4 \leq m^3 \left(\frac{d_n}{nh}\right)^2 \sum_{j=0}^{m-1} \mathbb{E}\left(\sum_{k=1}^n Y_{kj}\right)^4 \leq C_2m^3 \left(\frac{d_n}{nh}\right)^2 \sum_{j=0}^{m-1} \mathbb{E}\left(\sum_{k=1}^n Y_{kj}^2\right)^2 \leq c_0m^4,$$

for some $c_0 > 0$, which yields (a), where we have used the result (7.10) with $f^2(\cdot)$ replaced by Y_{kj}^2 .

We next prove (b) and (c). Let $g_k = \mathbb{E}[f(x_k/h, w_k)|\mathcal{F}_{k-m}]$. For some $q \geq 1$, we may write

$$\begin{aligned} \mathbb{E}S_{1n}^2 &= \frac{d_n}{nh} \left[\sum_{k=1}^n \mathbb{E}g_k^2 + 2 \sum_{k=1}^n \sum_{j=k+1}^{k+q} \mathbb{E}g_k g_j + 2 \sum_{k=1}^n \sum_{j=k+q}^n \mathbb{E}(g_k g_j) \right] \\ &= R_{n1} + R_{n2} + R_{n3}. \end{aligned} \tag{7.34}$$

Recall (6.3). It follows from (7.8) in Lemma 7.1 that $|g_k| \leq Ch/d_m$. On the other hand, $\mathbb{E}|g_k| \leq \mathbb{E}|f(x_k/h, w_k)| \leq Ch/d_k$. As a consequence, we have

$$|R_{n1}| + |R_{n2}| \leq Cqh/d_m \frac{d_n}{nh} \sum_{k=l_n}^n \mathbb{E}|q_k| \leq Cqhd_m^{-1}.$$

As for R_{n3} , by taking $m \geq B_0$ where B_0 is given in Lemma 7.6,

(i) under LM , it follows from (7.28) that, for any $q \geq B_0$,

$$\begin{aligned} |R_{n3}| &\leq \frac{2d_n}{nh} \sum_{k=1}^n \sum_{j=k+q}^n |\mathbb{E}(g_k g_j)| \leq C \frac{hd_n}{n} \sum_{k=1}^n \sum_{j=k+q}^n d_k^{-1} d_{j-k}^{-2} \\ &\leq Ch \int_q^\infty x^{2\mu-3} \rho^{-2}(x) dx, \end{aligned}$$

(ii) under SM , it follows from (7.29) that, for any $q \geq B_0$,

$$\begin{aligned} |R_{n3}| &\leq \frac{2}{\sqrt{nh}} \sum_{k=1}^n \sum_{j=k+q}^n |\mathbb{E}(g_k g_j)| \\ &\leq \frac{Ch}{\sqrt{n}} \sum_{k=1}^n \sum_{j=k+q}^n [(j-k)^{-1} k^{-1} + (j-k)^{-3/2} k^{-1/2}] \\ &\leq Ch(\log^2 n / \sqrt{n} + \int_q^\infty x^{-3/2} dx). \end{aligned}$$

Taking these estimates into (7.34), we obtain (b) and (c) by letting $q = \sqrt{d_n}$, as h is bounded. This completes the proof. \square

Proof of Proposition 7.2. We prove tightness of $S_{n,h}(t)$. Tightness of $Z_{n,h}(t)$ is shown in a similar way to Wang (2015, Thm. 2.20) and the details are omitted.

Recalling (6.1) and Lemma 7.8, it suffices to prove the following: for any fixed $s \in [0, 1]$, for each $\epsilon > 0$ and any bounded $h > 0$ satisfying $nh/d_n \rightarrow \infty$, there exists a sequence of $\alpha_n(\epsilon, \delta)$ satisfying $\lim_{\delta \rightarrow 0} \limsup_{n \rightarrow \infty} \alpha_n(\epsilon, \delta) = 0$ such that

$$I_n := \sup_{|t-s| \leq \delta} \mathbb{P}\left(\left| \sum_{k=\lfloor ns \rfloor + B_0}^{\lfloor nt \rfloor} f(x_k/h, w_k) \right| \geq \epsilon (nh/d_n)^{1/2} \mid \mathcal{F}_{\lfloor ns \rfloor} \right) \leq \alpha_n(\epsilon, \delta), \tag{7.35}$$

where B_0 is chosen as in Lemma 7.5. In fact, by noting

$$\begin{aligned} J_n(s, t) &:= \mathbb{E}\left[\left| \sum_{k=\lfloor ns \rfloor + B_0}^{\lfloor nt \rfloor} f(x_k/h, w_k) \right|^2 \mid \mathcal{F}_{\lfloor ns \rfloor} \right] \\ &\leq 2 \sum_{k=\lfloor ns \rfloor + B_0}^{\lfloor nt \rfloor} \sum_{l=k+1}^{\lfloor nt \rfloor} \mathbb{E}(|f(x_k/h, w_k)| |f(x_l/h, w_l)| \mid \mathcal{F}_{\lfloor ns \rfloor}) \\ &\quad + 2 \sum_{k=\lfloor ns \rfloor + B_0}^{\lfloor nt \rfloor} \sum_{l=k+2B_0}^n \left| \mathbb{E}\{f(x_k/h, w_k) f(x_l/h, w_l) \mid \mathcal{F}_{\lfloor ns \rfloor}\} \right|, \end{aligned}$$

it follows from (7.11) and Lemma 7.5 that, for some $\alpha > 0$:

(a) under *LM* [using (7.25)],

$$\begin{aligned}
 J_n(s, t) &\leq Cnh(t-s)^\alpha/d_n + Ch^2 \sum_{k=[ns]+1}^{[nt]} \sum_{l=k+1}^n d_{k-[ns]}^{-1} d_{l-k}^{-2} \\
 &\leq 2Cnh(t-s)^\alpha/d_n;
 \end{aligned}$$

(b) under *SM* [using (7.27)],

$$\begin{aligned}
 J_n(s, t) &\leq C\sqrt{nh}(t-s)^\alpha + \\
 &Ch^2 \sum_{k=[ns]+1}^{[nt]} \sum_{l=k+1}^n [(l-k)^{-1}(k-[ns])^{-1} + (l-k)^{-3/2}(k-[ns])^{-1/2}] \\
 &\leq 2C\sqrt{nh}(t-s)^\alpha.
 \end{aligned}$$

Now (7.35) follows by choosing $\alpha_n(\epsilon, \delta) = 2C\epsilon^{-2}\delta^\alpha$ and the fact that

$$I_n \leq \epsilon^{-2}d_n/(nh) \sup_{|t-s|\leq\delta} J_n(s, t) \leq \alpha_n(\epsilon, \delta).$$

□

Proof of Proposition 7.3. We start with (7.2). Due to the iid properties of λ_k , there exist constants μ_j with $|\mu_j| \leq C$,

$$\begin{aligned}
 &\left| \mathbb{E} e^{i(Z_{n3}+Z_{n2})} \{ [S_{n,h}(t) - S_{n,h}(s)]^2 - [Z_{n,h}(t) - Z_{n,h}(s)] \} \right| \\
 &\leq \mathbb{E} \left| \mathbb{E} \left[e^{i\sum_{j=[ns]+1}^{[nt]} \mu_j \epsilon_j} \{ [S_{n,h}(t) - S_{n,h}(s)]^2 - [Z_{n,h}(t) - Z_{n,h}(s)] \} \mid \mathcal{F}_{[ns]} \right] \right| \\
 &\leq \frac{d_n}{nh} \sum_{k=[ns]+1}^n \sum_{l=k+1}^n E |\mathbb{E} [II_{lk}(h) \mid \mathcal{F}_{[ns]}]| \\
 &\leq \frac{d_n}{nh} \sum_{k=[ns]+1}^n \left(\sum_{l=k+1}^{k+B_0} + \sum_{l=k+B_0}^n \right) E |\mathbb{E} [II_{lk}(h) \mid \mathcal{F}_{[ns]}]| \\
 &=: R_{n4} + R_{n5},
 \end{aligned} \tag{7.36}$$

where B_0 and $II_{lk}(h)$ are defined as in Lemma 7.5. Similar to (7.11) with minor modifications, under both *LM* and *SM*, we have $R_{n4} \leq Ch^{1/2}$. To estimate R_{n5} , under *LM*, it follows from (7.25) that

$$R_{n5} \leq \frac{Cd_n}{nh} h^2 \sum_{k=1}^n \sum_{l=k+B_0}^n d_k^{-1} (d_{l-k}^{-2} + d_{l-k}/\sqrt{n}) \leq Ch.$$

Similarly, under *SM*, we have $R_{n5} \leq Ch$ by (7.27). Taking these estimates into (7.36), we have (7.2) as $h \rightarrow 0$.

In a similar way for any $q \geq B_0$, we have

$$\begin{aligned} & \left| \mathbb{E} e^{i(Z_{n3}+Z_{n2})} \{ [S_n(t) - S_n(s)]^2 - [Y_{nq}(t) - Y_{nq}(s)] \} \right| \\ & \leq \frac{d_n}{n} \sum_{k=[ns]+1}^n \sum_{l=k+q}^n E | \mathbb{E} [I_{lk}(1) \mid \mathcal{F}_{[ns]}] | \\ & \leq \begin{cases} \frac{d_n}{\sqrt{n}} \sum_{k=[ns]+1}^n \sum_{l=k+q}^n \left[(l-k)^{-1} (k-[ns])^{-1} + (l-k)^{-3/2} (k-[ns])^{-1/2} \right], & \text{under LM,} \\ \frac{d_n}{n} \sum_{k=[ns]+1}^n \sum_{l=k+q}^n d_{k-[ns]}^{-1} d_{l-k}^{-2}, & \text{under SM,} \end{cases} \\ & \leq C \begin{cases} \int_q^\infty x^{2\mu-3} dx, & \text{under LM,} \\ \int_q^\infty x^{-3/2} dx + \log^2 n / \sqrt{n}, & \text{under SM,} \end{cases} \\ & \leq \epsilon + C \log^2 n / \sqrt{n}, \end{aligned}$$

by choosing q sufficiently large. This proves (7.3). The proof of (7.1) is similar and simpler, so the details are omitted. □

Proof of Proposition 7.5. With $\gamma = 0$ in assumption A1(i), we may write

$$x_r = \sum_{i=1}^r \sum_{j=0}^\infty \phi_j \epsilon_{i-j} = \sum_{j=1}^r a_{r-j} \epsilon_j + \sum_{j=0}^\infty [a_{r+j} - a_j] \epsilon_{-j}, \tag{7.37}$$

where $a_l = \sum_{s=0}^l \phi_s$ and $a_l = 0$ if $l < 0$. Let $z_r = \sum_{k=1}^r \epsilon_k a_{r-k}$ and $z_{1r} = \sum_{j=0}^{m_0} [a_{r+j} - a_j] \epsilon_{-j}$. We have $var(z_r) \sim d_r^2$ for $r \geq 2m_0$ and, when m_0 is fixed,

$$\begin{aligned} | \mathbb{E} \hat{f}(s, w_0) e^{-isz_{1r}} | & \leq \mathbb{E} | \hat{f}(s, w_0) (e^{-isz_{1r}} - 1) | + | \mathbb{E} \hat{f}(s, w_0) | \\ & \leq C(1 + |a_r|) \min\{1, |s|\}. \end{aligned}$$

Now it is readily seen from the i.i.d. properties of ϵ_k and (7.18) in Lemma 7.3 that

$$\begin{aligned} |G_r| & \leq \frac{1}{2\pi} \int_{-\infty}^\infty | \mathbb{E} \{ \hat{f}(s, w_0) e^{-isz_{1r}} \} | | \mathbb{E} \{ \hat{f}(s, w_r) e^{-isz_r} \} | ds \\ & \leq C(1 + |a_r|) \int_{-\infty}^\infty \min\{1, |s|\} | \mathbb{E} \{ \hat{f}(s, w_r) e^{-isz_r} \} | ds \\ & \leq C(1 + |a_r|) (d_r^{-3} + r^{-3}). \end{aligned}$$

Hence, $\sum_{r=2m_0}^\infty |G_r| < \infty$ due to $|a_r| \leq C$ under SM and $|a_r| \leq d_r$ under LM.

To prove (6.11) and $\sum_{r=1}^\infty | \tilde{G}_r | < \infty$, we make use of (7.33) in Lemma 7.7. In fact, for any $r \geq 1$ and $y \in R$, it follows from (7.33) that

$$\begin{aligned} & | \mathbb{E} \{ (u_{10} u_{1r} - u_0 u_r) K(y + x_r) \} | \\ & \leq \left(\sum_{k=m_0+1}^\infty \sum_{j=0}^\infty + \sum_{k=0}^\infty \sum_{j=m_0+1}^\infty \right) | \mathbb{E} \{ \psi_k \lambda'_{-k} \psi_j \lambda'_{r-j} K(y + x_r) \} | \end{aligned}$$

$$\begin{aligned}
 &\leq 2 \sum_{k=m_0+1}^{\infty} \sum_{j=r+k-1}^{r+k+1} d_r^{-1} \|\psi_k\| \|\psi_j\| \\
 &\quad + 2 \sum_{k=m_0+1}^{\infty} \sum_{j=0}^{\infty} \|\psi_k\| \|\psi_j\| \sum_{s=0}^j |\phi_s| \sum_{s=k}^{r+k} |\phi_s| (d_r^{-3} + r^{-3}) \\
 &\leq 2d_r^{-1} \sum_{k=m_0+1}^{\infty} \|\psi_k\| \sum_{j=-1}^1 \|\psi_{j+r+k}\| \\
 &\quad + 2C \sum_{k=m_0+1}^{\infty} \sum_{j=0}^{\infty} \|\psi_k\| \|\psi_j\| \sum_{s=0}^j |\phi_s| \sum_{s=k}^{r+k} |\phi_s| (d_r^{-3} + r^{-3}).
 \end{aligned}$$

Note that $\sum_{s=0}^j |\phi_s| \sum_{s=k}^{r+k} |\phi_s| (d_r^{-3} + r^{-3}) \leq Cj^{1/2} k^{1/2} r^{-3/2}$ under both *SM* and *LM*. It is readily seen from $\sum_{k=0}^{\infty} k^{1/2} \|\psi_k\| < \infty$ that

$$\begin{aligned}
 \sum_{r=1}^{\infty} |\tilde{G}_r - \tilde{G}_{r,m_0}| &\leq \int_{-\infty}^{\infty} K(y) \sum_{r=1}^{\infty} |\mathbb{E}\{(u_{10}u_{1r} - u_0u_r)K(y+x_r)\}| dy \\
 &\leq C \sum_{k=m_0+1}^{\infty} k^{1/2} \|\psi_k\| \int K(y) dy \rightarrow 0,
 \end{aligned} \tag{7.38}$$

as $m_0 \rightarrow \infty$. Similarly, we have $|\tilde{G}_0 - \tilde{G}_{0,m_0}| \rightarrow 0$, as $m_0 \rightarrow \infty$, and $\sum_{r=1}^{\infty} |\tilde{G}_r| < \infty$. The proof of Proposition 7.5 is then complete. □

Proof of Proposition 7.6. The proofs of (6.13) and (6.14) are simply established using Lemma 7.1. Indeed, by noting that

$$\begin{aligned}
 &\left| \sum_{k=1}^{n-r} K(x_k)K(x_{k+r}) (u_{1k}u_{1,k+r} - u_ku_{k+r}) \right| \\
 &\leq C \left(\sum_{l=m_0+1}^{\infty} \sum_{l_1=0}^{\infty} + \sum_{l=0}^{\infty} \sum_{l_1=m_0+1}^{\infty} \right) \sum_{k=1}^{n-r} K(x_k) |\psi_{l_1} \lambda'_{k-l} \psi_{l_1} \lambda'_{k+r-l_1}|,
 \end{aligned}$$

it follows from Lemma 7.1(i) and $\sum_{l=0}^{\infty} l \|\psi_l\| < \infty$ that, for some constant $A_0 > 0$,

$$\begin{aligned}
 \mathbb{E}|R_n| &\leq CM_1 \sum_{l=m_0+1}^{\infty} \sum_{l_1=0}^{\infty} \|\psi_{l_1}\| \|\psi_l\| \frac{d_n}{n} \left[(A_0 + 2) + \sum_{k=1}^n d_k^{-1} \right] \\
 &\leq C_1 M_1 \sum_{l=m_0+1}^{\infty} \|\psi_l\| \leq CM_1 m_0^{-1}.
 \end{aligned}$$

Hence, (6.14) follows if we take $M_1 = \sqrt{m_0}$. The proof of (6.13) is similar.

We next prove (7.6). Let $\sum_{j=k}^l = 0$ for $k > l$ and $\Gamma(\cdot)$ be a measurable function with $\Gamma(\lambda_1) = 0$ and $\mathbb{E}\Gamma^2(\lambda_1) < \infty$. Since $K(x)$ is bounded, for A_0 being chosen as in Lemma 7.7, we have

$$\begin{aligned} \Delta_n &\equiv \left| \sum_{k=1}^n \Gamma(\lambda_{k-j}) K(x_k/h) \right|^2 \\ &\leq 2 \left| \sum_{k=A_0}^n \Gamma(\lambda_{k-j}) K(x_k/h) \right|^2 + C \left(\sum_{k=1}^{A_0} |\Gamma(\lambda_{k-j})| \right)^2 \\ &= 2 \left(\sum_{k=A_0}^n \sum_{|k-l| < A_0} + 2 \sum_{k=A_0}^{n-1} \sum_{l=k+A_0}^n \right) \Gamma(\lambda_{k-j}) \Gamma(\lambda_{l-j}) K(x_k/h) K(x_l/h) \\ &\quad + C \left(\sum_{k=1}^{A_0} |\Gamma(\lambda_{k-j})| \right)^2 \\ &=: \Delta_{1n} + \Delta_{2n} + \Delta_{3n}, \quad \text{say.} \end{aligned} \tag{7.39}$$

It follows from Lemma 7.7 that

$$\begin{aligned} \mathbb{E}|\Delta_{1n}| &\leq Ch \sum_{k=1}^n \sum_{|k-l| < A_0} 1/d_k \leq C_1 nh/d_n, \\ \mathbb{E}|\Delta_{2n}| &\leq Ch^2 \begin{cases} \sum_{k=A_0}^{n-1} d_k^{-1} \left(\sum_{l=k+A_0}^{n \wedge (k+j)} d_{l-k}^{-1} + \sum_{k=0}^j |\phi_k| \sum_{l=k+j}^n d_{l-k}^{-2} \right) & \text{under LM} \\ \sum_{k=A_0}^{n-1} k^{-1/2} \sum_{l=k+A_0}^{n \wedge (k+j)} (l-k)^{-1/2} + \\ \sum_{k=A_0}^{n-1} \sum_{l=k+j}^n [k^{-1}(l-k)^{-1} + k^{-1/2}(l-k)^{-3/2}] & \text{under SM} \end{cases} \\ &\leq C(nh^2/d_n) \begin{cases} j/d_j + \sum_{k=0}^j |\phi_k| & \text{under LM,} \\ j^{1/2} + \log^2 n/\sqrt{n} + 1 & \text{under SM,} \end{cases} \\ &\leq Cj^{1/2} nh^2/d_n, \end{aligned}$$

where we have used the fact $\sum_{k=0}^j |\phi_k| \leq Cj/d_j \leq Cj^{1/2}$ under LM. On the other hand, it is readily seen that $\mathbb{E}|\Delta_{3n}| \leq CA_0^2$. Taking these estimates into (7.39), for any bounded h , we have

$$\mathbb{E} \left| \sum_{k=1}^n \Gamma(\lambda_{k-j}) K(x_k/h) \right|^2 \leq Cj^{1/2} nh/d_n. \tag{7.40}$$

The result (6.12) now follows from

$$\begin{aligned} \mathbb{E} \left| \sum_{k=1}^n u_{k,m_0} K(x_k/h) \right|^2 &= \mathbb{E} \left| \sum_{j=m_0}^{\infty} \sum_{k=1}^n \psi_j \lambda'_{k-j} K(x_k/h) \right|^2 \\ &\leq \sum_{j=m_0}^{\infty} j^{1/4} (|\psi_{1j}| + |\psi_{2j}|) \sum_{j=m_0}^{\infty} j^{-1/4} (|\psi_{1j}| + |\psi_{2j}|)^{-1} \mathbb{E} \left| \sum_{k=1}^n \psi_j \lambda'_{k-j} K(x_k/h) \right|^2 \end{aligned}$$

$$\begin{aligned} &\leq 2 \sum_{j=m_0}^{\infty} j^{1/4} (|\psi_{1j}| + |\psi_{2j}|) \sum_{j=m_0}^{\infty} j^{-1/4} (|\psi_{1j}| + |\psi_{2j}|) \\ &\quad \left(\mathbb{E} \left| \sum_{k=1}^n \epsilon_{k-j} K(x_k/h) \right|^2 + \mathbb{E} \left| \sum_{k=1}^n e_{k-j} K(x_k/h) \right|^2 \right) \\ &\leq C(nh/d_n) \left[\sum_{j=m_0}^{\infty} j^{1/4} (|\psi_{1j}| + |\psi_{2j}|) \right]^2, \end{aligned}$$

where we employ Hölder’s inequality and (7.40) with $\Lambda(\lambda_k) = \epsilon_k$ and e_k , respectively. The proof of Proposition 7.6 is complete. □

7.3. Proof of Proposition 7.4

We start with (7.4). The tightness of $Z_{n,h}(t)$ has been established in Proposition 7.2. It suffices to show that the finite-dimensional distributions of $Z_{n,h}(t)$ converge to those of $\tau^2 L_Z(t, 0)$. To this end, let $g(x) = \mathbb{E} f^2(x, w_1)$. Under **A2(b)** and **A3(I)**, $g(x)$ is bounded and integrable. Furthermore, by using Wang (2015, Thm. 2.20), we have

$$\frac{d_n}{nh} \sum_{k=1}^{\lfloor nt \rfloor} g(x_k/h) \Rightarrow \tau^2 L_Z(t, 0), \tag{7.41}$$

whenever $d_n/h \rightarrow \infty$ and $d_n/nh \rightarrow 0$. In terms of (7.41), the finite-dimensional distribution of $Z_{n,h}(t)$ will converge to those of $\tau^2 L_Z(t, 0)$ if we show that, for any fixed $0 < t \leq 1$,

$$\frac{d_n}{nh} \sum_{k=1}^{\lfloor nt \rfloor} [g(x_k/h) - f^2(x_k/h, w_k)] = o_P(1). \tag{7.42}$$

This is essentially the same as in the proof of (A.20) for $i = 2$ in Wang et al. (2021) (also see (4.8) in the article) and hence the details are omitted. (7.4) is now proved.

We next prove (7.5). It suffices to show the following:

- (a) for each $0 \leq r \leq m$, $\{\psi_{nr}(t)\}_{n \geq 1}$ is tight on $D[0, 1]$; and
- (b) the finite-dimensional distributions of $\{\psi_{n0}(t), \psi_{n1}(t), \dots, \psi_{nm}(t)\}$ converge to those of $\{G_0, G_1, \dots, G_m\} L_Z(t, 0)$.

The proof of part (a) is simple. Indeed, by noting

$$\begin{aligned} |\psi_{nr}(t) - \psi_{nr}(s)| &\leq \frac{d_n}{n} \sum_{k=\lfloor ns \rfloor + 1}^{\lfloor nt \rfloor} |f(x_k, w_k) f(x_{k+r}, w_{k+r})| \\ &\leq \frac{d_n}{n} \sum_{k=\lfloor ns \rfloor + 1}^{\lfloor nt \rfloor + r} f^2(x_k, w_k) \leq |Z_{n,1}(t) - Z_{n,1}(s)| + o_P(1), \end{aligned}$$

uniformly for $s < t$, the tightness of $\psi_{nr}(t)$ is implied by that of $Z_{n,1}(t)$.

To prove part (b), let $h_r(y) = \mathbb{E} \{f(y, w_0)f(y + x_r, w_r)\}$. We have $h_r(y)$ is bounded and integrable due to assumptions **A2(b)** and **A3(I)**. Hence, as in (7.41),

$$\frac{d_n}{n} \sum_{k=1}^{\lfloor nt \rfloor} [\alpha_0 h_0(x_k) + \dots + \alpha_m h_m(x_k)] \Rightarrow \sum_{r=0}^m \alpha_r G_r L_Z(t, 0),$$

on $D[0, 1]$, for any $(\alpha_0, \dots, \alpha_m) \in R^{m+1}$. The Cramér–Wold theorem now implies that part (b) will follow if we prove

$$|\psi_{nr}(t) - \frac{d_n}{n} \sum_{k=1}^{\lfloor nt \rfloor} h_r(x_k)| = o_P(1), \tag{7.43}$$

for any $r \geq 0$ and any fixed $0 \leq t \leq 1$.⁷

The proof of (7.43) is quite technical, starting with some preliminaries. Let $a_l = \sum_{s=0}^l \phi_s$ and $a_l = 0$ if $l < 0$. With $\gamma = 0$, we may write

$$x_k = \sum_{j=-\infty}^0 [a_{k-j} - a_{-j}] \epsilon_j + \sum_{j=1}^k a_{k-j} \epsilon_j, \tag{7.44}$$

and

$$\begin{aligned} x_{k+r} - x_k &= \sum_{j=-\infty}^k [a_{k+r-j} - a_{k-j}] \epsilon_j + \sum_{j=k+1}^{k+r} a_{k+r-j} \epsilon_j \\ &= \sum_{j=-\infty}^0 [a_{r-j} - a_{-j}] \epsilon_{j+k} + \sum_{j=1}^r a_{r-j} \epsilon_{j+k} \\ &= x_{1k,r} + x_{2k,r}, \end{aligned} \tag{7.45}$$

where

$$\begin{aligned} x_{1k,r} &= \sum_{j=-\infty}^{-A_0} [a_{r-j} - a_{-j}] \epsilon_{j+k}, \\ x_{2k,r} &= \sum_{j=-A_0+1}^0 [a_{r-j} - a_{-j}] \epsilon_{j+k} + \sum_{j=1}^r a_{r-j} \epsilon_{j+k}. \end{aligned}$$

It is readily seen that, for any $A_0 > 0$, $x_{1k,r}$ is independent of $x_{2k,r}$ and $x_{1k,r}$ is independent of w_k and w_{k+r} when $A_0 \geq m_0 + 1$. By letting $\gamma_j = a_{r+j} - a_j$, we further have $\sum_{j=1}^{\infty} \gamma_j^2 < \infty$ and

$$x_{1k,r} = \sum_{j=-\infty}^{-A_0} [a_{r-j} - a_{-j}] \epsilon_{j+k} = \sum_{q=1}^{k-A_0} \gamma_{k-q} \epsilon_q + \sum_{q=-\infty}^0 \gamma_{k-q} \epsilon_q. \tag{7.46}$$

⁷We remark that the r in (7.43) is allowed to depend on n and we have in fact established the convergence in (7.43) in L_1 rather than in probability. These enhanced properties will be useful in the proof of Proposition 7.7.

We next let $\hat{f}(t, s) = \int_{-\infty}^{\infty} e^{itx} f(x, s) dx$,

$$V_k(t, s) = \hat{f}(-t, w_k) \hat{f}(s, w_{k+r}) e^{-isx_{2k,r}},$$

$$A_r(t, s) = \mathbb{E} \{ \hat{f}(-t, w_0) \hat{f}(s, w_r) e^{-isx_r} \}.$$

Using the Fourier transformations, under assumption **A3(III)**, it is readily seen that

$$h_{1r}(y, s) := \frac{1}{2\pi} \int e^{i(t-s)y} \mathbb{E} V_0(t, s) dt = e^{-isy} \mathbb{E} \{ f(y, w_0) \hat{f}(s, w_r) e^{-isx_{20,r}} \},$$

$$h_{2r}(y, s) := \frac{1}{2\pi} \int e^{i(t-s)y} A_r(t, s) dt = e^{-isy} \mathbb{E} \{ f(y, w_0) \hat{f}(s, w_r) e^{-isx_r} \},$$

$$h_r(y) = \mathbb{E} \{ f(y, w_0) f(y + x_r, w_r) \} = \frac{1}{2\pi} \int h_{2r}(y, s) ds.$$

We are now ready to consider (7.43). Without loss of generality, assume $t = 1$. We have

$$\begin{aligned} \psi_{nr}(1) &= \frac{d_n}{2\pi n} \sum_{k=1}^n f(x_k, w_k) \int \hat{f}(s, w_{k+r}) e^{-isx_{k+r}} ds \\ &= \frac{d_n}{(2\pi)^2 n} \sum_{k=1}^n \int \int_{|s| \leq A} \hat{f}(-t, w_k) \hat{f}(s, w_{k+r}) e^{i(t-s)x_k - is(x_{k+r} - x_k)} ds dt + R_{0A} \\ &= \frac{d_n}{(2\pi)^2 n} \sum_{k=1}^n \int \int_{|s| \leq A} e^{i(t-s)x_k - isx_{1k,r}} \mathbb{E} V_k(t, s) ds dt + R_{1A} + R_{0A} \\ &= \frac{d_n}{2\pi n} \sum_{k=1}^n \int_{|s| \leq A} e^{-isx_{1k,r}} h_{1r}(x_k, s) ds + R_{1A} + R_{0A} \\ &= \frac{d_n}{2\pi n} \sum_{k=1}^n \int_{|s| \leq A} e^{-isx_{1k,r}} h_{2r}(x_k, s) ds + R_{2A} + R_{1A} + R_{0A} \\ &= \frac{d_n}{2\pi n} \sum_{k=1}^n \int_{|s| \leq A} h_{2r}(x_k, s) ds + R_{3A} + R_{2A} + R_{1A} + R_{0A} \\ &=: \frac{d_n}{n} \sum_{k=1}^n h_r(x_k) - R_{4A} + R_{3A} + R_{2A} + R_{1A} + R_{0A}, \end{aligned} \tag{7.47}$$

where

$$R_{0A} = \frac{d_n}{2\pi n} \sum_{k=1}^n f(x_k, w_k) \int_{|s| > A} \hat{f}(s, w_{k+r}) e^{-isx_{k+r}} ds,$$

$$R_{1A} = \frac{d_n}{(2\pi)^2 n} \sum_{k=1}^n \int_{|s| \leq A} \int e^{i(t-s)x_k - isx_{1k,r}} [V_k(t, s) - \mathbb{E} V_k(t, s)] dt ds,$$

$$\begin{aligned}
 R_{2A} &= \frac{d_n}{2\pi n} \int_{|s| \leq A} \sum_{k=1}^n e^{-isx_{1k,r}} [h_{1r}(x_k, s) - h_{2r}(x_k, s)] ds, \\
 R_{3A} &= \frac{d_n}{2\pi n} \int_{|s| \leq A} \sum_{k=1}^n (e^{-isx_{1k,r}} - 1) h_{2r}(x_k, s) dt ds \\
 &= \frac{d_n}{(2\pi)^2 n} \sum_{k=1}^n \int_{|s| \leq A} \int e^{i(t-s)x_k} (e^{-isx_{1k,r}} - 1) A_r(t, s) ds, \\
 R_{4A} &= \frac{d_n}{2\pi n} \sum_{k=1}^n \int_{|s| > A} h_{2r}(x_k, s) ds.
 \end{aligned}$$

Recalling w_k depends only on $\lambda_k, \dots, \lambda_{k-m_0}$, where m_0 is a fixed integer, it follows from Lemma 7.1(i) and $|f(y, w_0)| \leq T(y)(1 + \|w_0\|^\beta)$ that

$$\begin{aligned}
 \mathbb{E} |R_{0A}| &\leq C \frac{d_n}{n} \sum_{k=1}^n \int_{|s| > A} \mathbb{E} \{ |f(x_k, w_k)| |\hat{f}(s, w_{k+r})| \} ds \\
 &\leq C \frac{d_n}{n} \sum_{k=1}^n d_k^{-1} \int_{|s| > A} \int \mathbb{E} \{ |f(y, w_0)| |\hat{f}(s, w_r)| \} dy ds \\
 &\leq C \int T(y) dy \int_{|s| > A} \mathbb{E} \{ |\hat{f}(s, w_r)| (1 + \|w_0\|^\beta) \} ds \rightarrow 0,
 \end{aligned}$$

as $A \rightarrow \infty$. Similarly,

$$\begin{aligned}
 \mathbb{E} |R_{4A}| &\leq C \frac{d_n}{n} \sum_{k=1}^n \int_{|s| > A} \mathbb{E} |h_{2r}(x_k, s)| ds \\
 &\leq C \frac{d_n}{n} \sum_{k=1}^n d_k^{-1} \int_{|s| > A} \int |h_{2r}(y, s)| dy ds \\
 &\leq C \int_{|s| > A} \int \mathbb{E} \{ |f(y, w_0)| |\hat{f}(s, w_r)| \} dy ds \rightarrow 0,
 \end{aligned}$$

as $A \rightarrow \infty$. Hence, $|R_{0A}| + |R_{4A}| = o_P(1)$, as $n \rightarrow \infty$ first and then $A \rightarrow \infty$. This, together with (7.47), implies that (7.43) will follow if we prove: for any fixed $A > 0$,

$$R_{jA} = o_P(1), \quad j = 1, 2, 3, \tag{7.48}$$

as $n \rightarrow \infty$ first and then $A_0 \rightarrow \infty$.

The proof of (7.48) for $j = 2$ is simple. Indeed, due to the independence between $x_{10,r}$ and w_1, w_r , we have

$$\begin{aligned}
 &\int_{|s| \leq A} \int |h_{1r}(y, s) - h_{2r}(y, s)| dy ds \\
 &\leq \int_{|s| \leq A} \int \mathbb{E} \{ |f(y, w_0)| |\hat{f}(s, w_r)| |e^{-isx_{10,r}} - 1| \} dy ds
 \end{aligned}$$

$$\begin{aligned} &\leq A \int \int \mathbb{E} \{ |f(y, w_0)| |\hat{f}(s, w_r)| dy ds \mathbb{E} |x_{10, r}| \\ &\leq CA \left[\sum_{j=A_0}^{\infty} (a_{r+j} - a_j)^2 \right]^{1/2}, \end{aligned}$$

for any fixed $A > 0$. This yields that

$$\begin{aligned} \mathbb{E} |R_{2A}| &\leq \frac{d_n}{2\pi n} \int_{|s| \leq A} \sum_{k=1}^n \mathbb{E} |h_{1r}(x_k, s) - h_{2r}(x_k, s)| ds \\ &\leq \frac{d_n}{n} \sum_{k=1}^n d_k^{-1} \int_{|s| \leq A} \int |h_{1r}(y, s) - h_{2r}(y, s)| dy ds \\ &\leq CA \left[\sum_{j=A_0}^{\infty} (a_{r+j} - a_j)^2 \right]^{1/2} \rightarrow 0, \end{aligned}$$

as $n \rightarrow \infty$ first and then $A_0 \rightarrow \infty$, as required.

It is readily seen that (7.48) for $j = 1$ and 3 will follow if we prove: for any fixed $A > 0$,

$$\frac{d_n}{n} \sup_{|s| \leq A} \mathbb{E} \left| \sum_{k=1}^n \int e^{i(u-s)x_k - isx_{1k, r}} [V_k(u, s) - \mathbb{E} V_k(u, s)] du \right| = o(1), \tag{7.49}$$

$$\frac{d_n}{n} \sup_{|s| \leq A} \mathbb{E} \left| \sum_{k=1}^n \int e^{i(u-s)x_k} (e^{-isx_{1k, r}} - 1) A_r(u, s) du \right| = o(1), \tag{7.50}$$

as $n \rightarrow \infty$ first and then $A_0 \rightarrow \infty$.

We first prove (7.50). We may write, for any $B \geq 1$ and $|s| \leq A$,

$$\begin{aligned} &\sum_{k=1}^n \int e^{iux_k} (e^{-isx_{1k, r}} - 1) A_r(u + s, s) du \\ &= \sum_{k=1}^n \left(\int_{|u| \geq B/d_k} + \int_{|u| < B/d_k} \right) e^{iux_k} (e^{-isx_{1k, r}} - 1) A_r(u + s, s) du \\ &= \Delta_{1n}(s) + \Delta_{2n}(s), \quad \text{say.} \end{aligned} \tag{7.51}$$

Recalling $|f(x, y)| \leq T(x)(1 + \|y\|^\beta)$, where $T(x)$ is a bounded and integrable function, we have

$$\sup_{u, s} |A_r(u, s)| \leq \int \int \mathbb{E} \{ |f(x, w_0)| |f(y, w_r)| \} dx dy < \infty, \tag{7.52}$$

$$\begin{aligned} \sup_s \int |A_r(u, s)| du &\leq \int \int \mathbb{E} \{ |\hat{f}(t, w_0)| |f(x, w_r)| \} dt dx \\ &\leq \int T(x) dx \int \mathbb{E} \{ |\hat{f}(t, w_0)| (1 + \|w_r\|^\beta) \} dt < \infty, \end{aligned} \tag{7.53}$$

$$\begin{aligned} \frac{1}{2\pi} \sup_s \int \left| \int A_r(t+s, s) e^{ity} dt \right| dy &= \sup_s \int |h_{2r}(y, s)| dy \\ &\leq C \int \int \mathbb{E} \{ |f(t, w_0) \hat{f}(x, w_r)| \} dt dx < \infty. \end{aligned} \tag{7.54}$$

Due to (7.52), it is readily seen that, uniformly for $|s| \leq A$ and any $B > 0$,

$$\begin{aligned} \mathbb{E} |\Delta_{2n}(s)| &\leq C \sup_{|u|, |s| \leq A} |A_r(u+s, s)| B \sum_{k=1}^n d_k^{-1} \mathbb{E} |x_{1k, r}| \\ &\leq C B n / d_n \left[\sum_{k=A_0}^{\infty} (a_{r+k} - a_k)^2 \right]^{1/2}. \end{aligned} \tag{7.55}$$

To consider $\Delta_{1n}(s)$, writing $\Delta_{1n}(s) = \Delta_{1n,1}(s) + \Delta_{1n,2}(s)$, where

$$\begin{aligned} \Delta_{1n,1}(s) &= \sum_{k=1}^n \int_{|u| \geq B/d_k} e^{iux_k - isx_{1k, r}} A_r(u+s, s) du, \\ \Delta_{1n,2}(s) &= \sum_{k=1}^n \int_{|u| \geq B/d_k} e^{iux_k} A_r(u+s, s) du, \end{aligned}$$

then (7.50) will follow if we prove

$$\frac{d_n}{n} \sup_{|s| \leq A} \mathbb{E} |\Delta_{1n, i}(s)| \leq C (n/d_n) \sqrt{B^{-1} + BA_0^2 d_n/n}, \quad i = 1, 2. \tag{7.56}$$

Indeed, due to (7.51)–(7.56) and $\tau_{A_0} := \sum_{k=A_0}^{\infty} (a_{r+k} - a_k)^2 \rightarrow 0$ as $A_0 \rightarrow \infty$, (7.50) follows by taking $B = \tau_{A_0}^{-1/3}$.

We only prove (7.56) for $i = 1$ as the result for $i = 2$ is similar. We have

$$\begin{aligned} \mathbb{E} |\Delta_{1n,1}(s)|^2 &\leq \sum_{k=1}^n \sum_{j=1}^n \left| \int_{|t| \geq B/d_k} \int_{|u| \geq B/d_j} A_r(t+s, s) A_r(u+s, s) \mathbb{E} T_{kj} dt du \right| \\ &= \left(\sum_{|k-j| \geq A_0^2 + 1} + \sum_{|k-j| \leq A_0^2} \right) \left| \int_{|t| \geq B/d_k} \int_{|u| \geq B/d_j} A_r(t+s, s) A_r(u+s, s) \mathbb{E} T_{kj} dt du \right| \\ &=: \Omega_{1n} + \Omega_{2n}, \quad \text{say,} \end{aligned} \tag{7.57}$$

where $T_{k,j} = e^{itx_k + iux_j} e^{-is(x_{1k, r} + x_{1j, r})}$. Recalling (7.46), it follows that

$$\begin{aligned} &|\mathbb{E}(T_{kj} | \mathcal{F}_0)| \\ &\leq \left| \mathbb{E}(e^{itx_k + iux_j} e^{-is \sum_{q=1}^{k-A_0} \gamma_{k-q} \epsilon_q} e^{-is \sum_{q=1}^{j-A_0} \gamma'_{j-q} \epsilon_q} | \mathcal{F}_0) \right| \\ &= \left| \mathbb{E}(e^{itx_k + iux_j} e^{-i \sum_{q=1}^{k \vee j} s \gamma'_q \epsilon_q} | \mathcal{F}_0) \right|, \end{aligned} \tag{7.58}$$

where

$$\gamma'_q = \begin{cases} \gamma_{k-q} + \gamma_{j-q}, & \text{if } 1 \leq q < k \wedge j, \\ \gamma_{k \vee j - q}, & \text{if } k \wedge j \leq q < k \vee j - A_0, \\ 0, & \text{if } q \geq k \vee j - A_0, \end{cases}$$

satisfying $\sum_{q=1}^\infty \gamma'_q{}^2 < \infty$. Now, by noting (7.52) and using (7.23), we have that, uniformly for $|s| \leq A$,

$$\begin{aligned} \Omega_{1n} &\leq 2 \mathbb{E} \sum_{k-j \geq A_0^2+1} \int_{|t| \geq B/d_k} \int_{|u| \geq B/d_j} |A_r(t+s, s)A_r(u+s, s)| |\mathbb{E}(T_{kj} | \mathcal{F}_0)| dt du \\ &\leq C \sum_{l-k \geq A_0^2+1} [(l-k)^{-2} + B^{-1} d_{l-k}^{-1}] d_k^{-1} \\ &\leq C B^{-1} (n/d_n)^2. \end{aligned}$$

Turning to consider Ω_{2n} , note that

$$\begin{aligned} \mathbb{E} \left| \int_{|t| \geq B/d_k} A_r(t+s, s) e^{itx_k} dt \right| &\leq B/d_k \sup_{t,s} |A_r(t+s, s)| \\ &+ \mathbb{E} \left| \int A_r(t+s, s) e^{itx_k} dt \right| \\ &\leq C B/d_k + C d_k^{-1} \int \left| \int A_r(t+s, s) e^{ity} dt \right| dy \leq C B/d_k, \end{aligned}$$

due to (7.52) and (7.53). Uniformly for $|s| \leq A$, we have

$$\begin{aligned} |\Omega_{2n}| &\leq \sum_{|k-j| \leq A_0^2} \int_{|u| \geq B/d_j} |A_r(u+s, s)| du \mathbb{E} \left| \int_{|t| \geq B/d_k} A_r(t+s, s) T_{kj} dt \right| \\ &\leq \sum_{|k-j| \leq A_0^2} \int_{|u| \geq B/d_j} |A_r(u+s, s)| \mathbb{E} \left| \int_{|t| \geq B/d_k} A_r(t+s, s) e^{itx_k} dt \right| du \\ &\leq C B A_0^2 n/d_n. \end{aligned}$$

Taking this estimate into (7.57), for any fixed $A > 0$, we have

$$\sup_{|s| \leq A} \mathbb{E} |\Delta_{1n,1}(s)|^2 \leq C (B^{-1} + B A_0^2 d_n/n) (n/d_n)^2, \tag{7.59}$$

yielding (7.56). Then (7.50) is established.

Finally, we prove (7.49). Let $\sigma_k(t, s) = V_k(t, s) - \mathbb{E} V_k(t, s)$. Uniformly for $|s| \leq A$ where A is fixed, we have

$$\begin{aligned} \mathbb{E} \left| \sum_{k=1}^n \int e^{itx_k - isx_{1k,r}} \sigma_k(t+s, s) dt \right|^2 \\ = \sum_{k=1}^n \sum_{j=1}^n \mathbb{E} \int \int e^{-is(x_{1k,r} + x_{1j,r})} e^{itx_k + iux_j} \sigma_k(t+s, s) \sigma_j(u+s, s) dt du \end{aligned}$$

$$\begin{aligned}
 &= \left(\sum_{|j-k| \geq A_0^2 + 1} + \sum_{|j-k| \leq A_0^2} \right) \mathbb{E} \int \int e^{-is(x_{1k,r} + x_{1j,r})} e^{itx_k + iux_j} \sigma_k(t + s, s) \sigma_j(u + s, s) dt du \\
 &=: R_{n6} + R_{n7}, \quad \text{say.} \tag{7.60}
 \end{aligned}$$

Note that $\sigma_k(t + s, s)$ depends only on $\epsilon_{k+r}, \dots, \epsilon_{k-A_0}$, $\mathbb{E}\sigma_k(u + s, s) = 0$ and

$$\begin{aligned}
 \sup_{t,s} |\sigma_k(t + s, s)| &\leq C + \sup_t |\hat{f}(t, w_k)| \sup_t |\hat{f}(t, w_{k+r})| \\
 &\leq C(1 + \|w_k\|^{2\beta} + \|w_{k+r}\|^{2\beta}).
 \end{aligned}$$

As in the proof of (7.50), it follows from (7.20) in Lemma 7.4 that

$$\begin{aligned}
 |R_{n6}| &\leq \sum_{|j-k| \geq A_0^2 + 1} \left| \mathbb{E} \int \int e^{-is(x_{1k,r} + x_{1j,r})} e^{itx_k + iux_j} \sigma_k(t, s) \sigma_j(u, s) dt du \right| \\
 &\leq \sum_{|j-k| \geq A_0^2 + 1} \mathbb{E} \int \int \left| \mathbb{E} \left[e^{itx_k + iux_j - is \sum_{q=1}^{k \vee j} \gamma'_q \epsilon_q} \sigma_k(t + s, s) \sigma_j(u + s, s) \mid \mathcal{F}_0 \right] \right| dt du \\
 &\quad \text{(where } \gamma'_q \text{ is given as in (7.58))} \\
 &\leq C \sum_{|j-k| \geq A_0^2 + 1} d_k^{-1} d_{|j-k|}^{-2} \\
 &\leq C \begin{cases} n/d_n, & \text{under LM,} \\ n \log n/d_n, & \text{under SM.} \end{cases} \tag{7.61}
 \end{aligned}$$

To consider R_{n7} , let $l_k(y) = \int e^{ity} \sigma_k(t + s, s) dt$. It is readily seen that

$$\begin{aligned}
 |l_k(y)| &\leq |f(y, w_k)| |\hat{f}(s, w_{k+r})| + \mathbb{E} \{ |f(y, w_k)| |\hat{f}(s, w_{k+r})| \} \\
 &\leq C |f(y, w_k)| (1 + \|w_{k+r}\|^\beta) + C \mathbb{E} \{ |f(y, w_k)| (1 + \|w_{k+r}\|^\beta) \},
 \end{aligned}$$

and by Lemma 7.1,

$$\mathbb{E} |l_k(x_k)|^2 \leq C d_k^{-1} \mathbb{E} (1 + \|w_1\|^{4\beta}) \leq C_1 d_k^{-1}.$$

This yields that

$$|R_{n7}| \leq \sum_{|j-k| \leq A_0^2 + 1} \mathbb{E} \{ |l_k(x_k)| |l_j(x_j)| \} \leq C_1 \sum_{|j-k| \leq A_0^2 + 1} d_k^{-1} \leq C A_0^2 n/d_n. \tag{7.62}$$

It follows from (7.60) to (7.62) that

$$\begin{aligned}
 &\frac{d_n}{n} \mathbb{E} \left| \sum_{k=1}^n \int e^{itx_k - isx_{1k,r}} \sigma_k(t + s, s) dt \right| \\
 &\leq C (A_0^2 + \log n) \left(\frac{d_n}{n} \right)^{1/2} \rightarrow 0,
 \end{aligned}$$

as $n \rightarrow \infty$ first and then $A_0 \rightarrow \infty$. This proves (7.49) and also completes the proof of Proposition 7.4.

7.4. Proof of Proposition 7.7

Recall (6.14) and that $l(x)$ is continuous with $l(0) = 1$. It suffices to show that there exists $M \equiv M_n \rightarrow \infty$ so that, as $n \rightarrow \infty$ first and then $q \rightarrow \infty$,

$$\frac{d_n}{n} \sum_{r=q+1}^M \ell\left(\frac{r}{M}\right) \sum_{k=1}^{n-r} K(x_k)K(x_{k+r}) u_{1k} u_{1,k+r} = o_P(1), \tag{7.63}$$

where $u_{1j}(= u_{1,j}) = \sum_{i=0}^{m_0} \psi_i \lambda'_{j-i}$ for some $m_0 = m_0(n) \rightarrow \infty$ and $m_0 = o(\sqrt{n/d_n})$.

To this end, as in (7.45) and (7.46), for $A_0 = m_0 + 1$, we write

$$x_{k+r} - x_k = x_{1k,r} + x_{2k,r},$$

where, by using the notations $a_l = \sum_{s=0}^l \phi_s$ with $a_l = 0$ if $l < 0$ and $\gamma_l = a_{r+l} - a_l$,

$$x_{1k,r} = \sum_{j=-\infty}^{-A_0} [a_{r-j} - a_{-j}] \epsilon_{j+k} = \sum_{j=1}^{k-A_0} \gamma_{k-j} \epsilon_j + \sum_{j=-\infty}^0 \gamma_{k-j} \epsilon_j,$$

$$x_{2k,r} = \sum_{j=-A_0+1}^0 [a_{r-j} - a_{-j}] \epsilon_{j+k} + \sum_{j=1}^r a_{r-j} \epsilon_{j+k}.$$

Recall that $K(x) = \frac{1}{2\pi} \int e^{itx} \hat{K}(t) dt$ under the condition (a). For any $r \geq 0$ and $l_n \geq 0$, we have

$$\sum_{k=1}^{n-r} K(x_k)K(x_{k+r}) u_{1k} u_{1,k+r}$$

$$= \frac{1}{2\pi} \sum_{k=1}^{n-r} K(x_k) u_{1k} u_{1,k+r} \int_{|s| \leq l_n} \hat{K}(s) e^{-isx_{k+r}} ds + L_{1n}$$

$$= L_{1n}(r) + L_{2n}(r) + L_{3n}(r), \tag{7.64}$$

where, with $V_k(s) = e^{-isx_{2k,r}} u_{1k} u_{1,k+r}$,

$$L_{1n}(r) = \frac{1}{2\pi} \sum_{k=1}^{n-r} K(x_k) u_{1k} u_{1,k+r} \int_{|s| > l_n} \hat{K}(s) e^{-isx_{k+r}} ds,$$

$$L_{2n}(r) = \frac{1}{2\pi} \sum_{k=1}^{n-r} K(x_k) \int_{|s| \leq l_n} \hat{K}(s) e^{-is(x_k+x_{1k,r})} \mathbb{E} V_k(s) ds,$$

$$L_{3n}(r) = \frac{1}{2\pi} \sum_{k=1}^{n-r} K(x_k) \int_{|s| \leq l_n} \hat{K}(s) e^{-is(x_k+x_{1k,r})} [V_k(s) - \mathbb{E} V_k(s)] ds.$$

Using Lemma 7.1(i) and $\int |\hat{K}(s)| ds < \infty$, for any $m_0 \rightarrow \infty$ satisfying $m_0 = O(n/d_n)$, there exists $M_1 = M_{1n} \rightarrow \infty$ so that, whenever $l_n \rightarrow \infty$,

$$\begin{aligned} & \frac{d_n}{n} \sum_{r=q+1}^{M_1} \mathbb{E}|L_{1n}(r)| \\ & \leq C \sum_{r=q+1}^{M_1} \frac{d_n}{n} \left[\sum_{k=1}^{3m_0} \mathbb{E}|u_{1k} u_{1,k+r}| + \sum_{3m_0+1}^n d_k^{-1} \right] \int_{|s|>l_n} |\hat{K}(s)| ds \\ & \leq CM_1 \int_{|s|>l_n} |\hat{K}(s)| ds \rightarrow 0. \end{aligned} \tag{7.65}$$

To estimate $L_{2n}(r)$, let $\tilde{h}_r(y) = \mathbb{E}[K(y + x_{20,r})u_{10}u_{1r}]$. It is readily seen that $\tilde{h}_r(y)$ is bounded and integrable. Furthermore, using (7.33) in Lemma 7.7 with minor modifications, we have

$$\begin{aligned} |\tilde{h}_r(y)| & \leq \sum_{l=0}^{m_0} \sum_{v=0}^{m_0} |\mathbb{E}[K(y + x_{20,r})\psi_l \lambda'_{-l} \psi_v \lambda'_{r-v}]| \\ & \leq C \sum_{l=0}^{m_0} \sum_{v=0}^{m_0} \|\psi_l\| \|\psi_v\| \begin{cases} d_r^{-1}, & \text{if } |r-v+l| \leq 1, \\ \sum_{s=0}^v |\phi_s| \sum_{s=l}^{r+l} |\phi_s| (d_r^{-3} + r^{-3}), & \text{if } |r-v+l| \geq 2, \end{cases} \\ & \leq C \sum_{l=0}^{m_0} \|\psi_l\| \sum_{v=r+l-1}^{r+l+1} \|\psi_v\| d_r^{-1} + C \sum_{l=0}^{m_0} \sum_{v=0}^{m_0} l^{1/2} \|\psi_l\| v^{1/2} \|\psi_v\| d_r^{-3/2} \\ & \leq Cr^{-1} d_r^{-1} + Cr^{-3/2} \leq Cr^{-3/2}, \end{aligned}$$

uniformly in $y \in R$, where we have used the facts that $d_r^{-1} \leq Cr^{-1/2}$ and $\sum_{s=0}^v |\phi_s| \sum_{s=l}^{r+l} |\phi_s| (d_r^{-3} + r^{-3}) \leq Cv^{1/2} l^{1/2} r^{-3/2}$ under both SM and LM and $\sum_{v=0}^\infty v \|\psi_v\| < \infty$. Now, by noting that $\mathbb{E}V_k(s) = \mathbb{E}V_0(s)$, $\sup_s \mathbb{E}|V_0(s)| \leq \mathbb{E}|u_{10} u_{1r}| \leq C < \infty$ and

$$\tilde{h}_r(y) = \frac{1}{2\pi} \int \hat{K}(s) e^{-isy} \mathbb{E}V_0(s) ds,$$

standard calculations, together with the Hölder inequality, show⁸ that

$$\begin{aligned} \frac{d_n}{n} \mathbb{E}|L_{2n}(r)| & \leq \frac{d_n}{n} \sum_{k=1}^n \mathbb{E}[K(x_k) |\tilde{h}_r(x_k + x_{1k,r})|] + C \frac{d_n}{n} \sum_{k=1}^n \mathbb{E}K(x_k) \int_{|s|>l_n} |\hat{K}(s)| ds \\ & \leq \left[\frac{d_n}{n} \sum_{k=1}^n \mathbb{E}K^{4/3}(x_k) \right]^{3/4} \left[\frac{d_n}{n} \sum_{k=1}^n \mathbb{E}|\tilde{h}_r(x_k + x_{1k,r})|^4 \right]^{1/4} \\ & \quad + C \int_{|s|>l_n} |\hat{K}(s)| ds \end{aligned}$$

⁸Note that $x_k + x_{1k,r} = \sum_{j=-\infty}^k \tilde{a}_{k-j} \epsilon_j$ where $\tilde{a}_{k-j} = a_{k-j} + \gamma_{k-j} I(j \leq k - A_0)$ if $j \geq 1$ and $\tilde{a}_{k-j} = a_{k-j} - a_{-j} + \gamma_{k-j}$ if $j \leq 0$, and $\sum_{j=-\infty}^n \tilde{a}_j^2 \asymp d_n^2$. Lemma 7.1 still holds when the x_k is replaced by $x_k + x_{1k,r}$.

$$\begin{aligned} &\leq C \left[\int K^{4/3}(y) dy \right]^{3/4} \left[\int |\tilde{h}_r(y)|^4 dy \right]^{1/4} + C \int_{|s|>l_n} |\hat{K}(s)| ds \\ &\leq Cr^{-9/8} + C \int_{|s|>l_n} |\hat{K}(s)| ds. \end{aligned}$$

As a consequence, for any $l_n \rightarrow \infty$ and $M_1 \rightarrow \infty$ as given in (7.65), we have

$$\begin{aligned} \frac{d_n}{n} \sum_{r=q+1}^{M_1} \mathbb{E}|L_{2n}(r)| &\leq C \sum_{r=q+1}^{M_1} r^{-9/8} + CM_1 \int_{|s|>l_n} |\hat{K}(s)| ds \\ &\leq Cq^{-1/8} + CM_1 \int_{|s|>l_n} |\hat{K}(s)| ds \rightarrow 0, \end{aligned} \tag{7.66}$$

as $n \rightarrow \infty$ first and then $q \rightarrow \infty$.

We finally estimate $L_{3n}(r)$. It follows from the Fourier transformation that

$$\begin{aligned} L_{3n}(r) &= \frac{1}{(2\pi)^2} \sum_{k=1}^{n-r} \int \int_{|s| \leq l_n} \hat{K}(-t) \hat{K}(s) e^{i(t-s)x_k} e^{-isx_{1k,r}} [V_k(s) - \mathbb{E}V_k(s)] ds dt \\ &= \frac{1}{2\pi} \int_{|s| \leq l_n} \hat{K}(s) \mathcal{L}_n(s, r) ds, \end{aligned} \tag{7.67}$$

where $\mathcal{L}_n(s, r) = \sum_{k=1}^{n-r} \int \hat{K}(-t) e^{i(t-s)x_k} e^{-isx_{1k,r}} [V_k(s) - \mathbb{E}V_k(s)] dt$. Let $\sigma_k(s) = V_k(s) - \mathbb{E}V_k(s)$. Uniformly, for $|s| \leq l_n$, we have

$$\begin{aligned} \mathbb{E}\mathcal{L}_n^2(s, r) &= \mathbb{E} \left| \sum_{k=1}^n \int \hat{K}(t+s) e^{itx_k - isx_{1k,r}} \sigma_k(s) dt \right|^2 \\ &= \sum_{k=1}^n \sum_{j=1}^n \mathbb{E} \int \int \hat{K}(t+s) \hat{K}(u+s) e^{-is(x_{1k,r} + x_{1j,r})} e^{itx_k + iux_j} \sigma_k(s) \sigma_j(s) dt du \\ &= \left(\sum_{|j-k| \geq A_0^2 + 1} + \sum_{|j-k| \leq A_0^2} \right) \mathbb{E} \int \int \hat{K}(t+s) \hat{K}(u+s) e^{-is(x_{1k,r} + x_{1j,r})} \\ &\quad \times e^{itx_k + iux_j} \sigma_k(s) \sigma_j(s) dt du \\ &=: R_{n1}(s) + R_{n2}(s). \end{aligned} \tag{7.68}$$

Note that $\sigma_k(s)$ depends only on $\epsilon_{k+r}, \dots, \epsilon_{k-A_0}$, $\mathbb{E}\sigma_k(s) = 0$ and

$$\sup_s |\sigma_k(s)| \leq C(1 + |u_{1k}| |u_{1,k+r}|).$$

As in the proof of (7.50), it follows from (7.20) in Lemma 7.4 that

$$\begin{aligned}
 |R_{n1}(s)| &\leq \sum_{|j-k| \geq A_0^2+1} \left| \mathbb{E} \int \int e^{-is(x_{1k,r} + x_{1j,r})} e^{itx_k + iux_j} \sigma_k(s) \sigma_j(s) dt du \right| \\
 &\leq \sum_{|j-k| \geq A_0^2+1} \mathbb{E} \int \int \left| \mathbb{E} \left[e^{itx_k + iux_j - is \sum_{q=1}^{k \vee j} \gamma'_q \epsilon_q} \sigma_k(s) \sigma_j(s) \mid \mathcal{F}_0 \right] \right| dt du \\
 &\quad \text{(where } \gamma'_q \text{ is given as in (7.58))} \\
 &\leq C \sum_{|j-k| \geq A_0^2+1} d_k^{-1} d_{|j-k|}^{-2} (1 + |s|) \\
 &\leq C(1 + |s|) \begin{cases} n/d_n, & \text{under LM,} \\ n \log n/d_n, & \text{under SM.} \end{cases} \tag{7.69}
 \end{aligned}$$

As for $R_{n2}(s)$, by recalling $K(x) = \frac{1}{2\pi} \int \hat{K}(t) e^{itx} dx$ and $A_0 = m_0 + 1$, we have

$$\begin{aligned}
 |R_{n2}(s)| &\leq \sum_{|j-k| \leq A_0^2+1} \mathbb{E} \left[K(x_k) K(x_j) \sup_s |\sigma_k(s)| \sup_s |\sigma_j(s)| \right] \\
 &\leq C_1 \sum_{|j-k| \leq A_0^2+1} d_k^{-1} \leq C m_0^2 n/d_n. \tag{7.70}
 \end{aligned}$$

It follows from (7.67) to (7.70) that, for any $l_n \rightarrow \infty$ satisfying $l_n = o(\sqrt{n/d_n})$ and $m_0 = o(\sqrt{n/d_n})$, there exists $M_2 \equiv M_{2n} \rightarrow \infty$,

$$\begin{aligned}
 &\frac{d_n}{n} \sum_{r=q+1}^{M_2} \mathbb{E} |L_{3n}(r)| \\
 &\leq C M_2 \sup_{|s| \leq l_n} \mathbb{E} |\mathcal{L}_n(s, r)| \int_{|s| \leq l_n} |\hat{K}(s)| ds \leq C M_2 \sup_{|s| \leq l_n} [\mathbb{E} \mathcal{L}_n^2(s, r)]^{1/2} \\
 &\leq C M_2 [l_n(1 + \log n) + m_0^2]^{1/2} \left(\frac{d_n}{n}\right)^{1/2} \rightarrow 0. \tag{7.71}
 \end{aligned}$$

By virtue of (7.64), (7.65), (7.66), and (7.71), for any $M \equiv M_n \rightarrow \infty$ and $M_n \leq \min\{M_{1n}, M_{2n}\}$, we have

$$\begin{aligned}
 &\frac{d_n}{n} \sum_{r=q+1}^M \ell\left(\frac{r}{M}\right) \mathbb{E} \left| \sum_{k=1}^{n-r} K(x_k) K(x_{k+r}) u_{1k} u_{1,k+r} \right| \\
 &\leq C \frac{d_n}{n} \sum_{r=q+1}^{M_1} (\mathbb{E} |L_{1n}(r)| + \mathbb{E} |L_{2n}(r)|) + \frac{C d_n}{n} \sum_{r=q+1}^{M_2} \mathbb{E} |L_{3n}(r)| \rightarrow 0,
 \end{aligned}$$

as $n \rightarrow \infty$ first and then $q \rightarrow \infty$. This proves (7.63) and completes the proof of Proposition 7.7.

A. APPENDIX: PROOFS OF LEMMAS

A.1. Proof of Lemma 7.2

The idea of the proof is similar to that of Wang and Phillips (2011, pp. 246–247) and the following outline is provided here for completeness. We first prove (7.13). Write $\Omega_1 \equiv \Omega_1(t)$ (Ω_2 , respectively) for the set of $1 \leq v \leq k$ such that $|t\alpha_v + \beta_v| \geq 1$ ($|t\alpha_v + \beta_v| < 1$, respectively), and

$$B_2 = \sum_{v \in \Omega_2} \alpha_v \beta_v \quad \text{and} \quad B_3 = \sum_{v \in \Omega_2} \beta_v^2.$$

Since $B_2^2 \leq B_{1k} B_3$ by Hölder’s inequality, we have

$$\begin{aligned} \sum_{q \in \Omega_2} (t\alpha_q + \beta_q)^2 &= t^2 B_{1k} + 2tB_2 + B_3 \\ &= B_{1k}(t + B_2/B_{1k})^2 + (B_3 - B_2^2/B_{1k}) \geq B_{1k}(t + B_2/B_{1k})^2. \end{aligned}$$

On the other hand, there exist constants $\gamma_1 > 0$ and $\gamma_2 > 0$ such that

$$|\mathbb{E} e^{i\epsilon_1 l}| \leq \begin{cases} e^{-\gamma_1}, & \text{if } |l| \geq 1, \\ e^{-\gamma_2 l^2}, & \text{if } |l| \leq 1, \end{cases} \tag{A.1}$$

since $\mathbb{E}\epsilon_1 = 0$, $\mathbb{E}\epsilon_1^2 < \infty$ and ϵ_1 satisfies the Cramér’s condition due to $\int |\mathbb{E} e^{it\epsilon_0}| dt < \infty$ (see, e.g., Chapter 1 of Petrov (1995)). Without loss of generality, assume $\alpha_1 \neq 0$ and let $g(t) = \mathbb{E} e^{it\alpha_1 \epsilon_0}$. From these facts and the independence of ϵ_t , it follows that, for k sufficiently large and all t ,

$$\begin{aligned} |\mathbb{E} e^{iz_k(t)}| &\leq |g(t)| \prod_{q=2}^k |\mathbb{E} e^{i\epsilon_1(t\alpha_q + \beta_q)}| \\ &\leq |g(t)| \exp \left\{ -\gamma_1 \#(\Omega_1) - \gamma_2 \sum_{v \in \Omega_2} (t\alpha_v + \beta_v)^2 \right\} \\ &\leq |g(t)| \exp \left\{ -\gamma_1 \#(\Omega_1) - \gamma_2 B_{1k} (t + B_2/B_{1k})^2 \right\}. \end{aligned} \tag{A.2}$$

Hence, by recalling (7.12) and $|B_2| \leq \sum_{v=1}^k |\alpha_v \beta_v|$, simple calculations show that

$$\begin{aligned} &\int \min\{1, s_1 |t|^\delta + s_2\} |\mathbb{E} e^{iz_k(t)}| dt \\ &\leq \int_{\#(\Omega_1) \geq \sqrt{k}} |g(t)| e^{-\sqrt{k}} dt + C \int_{\#(\Omega_1) \leq \sqrt{k}} (s_1 |t|^\delta + s_2) e^{-\gamma_2 B_{1k} (t + B_2/B_{1k})^2} dt \\ &\leq C e^{-\sqrt{k}} + C s_1 \int (|t| + |B_2|/B_{1k})^\delta e^{-\gamma_2 B_{1k} t^2} I(B_{1k} \geq m_k^2) dt \\ &\quad + C s_2 \int e^{-\gamma_2 B_{1k} t^2} I(B_{1k} \geq m_k^2) dt \\ &\leq C (k^{-3} + s_1 [m_k^{-1-\delta} + m_k^{-1-2\delta} (\sum_{v=1}^k |\alpha_v \beta_v|)^\delta] + s_2 m_k^{-1}). \end{aligned}$$

Result (7.13) now follows from the fact that

$$\sum_{\nu=1}^k |\alpha_\nu \beta_\nu| \leq \left(\sum_{\nu=1}^k |\alpha_\nu|^2\right)^{1/2} \left(\sum_{\nu=1}^k |\beta_\nu|^2\right)^{1/2} \leq Cm_k \left(\sum_{\nu=1}^k |\beta_\nu|^2\right)^{1/2}.$$

The proof of (7.14) is similar and hence the details are omitted. We finally prove (7.15). In fact, by recalling $B_2^2/B_{1k} \leq B_3 \leq a$, i.e. $B_2/B_{1k} \leq a^{1/2}/m_k$ due to (7.12), it follows from (A.2) that

$$\begin{aligned} & \int_{|t| \geq B/m_k} |\mathbb{E} e^{iz_k(t)}| dt \\ & \leq \int_{\#\{\Omega_1\} \geq \sqrt{k}} |g(t)| e^{-\sqrt{k}} dt + C \int_{\#\{\Omega_1\} \leq \sqrt{k}, |t| \geq B/m_k} e^{-\gamma_2 B_{1k}(t+B_2/B_{1k})^2} dt \\ & \leq Ck^{-3} + \int_{|t| \geq 2^{-1}B/m_k} e^{-\gamma_2 B_{1k} t^2} I(B_{1k} \geq m_k^2) dt \\ & \leq C(k^{-3} + m_k^{-1} B^{-1}), \end{aligned}$$

as required.

A.2. Proof of Lemma 7.3

Let $V_k(t) = \sum_{\nu=k-m_0+1}^k (t\alpha_\nu + \beta_\nu)\epsilon_\nu$. Note that

$$\begin{aligned} |\mathbb{E} e^{iz_k(t/h)} g(t, w_k)| & \leq |\mathbb{E} e^{iz_{k-m_0}(t/h)}| |\mathbb{E} e^{iV_k(t/h)} g(t, w_k)| \\ & \leq \mathbb{E} |g(t, w_1)| |\mathbb{E} e^{iz_{k-m_0}(t/h)}|. \end{aligned}$$

It follows from (7.13) with $s_1 = 0$ and $s_2 = 1$ that

$$\int |\mathbb{E} e^{iz_k(t/h)} g(t, w_k)| dt \leq Ch \int |\mathbb{E} e^{iz_{k-m_0}(t)}| dt \leq Ch(k^{-3} + \tau_k^{-1}),$$

yielding (7.16). Similarly, by noting that

$$\begin{aligned} |\mathbb{E} e^{iV_k(t/h)} g(t, w_k)| & \leq |\mathbb{E} (e^{iV_k(t/h)} - 1)g(t, w_k)| + |\mathbb{E} g(t, w_k)| \\ & \leq 2 \min\{1, \alpha_{k0} |t|/h + \beta_{k0}\} \mathbb{E} \{| \epsilon_0 | |g(t, w_1)|\} + C \min\{1, |t|\} \\ & \leq C \beta_{k0} + C \min\{1, \alpha_{k0} |t|/h\} + C \min\{1, |t|\}, \end{aligned} \tag{A.3}$$

we have

$$\begin{aligned} & \int |\mathbb{E} \{e^{iz_k(t/h)} g(t, w_k)\}| dt \\ & \leq C \int \min\{1, \alpha_{k0} |t|/h\} |\mathbb{E} e^{iz_{k-m_0}(t/h)}| dt + C\beta_{k0} \int |\mathbb{E} e^{iz_{k-m_0}(t/h)}| dt \\ & \quad + C \int \min\{1, |t|\} |\mathbb{E} e^{iz_{k-m_0}(t/h)}| dt \\ & \leq Ch \left\{ (1 + \alpha_{k0}) \tau_k^{-2} \left[1 + \left(\sum_{\nu=1}^k \beta_\nu^2 \right)^{1/2} \right] + \beta_{k0} \tau_k^{-1} \right\}, \end{aligned}$$

as required in (7.17). As for (7.18), by noting that

$$|\mathbb{E} e^{iV_k(t/h)} g(t, w_k)| \leq C \beta_{k0} + C \min\{1, |t|\} + C \min\{1, |t|/h\},$$

due to (A.3) and $\sup_k \alpha_{k0} = O(1)$, it follows from (7.13) and (7.14) that

$$\begin{aligned} & \int \min\{1, |t|/h\} |\mathbb{E} \{e^{iz_k(t/h)} g(t, w_k)\}| dt \\ & \leq C \beta_{k0} \int \min\{1, |t|/h\} |\mathbb{E} e^{iz_k-m_0(t/h)}| dt + C \int \min\{1, (|t|/h)^2\} |\mathbb{E} e^{iz_k-m_0(t/h)}| dt \\ & \quad + C \int \min\{1, |t|\} \min\{1, |t|/h\} |\mathbb{E} e^{iz_k-m_0(t/h)}| dt \\ & \leq Ch \{k^{-3} + [\beta_{k0}(\tau_k^{-2} + k^{-3}) + \tau_k^{-3}](1 + \sum_{v=1}^k \beta_v^2)\}. \end{aligned}$$

This proves (7.18).

A.3. Proof of Lemma 7.4

We only prove (7.20) and (7.22). The other proofs are similar and simpler. Note that

$$\begin{aligned} x_k &= \sum_{j=1}^k \rho_n^{k-j} \xi_j = \sum_{j=1}^k \rho_n^{k-j} \left(\sum_{u=1}^j + \sum_{u=-\infty}^0 \right) \epsilon_u \phi_{j-u} \\ &= \sum_{u=1}^k \epsilon_u a_{k-u} + \sum_{u=0}^{\infty} \epsilon_{-u} b_{u,k}, \end{aligned} \tag{A.4}$$

where $a_{k-u} = \sum_{s=0}^{k-u} \rho_n^{k-u-s} \phi_s$ and $b_{u,k} = \sum_{s=1}^k \rho_n^{k-s} \phi_{s+u}$. It follows from the independence of the ϵ_j that

$$\begin{aligned} & |I_{k,l}(m)| \\ & \leq \int \int |\mathbb{E} \left\{ e^{i s \sum_{v=m+1}^k a_{k-v} \epsilon_v / h + i t \sum_{v=m+1}^l a_{l-v} \epsilon_v / h + i \sum_{j=m+1}^l \gamma_j \epsilon_j} g(s, w_k) g(t, w_l) \right\}| ds dt \\ & \leq C \int |\mathbb{E} \left\{ e^{i \sum_{v=k+1}^l (t a_{l-v} / h + \gamma_v) \epsilon_v} g(t, w_l) \right\}| \Lambda(t, k) dt, \end{aligned} \tag{A.5}$$

where

$$\Lambda(t, k) = \int |\mathbb{E} \left\{ e^{i \sum_{v=m+1}^k (s a_{k-v} / h + t a_{l-v} / h + \gamma_v) \epsilon_v} g(s, w_k) \right\}| ds.$$

As in Lemma 7.2, denote by Ω_1 a subset of $\Omega = \{m + 1, 2, \dots, k\}$ and $\Omega_2 = \Omega - \Omega_1$. Note that, for any $k - m \geq 1$, $\sum_{v \in \Omega_2} a_{k-v}^2 \asymp d_{k-m}^2$ whenever $\#\Omega_1 \leq \sqrt{k - m}$. It is readily seen from (7.16) with $\alpha_v = a_{k-v}$ and $\beta_v = t a_{l-v} / h + \gamma_v$ that

$$\Lambda(t, k) \leq Ch d_{k-m}^{-1}. \tag{A.6}$$

By similar arguments, it follows from (7.17) with $\alpha_v = a_{l-v}$ and $\beta_v = \gamma_v$ that

$$\begin{aligned} & \int |\mathbb{E} \{ e^{i \sum_{v=k+1}^l (ta_{l-v}/h + \gamma_v) \epsilon_v} g(t, w_l) \}| dt \\ & \leq Ch \{ (l-k)^{-3} + \alpha_{l0} d_{l-k}^{-2} [1 + (\sum_{v=k+1}^l \gamma_v^2)^{1/2}] + \beta_{l0} d_{l-k}^{-1} \} \\ & \leq Ch [d_{l-k}^{-2} (1 + a^{1/2}) + \beta_{l0} d_{l-k}^{-1}], \end{aligned} \tag{A.7}$$

where $a = \sum_{v=1}^l \gamma_v^2$, $\beta_{l0} = \max_{0 \leq j \leq m_0} |\gamma_{l-j}|$ and we have used the fact:

$$\alpha_{l0} = \max_{0 \leq i \leq m_0} |\alpha_{l-i}| = \max_{0 \leq i \leq m_0} |a_i| = O(1).$$

It follows from (A.5) to (A.7) that

$$\begin{aligned} |I_{k,l}(m)| & \leq Ch d_{k-m}^{-1} \int |\mathbb{E} e^{i \sum_{v=k+1}^l (ta_{l-v}/h + \gamma_v) \epsilon_v} g(t, w_l)| dt \\ & \leq Ch^2 d_{k-m}^{-1} [d_{l-k}^{-2} (1 + a^{1/2}) + \beta_{l0} d_{l-k}^{-1}], \end{aligned}$$

implying (7.20).

The proof of (7.22) requires some modifications. First notice that, under *SM*, we have

$$\Lambda(t, k) \leq Ch [(k-m)^{-1} + \min\{1, |t|/h\} (k-m)^{-1/2}], \tag{A.8}$$

rather than (A.6). Indeed, under *SM*, it follows that:

- (a) $\Lambda(t, k) \leq Ch(k-m)^{-1/2}$ by (7.16) and, for any $t \in R$,
- (b) $\Lambda(t, k) \leq Ch [(k-m)^{-1} + |t|/h (k-m)^{-1/2}]$ by (7.17) with $\alpha_v = a_{k-v}$ and $\beta_v = ta_{l-v}/h + \mu_v/\sqrt{n}$,

implying (A.8). Now, by using (A.5) first and then (7.17) and (7.18), we have

$$\begin{aligned} & |I_{k,l}(m)| \\ & \leq Ch(k-m)^{-1} \int |\mathbb{E} \{ e^{i \sum_{v=k+1}^l (ta_{l-v}/h + \gamma_v) \epsilon_v} g(t, w_l) \}| dt \\ & + Ch(k-m)^{-1/2} \int \min\{1, |t|/h\} |\mathbb{E} \{ e^{i \sum_{v=k+1}^l (ta_{l-v}/h + \gamma_v) \epsilon_v} g(t, w_l) \}| dt \\ & \leq Ch^2 [(l-k)^{-1} (k-m)^{-1} + (l-k)^{-3/2} (k-m)^{-1/2}], \end{aligned}$$

which yields (7.22).

A.4. Proof of Lemma 7.5

We only prove (7.25). The other proofs are similar and use the corresponding results in Lemma 7.4. Recalling (2.4), we may write

$$II_{lk}(h) = \frac{1}{(2\pi)^2} \int \int \hat{f}(t, w_k) \hat{f}(s, w_l) e^{itx_k/h + isx_l/h} e^{i \sum_{j=m+1}^n \mu_j \epsilon_j / \sqrt{n}} dt ds. \tag{A.9}$$

It follows from (A.4), the independence of ϵ_j and (7.20) with $\gamma_j = \mu_j/\sqrt{n}$ and $g(s, w_k) = \hat{f}(s, w_k)$ that

$$\begin{aligned} & |\mathbb{E}[II_k(h) \mid \mathcal{F}_m]| \\ & \leq \frac{1}{(2\pi)^2} \int \int \mathbb{E}(e^{isx_k/h + itx_l/h + i \sum_{j=m+1}^l \mu_j \epsilon_j / \sqrt{n}} \hat{f}(s, w_k) \hat{f}(t, w_l) \mid \mathcal{F}_m) ds dt \\ & \leq Ch^2 d_{k-m}^{-1} (d_{l-k}^{-2} + d_{l-k}/\sqrt{n}), \end{aligned}$$

as required.

A.5. Proof of Lemma 7.6

Recalling (A.4), as in (A.9), we have

$$\begin{aligned} & |\mathbb{E}\{f(x_l/h, w_l)\mathbb{E}[f(x_k/h, w_k) \mid \mathcal{F}_{k-m}]\}| \\ & = \int \int |\mathbb{E}\{e^{itx_l/h} \hat{f}(-t, w_l)\mathbb{E}[e^{isx_k/h} \hat{f}(s, w_k) \mid \mathcal{F}_{k-m}]\}| ds dt \\ & \leq \int \int |\mathbb{E}\{e^{ith^{-1} \sum_{v=k}^l a_{l-v} \epsilon_v} \hat{f}(-t, w_l)\}| \\ & \quad \mathbb{E}[e^{(ish^{-1} \sum_{v=1}^k a_{k-v} \epsilon_v + ith^{-1} \sum_{v=1}^{k-m} a_{l-v} \epsilon_v)} \hat{f}(s, w_k)] ds dt \\ & \leq C \int |\mathbb{E}\{e^{ith^{-1} \sum_{v=k}^l a_{l-v} \epsilon_v} \hat{f}(-t, w_l)\}| \Lambda(t, k) dt, \end{aligned}$$

where, by letting $a_{l-v}^* = 0$ if $k - m + 1 \leq v \leq k$ and $a_{l-v}^* = a_{l-v}$ if $1 \leq v \leq k - m$, we have

$$\Lambda(t, k) = \int |\mathbb{E}\{e^{i \sum_{v=1}^k (sa_{k-v}/h + ta_{l-v}^*/h) \epsilon_v} \hat{f}(s, w_k)\}| ds.$$

The remainder of the proof is the same as that of Lemma 7.4 and is omitted.

A.6. Proof of Lemma 7.7

Take A_0 as required in Lemma 7.1. Recalling $K(x)$ is bounded, (7.30) follows immediately from Lemma 7.1(i). If $k \geq A_0$, $l - k \geq A_0$, and $l - j \leq k$, it follows from Lemma 7.1(ii) and the conditional arguments that

$$\begin{aligned} I & := |\mathbb{E}\{\Gamma(\lambda_{k-j})\Gamma(\lambda_{l-j})K(x_k/h)K(x_l/h)\}| \\ & \leq \mathbb{E}\{|\Gamma(\lambda_{k-j})\Gamma(\lambda_{l-j})K(x_k/h)| |\mathbb{E}[K(x_l/h) \mid \mathcal{F}_l]|\} \\ & \leq C\mathbb{E}\Gamma^2(\lambda_1)h^2 d_k^{-1} d_{l-k}^{-1}, \end{aligned}$$

indicating (7.31).

We next assume that $k \geq A_0$, $l - k \geq A_0$, and $l - j > k$. Recalling (A.4), as in (A.9), we have

$$I = \int \int |\mathbb{E} \left\{ e^{itx_l/h} e^{isx_k/h} \Gamma(\lambda_{k-j}) \Gamma(\lambda_{l-j}) \right\} | |\hat{K}(-s)| |\hat{K}(-t)| ds dt$$

$$\leq C \int |\mathbb{E} \left\{ e^{ith^{-1} \sum_{v=k}^l a_{l-v} \epsilon_v} \Gamma(\lambda_{l-j}) \right\} | \Lambda(t, k) dt,$$

where

$$\Lambda(t, k) = \begin{cases} \int |\mathbb{E} \left\{ e^{i \sum_{v=1}^k (sa_{k-v}/h + ta_{l-v}/h) \epsilon_v} e^{-i(s\epsilon_{k-j} b_{j-k, k/h} + t\epsilon_{l-k} b_{l-k, k/h})} \Gamma(\lambda_{k-j}) \right\} | ds, & \text{if } k - j \leq 0, \\ \int |\mathbb{E} \left\{ e^{i \sum_{v=1}^k (sa_{k-v}/h + ta_{l-v}/h) \epsilon_v} \Gamma(\lambda_{k-j}) \right\} | ds, & \text{if } k - j \geq 1. \end{cases}$$

It follows from arguments similar to those given in the proof of Lemma 7.4 with some minor modifications⁹ that:

(a) under *LM*, $\Lambda(t, k) \leq Ch d_k^{-1}$ and

$$I \leq Ch d_k^{-1} \int |\mathbb{E} \left\{ e^{ith^{-1} \sum_{v=k}^l a_{l-v} \epsilon_v} \Gamma(\lambda_{l-j}) \right\} | dt \leq C \sum_{s=0}^j |\phi_s| h^2 d_k^{-1} d_{l-k}^{-2};$$

(b) under *SM* (noting $|b_{j-m, m}| \leq \sum_{i=j-m}^j |\phi_i| \leq C < \infty$ for any $m \geq 0$ and $\max_{1 \leq v \leq k} |a_v| \leq C < \infty$),

$$\Lambda(t, k) \leq \int |\mathbb{E} \left\{ e^{i \sum_{v=1, v \neq k-j}^k (sa_{k-v}/h + ta_{l-v}/h) \epsilon_v} \right\} | (\min\{1, |s|/h\} + \min\{1, |t|/h\}) ds$$

$$\leq Ch (k^{-1} + \min\{1, |t|/h\} k^{-1/2})$$

and

$$I \leq Ch k^{-1} \int |\mathbb{E} \left\{ e^{ith^{-1} \sum_{v=k}^l a_{l-v} \epsilon_v} \Gamma(\lambda_{l-j}) \right\} | dt$$

$$+ Ch k^{-1/2} \int \min\{1, |t|/h\} |\mathbb{E} \left\{ e^{ith^{-1} \sum_{v=k}^l a_{l-v} \epsilon_v} \Gamma(\lambda_{l-j}) \right\} | dt$$

$$\leq Ch^2 k^{-1} (l - k)^{-1} + Ch^2 k^{-1/2} (l - k)^{-3/2}.$$

This proves (7.32).

Similarly, by letting $z_{2r} = \sum_{k=1, k \neq r-j}^r \epsilon_k a_{r-k}$, we have

$$|\mathbb{E} \left\{ \Gamma(\lambda_{r-j}) \Gamma(\lambda_{-k}) e^{isx_r/h} \right\} |$$

$$\leq C |\mathbb{E} e^{isz_{2r}/h} | \begin{cases} 1, & \text{if } |r - j + k| \leq A_0, \\ |a_j| |a_{r+k} - a_k| \min\{1, |s|^2\}, & \text{if } |r - j + k| > A_0, \end{cases}$$

⁹Replace m_0 by j , set $\gamma_v = 0$ and take $m = 0$. In this case, α_0 used in (A.7) satisfies

$$\alpha_0 = \max_{0 \leq i \leq j} |\alpha_{l-i}| = \max_{0 \leq i \leq j} |a_i| \leq \sum_{s=0}^j |\phi_s|,$$

which can not be eliminated.

implying that, uniformly for $y \in R$,

$$\begin{aligned} & \left| \mathbb{E} \left\{ K(y + x_l/h) \Gamma(\lambda_{l-j}) \Gamma(\lambda_{-k}) \right\} \right| \\ & \leq \int |\hat{K}(s)| \left| \mathbb{E} \left\{ e^{isx_l/h} \Gamma(\lambda_{l-j}) \Gamma(\lambda_{-k}) \right\} \right| ds \\ & \leq Ch \begin{cases} d_l^{-1}, & \text{if } |l-j+k| \leq A_0, \\ \sum_{s=0}^j |\phi_s| \sum_{s=k}^{l+k} |\phi_s| (d_l^{-3} + l^{-3}), & \text{if } |l-j+k| > A_0, \end{cases} \end{aligned}$$

as required in (7.33).

REFERENCES

Akonom, J. (1993). Comportement asymptotique du temps d'occupation du processus des sommes partielles. *Annales de l'IHP Probabilités et Statistiques*, 29, 57–81.

Billingsley, P. (1968). *Convergence of probability measures*. John Wiley & Sons.

Billingsley, P. (1974). Conditional distributions and tightness. *The Annals of Probability*, 2, 480–485.

Borodin, A. N., & Ibragimov, I. A. (1995). *Limit theorems for functionals of random walks*, 195. American Mathematical Society.

Cai, Z., Li, Q., & Park, J. Y. (2009). Functional-coefficient models for nonstationary time series data. *Journal of Econometrics*, 148, 101–113.

Chan, N., & Wang, Q. (2015). Nonlinear regressions with nonstationary time series. *Journal of Econometrics*, 185, 182–195.

Dong, C., & Linton, O. (2018). Additive nonparametric models with time variable and both stationary and nonstationary regressors. *Journal of Econometrics*, 207, 212–236.

Duffy, J. A. (2016). A uniform law for convergence to the local times of linear fractional stable motions. *Annals of Applied Probability*, 26, 45–72.

Duffy, J. A. (2020). Asymptotic theory for kernel estimators under moderate deviations from a unit root, with an application to the asymptotic size of nonparametric tests. *Econometric Theory*, 36, 559–582.

Duffy, J. A., & Kasparis, I. (2021). Estimation and inference in the presence of fractional $d = 1/2$ and weakly nonstationary processes. *Annals of Statistics*, 49, 1195–1217.

Fiorio, C. V., Hajivassiliou, V. A., & Phillips, P. C. B. (2010). Bimodal t-ratios: The impact of thick tails on inference. *The Econometrics Journal*, 13, 271–289.

Gao, J., & Phillips, P. C. B. (2013). Semiparametric estimation in triangular system equations with nonstationarity. *Journal of Econometrics*, 176, 59–79.

Geman, D., & Horowitz, J. (1980). Occupation densities. *The Annals of Probability*, 8, 1–67.

Gozalo, P., & Linton, O. (2000). Local nonlinear least squares: Using parametric information in nonparametric regression. *Journal of Econometrics*, 99, 63–106.

Hu, Z., Phillips, P. C. B., & Wang, Q. (2021). Nonlinear cointegrating power function regression with endogeneity. *Econometric Theory*, 37, 1173–1213.

Jeganathan, P. (2004). Convergence of functionals of sums of r.v.s to local times of fractional stable motions. *Annals of Probability*, 32, 1771–1795.

Jeganathan, P. (2008). *Limit theorems for functionals of sums that converge to fractional stable motions*. Cowles Foundation Discussion Paper No. 1649. Yale University.

Karlsen, H. A., & Tjøstheim, D. (2001). Nonparametric estimation in null recurrent time series. *Annals of Statistics*, 29, 372–416.

Li, D., Tjøstheim, D., & Gao, J. (2016). Estimation in nonlinear regression with Harris recurrent Markov chains. *Annals of Statistics*, 44, 1957–1987.

Liang, H.-Y., Shen, Y., & Wang, Q. (2023). Functional-coefficient cointegrating regression with endogeneity. In *Essays in Honor of Joon Y. Park: Econometric Theory* (pp. 157–186). Emerald Publishing Limited.

- Logan, B. F., Mallows, C. L., Rice, S., & Shepp, L. A. (1973). Limit distributions of self-normalized sums. *The Annals of Probability*, 1, 788–809.
- Park, J. Y. (2014). Nonstationary nonlinearity: A survey on Peter Phillips's contributions with a new perspective. *Econometric Theory*, 30, 894–922.
- Park, J. Y., & Phillips, P. C. B. (1999). Asymptotics for nonlinear transformations of integrated time series. *Econometric Theory*, 15, 269–298.
- Park, J. Y., & Phillips, P. C. B. (2000). Nonstationary binary choice. *Econometrica*, 68, 1249–1280.
- Park, J. Y., & Phillips, P. C. B. (2001). Nonlinear regressions with integrated time series. *Econometrica*, 69, 117–161.
- Petrov, V. V. (1995). *Limit theorems of probability theory; sequences of independent random variables*. Oxford Studies in Probability. Clarendon Press.
- Phillips, P. C. B. (2009). Local limit theory and spurious nonparametric regression. *Econometric Theory*, 25, 1466–1497.
- Phillips, P. C. B., & Park, J. Y. (1998). *Nonstationary density estimation and kernel autoregression*. Cowles Foundation Discussion Paper, No. 1181. Yale University.
- Phillips, P. C. B., & Wang, Y. (2023). When bias contributes to variance: True limit theory in functional coefficient cointegrating regression. *Journal of Econometrics*, 232, 469–489.
- Sun, Y., Cai, Z., & Li, Q. (2016). A consistent nonparametric test on semiparametric smooth coefficient models with integrated time series. *Econometric Theory*, 32, 988–1022.
- Sun, Y., & Li, Q. (2011). Data-driven bandwidth selection for nonstationary semiparametric models. *Journal of Business & Economic Statistics*, 29, 541–551.
- Tjøstheim, D. (2020). Some notes on nonlinear cointegration: A partial review with some novel perspectives. *Econometric Reviews*, 39, 655–673.
- Tu, Y., & Wang, Y. (2022). Spurious functional-coefficient regression models and robust inference with marginal integration. *Journal of Econometrics*, 229, 396–421.
- Wang, Q. (2015). *Limit theorems for nonlinear cointegrating regression* (Vol. 5). World Scientific.
- Wang, Q., Lin, Y.-X., & Gulati, C. M. (2003). Strong approximation for long memory processes with applications. *Journal of Theoretical Probability*, 16, 377–389.
- Wang, Q., & Phillips, P. C. B. (2009a). Asymptotic theory for local time density estimation and nonparametric cointegrating regression. *Econometric Theory*, 25, 710–738.
- Wang, Q., & Phillips, P. C. B. (2009b). Structural nonparametric cointegrating regression. *Econometrica*, 77, 1901–1948.
- Wang, Q., & Phillips, P. C. B. (2011). Asymptotic theory for zero energy functionals with nonparametric regression applications. *Econometric Theory*, 27, 235–259.
- Wang, Q., & Phillips, P. C. B. (2016). Nonparametric cointegrating regression with endogeneity and long memory. *Econometric Theory*, 32, 359–401.
- Wang, Q., Phillips, P. C. B., & Kasparis, I. (2021). Latent variable nonparametric cointegrating regression. *Econometric Theory*, 37, 138–168.
- Xiao, Z. (2009). Functional-coefficient cointegration models. *Journal of Econometrics*, 152, 81–92.