
INSTRUCTIONS TO CONTRIBUTORS

The *Journal of Helminthology* publishes papers on all aspects of animal parasitic helminths, particularly those of medical or veterinary importance, but only in exceptional circumstances will systematic or taxonomic studies be acceptable.

Manuscripts, which must be in English or French (with an English summary) should be accompanied by a letter signed by *all* the authors and should be addressed to:

The Editor, Journal of Helminthology,
CAB International Institute of Parasitology,
395A Hatfield Road,
St Albans, Herts AL4 0XQ.
England.

Three copies of a typescript, on size A4 paper with double spacing, should be submitted. Papers should be preceded by a short abstract and will normally have the following sections: brief Introduction; Materials and Methods; Results; Discussion; Acknowledgements; References. However, the form of the paper may vary, depending on its subject matter; recent past issues should be consulted for a suitable form. Research Notes should also be preceded by a brief abstract. Illustrations should be drawn in Indian ink, preferably not more than double the final size. Care should be taken that all illustrations fit into the format of the Journal. The maximum size an illustration will be printed is 12.0×20.0 cm. Where many separate drawings are made, some indication of how they may be grouped to make a corporate plate without undue wastage of space, should be indicated. Some indication of scale (preferably a scale bar) should normally be given on the figure. Photocopies of illustrations should be enclosed for refereeing purposes. Lettering and numbering, which must be of a high standard, should be added by the author, with due regard for subsequent reduction.

Photographs should be glossy prints of the same size as they are to appear in the Journal (maximum size 12.0×20.0 cm). Composite prints must be mounted and can have the separate photographs abutting; they will then have a separating line inserted by the printers. All figures and letters on photographs must be inserted by the author.

Information should not be repeated in the text and in tables or figures. The legends to tables and to figures should be sufficiently detailed for the information to be understood without reference to the text.

References should be given in alphabetical order with the full title of the journal. The following are examples:

DUKE, B. O. L. (1971) The ecology of onchocerciasis in man and animals. In: *Ecology and Physiology of Parasites* (editor, A. M. Fallis) pp. 213–222. Adam Hilger Ltd.: London.

JAMES, C. & WEBBE, G. (1973) A comparison of Egyptian and East African strains of *Schistosoma haematobium*. *Journal of Helminthology*, **47**, 49–59.

25 offprints are provided free of charge; additional copies may be ordered at the proof stage.

Contents

	Pages
The capability of the predacious fungus <i>Arthrobotrys oligospora</i> (Hyphomycetales) to reduce numbers of infective larvae of <i>Ostertagia ostertagi</i> (Trichostrongylidae) in cow pats and herbage during the grazing season in Denmark. J. GRØNVOLD, P. NANSEN, S. A. HENRIKSEN, J. THYLIN and J. WOLSTRUP	271-280
The effect of constant and changing temperatures on the development of the eggs and larvae of <i>Oswaldocruzia filiformis</i> (Nematoda: Trichostrongyloidea). C. T. GRIFFIN	281-292
<i>In vitro</i> transformation of <i>Diplostomum spathaceum</i> (Digenea) cercariae and short term maintenance of post-penetration larvae <i>in vitro</i> . S. K. WHYTE, L. H. CHAPPELL and C. J. SECOMBES	293-302
On the viability of <i>Fasciola gigantica</i> metacercariae ingested by <i>Lymnaea auricularia</i> . S. C. YADAV and S. C. GUPTA (Research Note)	303-304
The tegument in Hemiuridae (Digenea: Hemiuroidea): structure and function in the adult. B. F. MATTHEWS and R. A. MATTHEWS	305-316
The ecsoma in Hemiuridae (Digenea: Hemiuroidea): tegumental structure and function in the mesocercaria and the metacercaria of <i>Lecithochirium furcolum</i> (Jones, 1933) Dawes, 1947. B. F. MATTHEWS and R. A. MATTHEWS	317-330
Ultrastructure and cytochemistry of the tegument of <i>Orthocoelium scolicoelium</i> and <i>Paramphistomum cervi</i> (Trematoda: Digenea). P. N. SHARMA and R. E. B. HANNA	331-343
Morphological differences between Venezuelan and African microfilariae of <i>Onchocerca volvulus</i> . C. BOTTO, A. ESCALANTE, M. ARANGO and L. YARZABAL	345-351
Announcements	344
Subject Index	353-356
Author Index	357-359
Contents of Volume	i-iv