

2. Department of Traumatology and Acute Critical Medicine, Osaka, University Medical School, Osaka, Japan
3. Department of Emergency Medicine, Osaka Prefectural General Hospital, Osaka, Japan

A total of 14 patients with crush syndrome were transferred to Osaka University Hospital after the 1995 Hanshin-Awaji earthquake. Clinical aspects of three cases representing the typical and atypical crush syndrome were presented.

**[Case 1]** A 66-year-old female was buried under a collapsed house and was rescued one hour after entrapment. She was conscious on the scene and was evaluated as yellow (severe, but not critical). However, she developed cardiopulmonary arrest during transport to the hospital. She was successfully resuscitated, and was referred to this Center. She had unstable pelvic fractures as well as crush injury to the buttock. Bleeding from the pelvic fracture was the cause of her sudden collapse. Transcatheter embolization controlled the bleeding and external fixation of the pelvis was provided. She did not develop renal failure or other organ dysfunctions. In this patient, the crush injury was not severe, but the associated pelvic fracture and hemorrhage caused critical situation.

**[Case 2]** A 25-year-old female was buried under the debris for three hours and was transferred to this hospital three hours following her extrication. On admission, severe hyperkalemia, severe rhabdomyolysis, and increased intracompartmental pressure were detected. Emergency hemodialysis and fasciotomy were performed. Aggressive fluid supplementation reaching 24 liters for the initial 24 hours was needed to counteract massive edema and fluid loss to the injured legs. This fluid support was sustained until the fasciotomy wound was closed. The clinical course was not complicated by acute renal failure, and now she can walk without assistance.

**[Case 3]** A 25-year-old male was transferred to this hospital 35 hours after injury. He had been compressed under the debris for eight hours, and had renal failure on admission. He was hemodialyzed from the day of arrival. After several days, he had unexpected complications including necrosis of the gall bladder that required surgical intervention. His general condition and renal function recovered rapidly after the operation.

**Keywords:** acute renal failure; cardiopulmonary arrest; crush syndrome; earthquake; fasciotomy; fracture; hemodialysis; hyperkalemia; rhabdomyolysis

*Prehosp Disast Med 2002;17:s25-26.*

### Long-Term Physical Outcome of Patients Who Suffered Crush Syndrome: Prognostic Indicators

Tetsuya Matsuoka

Senshu Critical Care Medical Center, Japan

**Objective:** To identify independent predictors of physical outcome in patients suffering from crush syndrome

**Methods:** Sensory and motor functions were examined two years after the 1995 Hanshin-Awaji earthquake in 42

patients with a combined total of 58 compressed lower extremities. The influences of time to rescue, fasciotomy, and radical debridement on lower leg muscle strength were evaluated by stepwise regression analysis.

**Results:** Severe disabilities related to the lower leg compartment were present, and the anterior compartment was damaged more severely than was the posterior compartment. Stepwise regression analysis showed the performance of fasciotomy and debridement to be an independent predictor of long-term, lower leg muscle survival, and indicated that it was an independent predictor when the debrided compartments were not included in the analysis. In all debrided anterior compartments, muscle contractility was abolished completely. There was a significant negative correlation between time to fasciotomy and lower leg muscle strength.

**Conclusions:** Secondary compartment syndrome affected physical outcome in crush syndrome patients. There was no evidence that fasciotomy improves functional outcome. Delayed rescue, delayed fasciotomy, and radical debridement may worsen the physical prognosis. Indications for fasciotomy in crush syndrome during the acute phase need further study and evaluation.

**Keywords:** debridement; earthquake; fasciotomy; Hanshin-Awaji; morbidity; predictors; rescue

*Prehosp Disast Med 2002;17:s26.*

### Penetrating Cardiac Injury following Sternal Fracture

Francis Fu Sheng Wu,<sup>1</sup> Chi-Wen Juan,<sup>1</sup> Fang-Chu Chen,<sup>1</sup> Sai-Kit Ho,<sup>1</sup> Yao-Ru Hu,<sup>2</sup> Tzu-Hui Tung<sup>2</sup>

1. Department of Emergency Medicine, Changhua Christian Hospital, Changhua, Taiwan, ROC
2. Department of Nursing, Changhua Christian Hospital, Changhua, Taiwan, ROC

**Objective:** Myocardial injury used to be considered as one of the major complications associated with sternal fracture, even though recent studies on injuries associated with fracture of sternum are contrary to this belief. Many authors now believe the presence of sternal fracture no longer is indicative of occult injuries to the underlying structure such as the heart.

**Methods:** A case was presented of a 38-year-old female patient transferred to our hospital after being injured in a motor vehicle accident. On arrival, her blood pressure (BP) was 90/50 mmHg but two hours later, it dropped to 60/30 mmHg. Although her chest roentgenography and electrocardiography (ECG) did not reveal any significant findings, chest computerized tomography (CT) scan later revealed a sternal fracture and cardiac tamponade.

**Results:** A diagnosis of cardiac rupture resulting from sternal fracture following blunt chest trauma was made. Under midline sternotomy, her right atrial rupture was repaired. The patient was doing well during a three months post-operative follow-up.

**Conclusion:** Clinicians should maintain a high index of suspicion for the presence cardiac tamponade in cases presented as blunt chest trauma as early diagnosis and surgical intervention is vital to the patient's survival.

**Keywords:** cardiac tamponade; right atrial rupture; sternal fracture.  
*Prehosp Disast Med 2002;17:s26-27.*

### The Effects on the Central Nervous System Induced by Tetrodotoxin Poisoning

Francis Fu Sheng Wu;<sup>1</sup> Chi-Wen Juan;<sup>1</sup> Pang-Chi Lai;<sup>2</sup>  
Kuo-Chou Cheng;<sup>1</sup> Chia-Ju Wu;<sup>2</sup> Hsiu-Hui Hou<sup>2</sup>

1. Department of Emergency Medicine, Changhua Christian Hospital, Changua, Taiwan, ROC
2. Department of Nursing, Changhua Christian Hospital, Changua, Taiwan, ROC

**Objective:** Tetrodotoxin (TTX) is a neurotoxin known to cause food poisoning. It is found in a variety of both freshwater and marine species. Patients with tetrodotoxin intoxication generally have a very typical course of illness involving the cardiovascular, respiratory, peripheral and central nervous systems.

**Methods:** A 59-year-old, male, taxi driver presented to the emergency department with dyspnea and numbness after eating some porridge cooked with Takifugu niphobles. On arrival, he was tachypneic, hypertensive, consciousness clear, and was able to give a coherent history. He immediately was intubated, but respiratory effort ceased several minutes later. About one hour later, he was in deep coma with Glasgow Coma Scale (GCS) Score = 3, had a sinus bradycardia, and became severely hypotensive.

**Results:** During the next two days, he continued to have fluctuating blood pressure, hypothermia, and polyuria. At 36 hours after admission, he regained consciousness and some upper limb motor function, but still had no spontaneous breathing effort. An EEG showed "minor diffuse cortical dysfunction likely to be metabolic in origin". Later in the day his vision recovered spontaneously. Ventilatory function returned and the patient self-extubated on the following day. On day 10, he was discharged after a follow-up EEG that showed "no evidence of focal cortical dysfunction". Tetrodotoxin measured by bioassay was 300 mouse units per gram (MU/g).

**Conclusion:** To our knowledge this is the first report of a human that demonstrated this cortical dysfunction on EEG during tetrodotoxin poisoning.

**Keywords:** central nervous system; intoxication; takifugu niphobles; tetrodotoxin  
*Prehosp Disast Med 2002;117:s27.*

### Six Years Experience with Telemedicine in an Emergency Department in Taiwan

Lee-Min Wang, MD;<sup>1</sup> Hong-Chang Lo, MD;<sup>1</sup>  
Chen-Hsen Lee, MD;<sup>1</sup> Dab-Dian Tang;<sup>2</sup> Pi-Chin Yu;<sup>3</sup>  
Low-Tone Ho, MD<sup>3</sup>

1. Emergency Department
2. Information Services Center
3. Department of Medical Research and Education, Veterans General Hospital-Taipei; National Defense Medical Center. National Yang-Ming University, Taipei, Taiwan ROC

**Objective:** To observe telemedicine in an emergency

department between Taipei-Veterans General Hospital and Kinmen-Granite, County Hospitals in Taiwan.

**Methods:** This clinical study was carried out from July 1996 through to September 2001. Data for total of 920 teleconsultation cases were collected using a questionnaire. Evaluation elements included: 1) consulting quality; 2) duration of the teleconsultation; the benefit to the patients; and 4) the charges. The purposes of the teleconsultation project were to establish electronically an exchange of medical information and to conduct clinical examinations and consultations.

**Results:** The peak interval of the ages of the patients was 21–30 years. Radiological consultation was the most frequently involved area of consultation, followed by chest medicine, neurosurgery, orthopedics, dermatology, and general surgery. The questionnaire indicated that 74% of the time, teleconsultations was used to obtain a the second opinion. More than 95% of the physicians involved believe that the telemedicine system will affect the quality of the consultation and that the consultation is helpful. The doctors in Taipei-VGH were more satisfied with the teleconsultation facility than were the doctors in Kinmen Hospitals.

**Conclusion:** The consultation doctors confirm that the clinical evaluation using the telemedicine is positive and has the potential to help during a disaster period.

**Keywords:** consultation; questionnaire; telecommunication, telemedicine.  
*Prehosp Disast Med 2002;17:s27.*

### Covering Disasters: A Look at Disaster News Reporting in the Philippine Media

Lirio Racho Luyun

Cemshad, Inc., The Philippines

**Objectives:** The study aims to find out how disaster news are framed in the Philippine print media and the media practices that inform the framing of such news.

**Method:** The study qualitatively content analyzed several disaster news reports from different broadsheets within 2001.

**Results:** The regularity of events that result in disasters that strike the Philippines for the past years reveals a pattern that the press applies in framing disaster news. Emphasis is given to statistics regarding the number of persons affected by these events. It also showed a tendency to place emphasis on personalities, spending precious ink to highlight the activities of politicians and officials who often use such disasters as photo opportunities to further their political careers. Disaster news, on the whole, is presented as an 'event', given prominence for a while until a new event takes its place in the public spotlight.

**Conclusions:** The tendency of the press to treat disaster news as 'events' has the potential to instill a feeling of fatigue among the populace towards disasters. Instead of communicating information that will help people make critical and informed choices during such disasters, the press focus on simple cause and effect explanations as to why such disasters happen. This has not translated into