

Cambridge Core

The new
home of
academic
content

cambridge.org/core

Cambridge Core



CAMBRIDGE
UNIVERSITY PRESS



Life Sciences

Books and Journals from
Cambridge University Press

Cambridge is one of the leading publishers in ecology and conservation biology and publishes high quality texts and research across the breadth of the life sciences, focusing particularly on animal behaviour, biological anthropology, evolutionary biology, computational and systems biology, as well as statistics and professional development titles for biologists.

We also have an extensive portfolio of established journals in agriculture, ecology and conservation, and animal science.

For further details visit:

cambridge.org/core-life-sciences

**Cambridge
Core**



**CAMBRIDGE
UNIVERSITY PRESS**

THE MARINE BIOLOGICAL ASSOCIATION OF THE UNITED KINGDOM

TERMS OF MEMBERSHIP FOR 2019

MBA MEMBERSHIP (prices per annum)	Price from 1st March 2019
Young Marine Biologist (YMB)	£12.50
Student Member	£25.00
Associate Member	£40.00 (direct debit discount £5)
Professional Member, Mem.MBA	£45.00 (direct debit discount £5)
MBA Fellow, FMBA	£120.00 (direct debit discount £5)
Institutional Member	£340.00

JOURNAL SUBSCRIPTION (MBA price added to the above membership fees, with the exception of YMB as no Journal option for this category)	
Online Journal	£20.00
Print & Online Journal	£80.00

The Association is a learned society founded in 1884 and incorporated by Royal Charter in 2013. The Marine Biological Association (MBA) provides a unified, clear, independent voice on behalf of the marine biological community. In 1888, the MBA opened the Plymouth laboratory at Citadel Hill.

The MBA is also a registered charity with its key aims being to promote scientific research into all aspects of life in the sea, including the environment on which it depends, and to disseminate to the public the knowledge gained. The Patron of the Association is His Royal Highness the Duke of Edinburgh, who is also an Honorary Fellow. The other Honorary Fellows are His Serene Highness Prince Albert of Monaco, Dr Sylvia Earle, Sir Tim Hunt and Professor James Lovelock.

The work of the Association is controlled by the Council elected annually by its subscribing members.

The Marine Biological Association, under the direction of its Council, has undertaken research in all branches of marine science and many of the results have been published in this Journal. An account of the history of the first hundred years of the Association and Journal may be found in Volume 67 (pp. 463 and 465).

Since its foundation the Association has been supported by grants from Research Councils and by subscriptions and donations

from private members, universities, learned societies, the Fishmongers' Company and other public bodies.

The Marine Biological Association is a very active scientific learned society with a large international membership. The MBA also manages an internationally renowned research programme.

Subscriptions can be paid by cash/cheque, direct debit from a UK bank account or credit card. *Subscriptions are reduced by £5 per annum if paying by direct debit.

Members of the Association have a wide range of rights and privileges including the use of post-nominals (FMBA for MBA Fellows, Mem.MBA for Professional Members) and being entitled to receive The Marine Biologist Magazine. The Commissioners of Inland Revenue have approved the Association for the purposes of Section 16, Finance Act, 1958, and that the whole of the annual subscriptions paid by a member who qualifies for relief under the section will be allowable as a deduction from his emoluments assessable to income tax.

All correspondence should be addressed to: The Membership Secretary, Marine Biological Association, The Laboratory, Citadel Hill, Plymouth, PL1 2PB, UK (Tel. +44 (0) 1752 426493; e-mail: membership@mba.ac.uk).

PUBLISHED BY THE PRESS SYNDICATE OF THE UNIVERSITY OF CAMBRIDGE
The Pitt Building, Trumpington Street, Cambridge, CB2 1RP, United Kingdom

This journal issue has been printed on FSC-certified paper and cover board. FSC is an independent, non-governmental, not-for-profit organization established to promote the responsible management of the world's forests. Please see www.fsc.org for information.

Printed by Bell and Bain Limited, Glasgow

CAMBRIDGE UNIVERSITY PRESS

Journals Fulfillment Department, UPH, Shaftesbury Road, Cambridge CB2 8BS, United Kingdom
1 Liberty Plaza, Floor 20, New York, NY 10006, USA
477 Williamstown Road, Port Melbourne, VIC 3207, Australia
C/ Orense, 4, planta 13, 28020, Madrid, Spain
Dock House, The Waterfront, Cape Town 8001, South Africa

CONTENTS

ARTICLES

Talbot, E., Bruggeman, J., Hauton, C. and Widdicombe, S. Uncovering the environmental drivers of short-term temporal dynamics in an epibenthic community from the Western English Channel	1467
Kim, H.-G., Hawkins, L.E., Godbold, J.A., Maclean, M., Oh, C.-W., Rho, H.S. and Hawkins, S.J. A comparison of epiphytic nematode diversity and assemblages in <i>Corallina</i> turves on British and South Korean coasts across hierarchical spatial scales	1481
Alvarado-Rodríguez, J.F., Nava, H. and Carballo, J.L. Spatio-temporal variation in rate of carbonate deposition by encrusting organisms in different reef microhabitats from Eastern Pacific coral reefs	1495
Quimpo, T.J.R., Cabaitan, P.C., Go, K.T.B., Dumalagan Jr, E.E., Villanoy, C.L. and Siringan, F.P. Similarity in benthic habitat and fish assemblages in the upper mesophotic and shallow water reefs in the West Philippine Sea	1507
Costa, G., Bavestrello, G., Micaroni, V., Pansini, M., Strano, F. and Bertolino, M. Sponge community variation along the Apulian coasts (Otranto Strait) over a pluri-decennial time span. Does water warming drive a sponge diversity increasing in the Mediterranean Sea?	1519
Laverick, J.H., Green, T.K., Burdett, H.L., Newton, J. and Rogers, A.D. Depth alone is an inappropriate proxy for physiological change in the mesophotic coral <i>Agaricia lamarcki</i>	1535
Hernández-Alcántara, P., Cruz-Pérez, I.N. and Solís-Weiss, V. Composition and diversity patterns of Eunicida and Amphinomida (Annelida) associated to dead coral in the Chinchorro Bank Biosphere Reserve, Caribbean Sea	1547
Méndez, N., Hilliard, J. and Schulze, A. Early development of two <i>Capitella</i> species (Annelida: Capitellidae) from the Gulf of Mexico	1557
Righi, S., Maletti, I., Maltagliati, F., Castelli, A., Barbieri, M., Fai, S., Prevedelli, D. and Simonini, R. Morphometric and molecular characterization of an expanding Ionian population of the fireworm <i>Hermodice carunculata</i> (Annelida)	1569
Purroy, A., Bukša, F., Puljas, S. and Peharda, M. Variations in the reproductive investment of a venerid bivalve, <i>Callista chione</i>	1579
Zhao, D., Kong, L.-F., Sasaki, T., and Li, Q. Shell variations in the gastropod, <i>Monodonta labio</i> , in the North-western Pacific: the important role of temperature in the evolution process	1591
Ruiz, J.M., Carro, B., Albaina, N., Couceiro, L., Quintela, M. and Barreiro, R. A decadal study of biometric and imposex indices in two gastropods	1601
Sousa, R., Henriques, P., Vasconcelos, J., Faria, G., Riera, R., Pinto, A.R., Delgado, J. and Hawkins, S.J. First observations of hermaphroditism in the patellid limpet <i>Patella piperata</i> Gould, 1846	1615
Sajikumar, K.K., Sasikumar, G., Mohan, G., Kripa, V., Alloyicious, P.S. and Mohamed, K.S. Age and growth of the little Indian squid, <i>Loliolus hardwickei</i> (Gray, 1849) in the Arabian Sea	1621
Bosley, K.M., Coleman, N. and Dumbauld, B.R. Verification of ageing methods for the burrowing shrimp, <i>Neotrypaea californiensis</i> , using extractable lipofuscin and gastric mill cuticular features	1627
Canales, T.M., Wiff, R., Quiroz, J.C. and Queirolo, D. Estimation of natural mortality in two demersal squat lobster species off Chile	1639
Şensurat-Genç, T., Akyol, O., Özgül, A. and Özden, U. Food composition of whiting <i>Merlangius merlangus</i> , captured around the sea-cage fish farms in Ordu, south-eastern Black Sea	1651
Ehemann, N.R., Abitia-Cardenas, L.A., Navia, A.F., Mejía-Falla, P.A. and Cruz-Escalona, V.H. Zeros as a result in diet studies, is this really bad? <i>Rhinoptera steindachneri</i> as a case study	1661
Condini, M.V., Pereyra, P.E.R., Garcia, A.M., Saint'pierre, T.D., Ceni, G., Lugo, R., Fontoura, N.F., Vieira, J.P. and Albuquerque, C.Q. Use of fresh water by an estuarine-resident marine catfish: evidence from gonadal and otolith chemistry analyses	1667
de Carvalho, B.M., Spach, H.L., Vaz-Dos-Santos, A.M. and Volpedo, A.V. Otolith shape index: is it a tool for trophic ecology studies?	1675
Orlova, S.Y., Schepetov, D.M., Mogue, N.S., Smirnova, M.A., Senou, H., Baitaliuk, A.A. and Orlov, A.M. Evolutionary history told by mitochondrial markers of large teleost deep-sea predators of family Anoplopomatidae Jordan & Gilbert 1883, endemic to the North Pacific	1683
CORRIGENDUM	
Nekliudova, U.A., Shunkina, K.V., Grishankov, A.V., Varfolomeeva, M.A., Granovitch, A.I. and Ostrovsky, A.N. Colonies as dynamic systems: reconstructing the life history of <i>Cribrilina annulata</i> (Bryozoa) on two algal substrates – CORRIGENDUM	1693