

Research Article

Cite this article: Mohapatra SR, Das SK, Kar MR, Das H (2024). Floral biology and breeding behaviour of *Melia azedarach* L.: imperative for hybridization. *Plant Genetic Resources: Characterization and Utilization* 1–10. <https://doi.org/10.1017/S1479262124000376>

Received: 25 April 2024

Revised: 6 July 2024

Accepted: 8 July 2024


Keywords:

breeding system; hybridization; maha-neem; *Melia azedarach* L.; reproductive biology

Corresponding author:

Sourav Ranjan Mohapatra;
Email: srmohapatra.forestry@ouat.ac.in

Floral biology and breeding behaviour of *Melia azedarach* L.: imperative for hybridization

Sourav Ranjan Mohapatra¹ , Susanta Kumar Das², Manas Ranjan Kar² and Harekrishna Das²

¹Department of Forest Biology and Tree Improvement, Odisha University of Agriculture and Technology, Bhubaneswar- 751 003, Odisha, India and ²Department of Forest Resource Management, Odisha University of Agriculture and Technology, Bhubaneswar- 751 003, Odisha, India

Abstract

Melia azedarach L. commonly called Maha Neem is an economically and industrially important tree species with global significance. Although species possess versatile importance worldwide, information on reproductive biology and breeding system is scarce and limited for eastern coastal plain of India. Therefore, current study provides a detailed report on reproductive biology of *M. azedarach*. Maha Neem was found to bear violet to whitish violet, slightly fragrant, 14.99 ± 0.05 mm. long and 17.01 ± 0.08 mm. wide flowers. This tree commenced opening of floral buds during March and continued until May with a peak during March-end. The anthesis of species peaked between 08:00 and 10:00 am, which coincides with insect activity. Anthers were observed to dehisce during or shortly before anthesis; however, stigma started receptivity before anthesis and continued invitation up to 12 h after anthesis. Pollen viability ranged from $96.67 \pm 1.6\%$ to 98.26 ± 1.2 at the time of anthesis; after that decreased rapidly. Pollen: ovule ratio of 1096.38 ± 108.70 indicated the possibility of autogamy. The breeding system of *M. azedarach* revealed that fruit sets under natural pollination (NP) were significantly higher than Xenogamy. However, substantial difference was not reported in fruit set percentage under natural pollination and Autogamy, which is strong evidence in favour of self-pollination. Moreover, Maha-Neem is entomophilous with frequent visits by Apis and Syrphid flies. Current findings will be helpful in designing potent conservation strategies and planning successful breeding programmes.

Introduction

The ever-increasing population and rapid decline in quality stock are putting pressure on forest stands and causing climatic disturbances. In 2018, India imported forest products, logs and wood products (other than logs), amounting to USD 2073 million, USD 1052 million, and USD 1021 million, respectively (Global Agriculture Information Network, 2019). By 2025, the gap between demand and supply for round wood will be approximately 22.4 million m³ (Srinivasan *et al.*, 2018). Demand is expected to rise further as technology advances and trade, and tariffs become more convenient. To fill this void, short rotation and fast-growing species must be the focus (Khan and Chaudhry, 2007). Furthermore, such species may be an essential keystone for conservation and climate change. Hence, there is a necessity to adopt species like *Melia azedarach* L., which possess highly desirable features like high productivity, greater biomass, short rotation, wider adaptability and faster-growing habit.

Melia azedarach L. commonly known as Persian lilac or China tree (English), Bakain (Hindi), Maha Neem (Odia), belongs to the prime family Meliaceae is a fairly large, deciduous tree with a height of 7–12 m. but exceptionally can grow up to 45 m. (Duong *et al.*, 2017), and cylindrical straight bole up to 9 m. (Troup, 1921). Being a light-demanding species, it grows vigorously in wide forest types and can thrive in various edaphic and environmental conditions. Further, tree requires least cultural efforts and low maintenance costs, making them more adaptable to small and marginal farmers. This species has been acknowledged as a source for plywood industries (Rahman *et al.*, 2014) and recognized suitable for pulp, paper, furniture, agricultural tools, home construction and packaging case industries (Venson *et al.*, 2008; Duong *et al.*, 2017). Additionally, it has been revealed that species contains analgesic, antibacterial, antifungal, antifeedant, antioxidant, anti-inflammatory, anti-diabetic and hepatoprotective properties (Nivedita *et al.*, 2019).

Reproductive knowledge of tree species is crucial for comprehending pollination and breeding systems (Tandon *et al.*, 2003), managing and recovering threatened species (Kuniyal *et al.*, 2003; Murugan *et al.*, 2006), and improving desired traits and developing new varieties (Khanduri *et al.*, 2016). Details related to floral biology permit learning about population viability and its gene flow, which is necessary for wild species conservation (Marten and Quesada, 2001; Lee *et al.*, 2006). Similarly, the successfulness of reproductive cycle depends upon



detection of phenology, intensity and duration of flowering, which may impact the population of pollinators and frugivores relying on plant species (Newstrom *et al.*, 1994). Apart from this, reproductive biology is a prerequisite for developing breeding strategies. Henceforth thorough information on reproductive biology and plant breeding systems is crucial for creating an efficient approach to developing successful breeding and conservation strategies. However, Persian lilac has not been explored for varying research dimensions, including phenology, floral architecture and breeding system. Moreover, hybridization-related techniques, including phenology, pollen storage, viability and cross-ability assessment, are minutely uncovered and have minimal information available in public domain with special reference to eastern coastal plains of India. Therefore, it needs systematic investigation and multifaceted research activities to uncover crucial information. Keeping the preceding in mind, a detailed study on reproductive biology of *M. azedarach* has been undertaken. On that account, the present investigation on *M. azedarach* intended to elucidate (a) floral morphology, structure, period of anthesis and stigma receptivity, (b) pollen studies and (c) pollination and breeding system. The outcome of the study aid in filling the knowledge gaps and unravelling crucial facts, which will assist in planning and developing strategies for effective and efficient conservation. This would also pave the way to a sustained generation of raw material for commercial usage in future industries and regeneration for habitat enrichment.

Materials and methods

Study location

The current investigation was undertaken in and around Odisha University of Agriculture and Technology, Bhubaneswar. The area is situated between 20° 15' 52" N latitude and 85° 48' 43" E longitude and receives an average annual rainfall of 1550.2 mm. During summer, the mean monthly maximum and minimum temperature ranged from 45.6 to 34.5°C, respectively, while corresponding winter temperatures varied from 28.3°C maximum and 16.7°C minimum. Five trees of *M. azedarach*, were selected for their accessibility, convenience and representing normal growth conditions free from biotic disturbances. These trees, aged approximately 7 to 8 years old exhibited height ranging from 10 to 15 metres and diameters from 22 to 43 cm. They were studied continuously for two years (2021 and 2022) with a special focus during flowering season.

Phenology

Phonological events of initiation of flowering primordia, bud break, flowering duration, peak flowering period, fruit set, and maturity were recorded from five labelled trees. Floral observations were performed every morning throughout the flowering season, while fruit features were observed once a week starting from mid-February 2022 and continuing throughout the current and subsequent year till the ripening of fruits. In addition, flower colour was observed by photographing flowers of the flowering plant using a good quality digital SLR (Nikon D300).

Floral architecture

The qualitative and quantitative pattern of floral morphology was characterized by inspecting 150 flowers in 15 randomly selected

inflorescences (10 flowers per inflorescence) from five different plants. Branches were tagged prior to the formation of floral buds, and observations were made regularly. Morphological characteristics including inflorescence type, flower colour, number of sepals, and petals per flower were inspected thoroughly. The structure and dimensions of floral components such as sepal, petal, stamen and pistil were observed and recorded under a microscope. In contrast, the number of floral buds per inflorescence was determined with simple counting.

Breeding system

The following characteristics were investigated to understand the nature of breeding system operative in this species:

Anthesis and anther dehiscence

Anthesis was observed in 20 randomly selected inflorescences (04 from each sampled tree) during the peak season of flower opening. Flowers were inspected at various time intervals throughout the day to record number of flowers opening in each time interval, *i.e.*, 06:00–08:00, 08:00–10:00, 10:00–12:00, 12:00–14:00, and 14:00–16:00. To minimize duplication and overcounting, opened flowers were marked with permanent markers at 2-hour intervals, and that same flower was used for recording the time of anther dehiscence.

Pollen production

The noon loop method was applied to assess pollen production. Anthers were collected from mature floral bud shortly before the anther dehiscence and were crushed to liberate total pollens in 0.5 ml of distilled water along with a drop of Tween-20. The solution was made homogenized, and pollen count was carried with Haemocytometer (Waller *et al.*, 1998). The average pollen production (P_a) per anther was calculated by applying following formula:

$$P_a = \left(\sum T_p / N \right) \times n$$

Where T_p = Total number of pollens in five drops; N = Total number of counting samples (slides); and n = Total number of drops.

Pollen grains per flower were computed by multiplying the number of pollen grains present per anther with number of anthers per flower. Similarly, pollen production within a tree (P_t) was estimated by multiplying number of inflorescences/trees (I_t), number of flowers per inflorescence (F_f), number of anthers/flowers (A_f), and number of pollen grains per anther (P_a) together.

$$P_t = \sum I_t F_f A_f P_a$$

Pollen viability and germination

Pollen viability was examined using *in vitro* aceto-carmin staining. Pollen grains were properly mixed with acetocarmine solution and allowed to stain for 10 min. Deeply stained, normal-looking pollen grains were tallied as viable, whereas faintly stained, empty, shriveled grains were recorded as non-viable.

To study the pollen germination, pollens were collected from freshly opened flowers of sampled trees and were immediately tested in customized media such as indole 3 acetic acid (IAA),

indole 3 butyric acid (IBA), and naphthalene acetic acid (NAA) at concentration of 50, 100 and 150 $\mu\text{l l}^{-1}$ each. After being placed in a medium, pollen grains from each solution were kept in an incubator and tested for germination at 2, 4, 6, and 8 h. The pollen germination was observed using a microscope (Brewbaker and Kwack, 1963).

Pollen: ovule ratio

Pollen-ovule ratio was calculated using Cruden's (1977) method, which is as follows:

$$P/O : \frac{\text{Pollen count per anther} \times \text{No. of anthers}}{\text{Number of ovules per flower}}$$

Stigma receptivity

Stigma receptivity was analysed with H_2O_2 following the method of Dafni 1992. Bubbling in the presence of hydrogen peroxide is considered a positive result (Osborn *et al.*, 1988).

Floral visitors

On each sampled tree, flowers were observed for pollinator visits between 06:00 and 18:00. The pollinators' foraging habits, number of visits and relative abundance were recorded. During the foraging time, i.e., between 06:00–18:00, insects were trapped using insect capturing nets (sweep nets) and polythene bags. Dried specimens of unknown insects were put on distinct rectangular papers for further identification, while the double mounting technique was employed for little insects.

Breeding behaviour

Determination of breeding behaviour was established with the following pollination treatments: (1) Natural pollination (NP): Buds were selected, counted, tagged and left as such for natural pollination; (2) Self-pollination (SP)/Autogamy: Buds in an inflorescence were selected, and covered with butter paper bag without emasculation prior to anthesis; (3) Open pollination (OP)/

Xenogamy: Mature buds were emasculated, tagged and left as such for pollination. Concurrently, surrounding buds and flowers were removed to reduce the possibility of geitonogamy. The Fruit set percentage was calculated by counting number of fruits set in each pollination treatment.

Data analysis

The SPSS version 29 (SPSS, Chicago) statistical package was used for data analysis. The standard error of the mean was calculated for each floral architectural trait and presented as \pm SE ($N=150$). As the difference in flower production among studied years was insignificant and statistical at par the data were pooled and presented. The result of the fruit set percentage was transformed (Sokal and Rohlf, 1995) and then ANOVA was performed to compare the fruit set percentage between various treatments at a 5% probability level.

Results

Phenology

The observations on flowering phenology indicated that *M. azedarach* formed floral buds as small protruding structures with the commencement of new leaves in the last week of February. The Proportion of floral buds increased gradually in number and reached a peak during the first week of March. Floral buds were purplish-blue and commenced their opening during mid-March and continued until May with a peak in late March (Table 1). The trees remained in bloom for 14 to 18 days, with an average of 15 days. Fruiting commenced in the first week of April and matured during November, where it was utterly ripened from December to January. Hence, completion of a single reproductive cycle takes an average duration of 12–13 months (Table 1). In addition, it was observed that abundant flowering and fruiting occur in both the studied years, consequently, seeds were available each year profusely.

Table 1. Phenological events of *M. azedarach* L

Tree No. (Location)	Character									
	Panicle initiation		Commencement of flowering		Peak period of flowering		Initiation of fruiting		Fruit ripening	
	I	II	I	II	Year		I	II	I	II
1 (20°26'51"E) (85°80'80"N)	Feb 27	Feb 28	Mar 14	Mar 16	Mar 26	Mar 29	Apr 3	Apr 2	Dec 29	Dec 28
2 (20°26'28"E) (85°81'19"N)	Feb 25	Feb 25	Mar 12	Mar 13	Mar 26	Apr 01	Apr 1	Apr 4	Dec 31	Dec 24
3 (20°20'47"E) (85°50'61"N)	Feb 26	Feb 28	Mar 15	Mar 14	Mar 29	Apr 2	Apr 1	Apr 5	Dec 27	Jan 5
4 (20°23'77"E) (85°72'15"N)	Feb 23	Feb 26	Mar 15	Mar 17	Mar 28	Mar 27	Apr 3	Apr 2	Jan 3	Dec 22
5 (20°24'54"E) (85°63'12"N)	Feb 24	Feb 23	Mar 12	Mar 18	Mar 26	Mar 29	Apr 4	Apr 6	Dec 30	Dec 27
Range	Feb 23 – Feb 28		Mar 13 – Mar 18		Mar 26 – Apr 02		Apr 1 – Apr 5		Dec 27- Jan 05	

Notes: I: Period of 2021–22; II: Period of 2022–23

Floral architecture

Maha Neem was found to bear deep violet to whitish violet, slightly fragrant, 14.99 ± 0.05 mm long and 17.01 ± 0.08 mm wide flowers arranged in a cymose inflorescence, which were recognized as a hermaphrodite, actinomorphic and pedicelled. Calyx of the flower was discovered to be 5–7 lobed, 1.70 ± 0.05 mm long, 1.00 ± 0.05 mm wide, ovate to oblong in shape, and tomentose (Fig. 1A, Table 2). The corolla appeared to have 4 to 6 petals, each measuring 14.53 ± 0.18 mm in length and 3.52 ± 0.04 mm in breadth (Fig. 1B, Table 2). Petals were found to be linear, spatulate, concave, pubescent on the exterior, and puberulous inside. Stamens were featured as ten dentate with a staminal tube rough textured, ribbed, basically gibbous and apically dilated or wider by bearing exerted and pubescent anthers. The most striking feature was alteration in staminal tube colour with age. The staminal tube of Maha Neem was pale purple during anthesis and turned moderate purple and finally dark before withering (Fig. 2). Likewise, the pistil had 3–8 locules (Fig. 1H), a superior and rectangular ovary, a cylindrical or slightly tapering style, and five serrated capitate stigmas. The stamen length was measured between 6.97 and 7.39 mm, with a mean value of 7.16 ± 0.05 mm, while width ranged from 2.66 to 2.91 mm, with a mean value of 2.79 ± 0.02 mm (Fig. 1C). Similarly, the pistil length varied from 6.72 to 7.01 mm, with a mean value of 6.87 ± 0.02 mm, while pistil width was assessed between 0.64 and 0.95 mm, with a mean value of 0.83 ± 0.02 mm (Fig. 1D; Table 3).

Anthesis and anther dehiscence

The observation started with the flower opening initiated with a slit at the bud's top, which widened gradually and took 2 h for complete opening (Fig. 3). Maximum flowers per inflorescence were opened during 6:00–10:00 am, which accounted for 81.33 of the totals and peaked between 8:00 and 10:00, while no substantial flowers opened after 12 pm (Fig. 4). Anthers were

observed to dehisce either during or shortly before anthesis. Moreover, the flowering pattern was asynchronous and took an average of 17 to 20 days to reach the flower opening stage.

Pollen biology

The average number of inflorescences produced per tree was 42.41 ± 5.26 , whereas mean total flower production per inflorescence was 383.47 ± 16.33 , and the average flower production per tree was $15,927.73 \pm 1470.82$. The pollen production per anther oscillated between 594 and 663 with an average of 636 ± 12.21 , while pollen grains per flower and inflorescence were 6370 ± 121.94 and $24.49 \times 10^5 \pm 99,855$, respectively. Likewise, number of pollens per tree was $10.19 \times 10^7 \pm 95.37 \times 10^5$ (Table 3). The number of ovules in each flower varied from 3 to 8, with a mean value of 5.81 ± 1.14 (Fig. 1H, Table 3). Hence, Pollen: ovule ratio was 1096.38 ± 108.70 , which indicated the possibility of autogamy (Table 3).

The shape of pollen grains appeared to be spherical in shape. Pollen viability was found $96.67 \pm 1.6\%$ to $98.26 \pm 1.2\%$ viable at the time of anthesis. Subsequently, pollen viability was examined with a 3 h difference. It remained almost identical (95.12 ± 1.14) after 3 h of anthesis. Pollen viability was reduced to 57.18 ± 5.91 and 33.38 ± 6.2 after 6 and 9 h of anthesis, respectively. No further viable pollen was recorded after 9 h of anthesis. Additionally, pollen germination was not observed in any of the concentrations mentioned.

Stigma receptivity

The stigma of a freshly opened flower was found receptive and continued up to 15 h after anthesis. To investigate stigma receptivity prior to anthesis, bud was divided into nine stages depending on their maturity and development (Fig. 5). Genital organs, i.e., stamen and pistil, marked development during stage 3 and thereafter (Figs 2 and 6). Stigma started active bubbling during stage 8





Figure 1. Microscopic view of floral whorls (Sepal (A), Petal (B), Pistil (C), Stamen (D)); Longitudinal view of mature anther (E); Longitudinal view of dehiscent anther (F); Stigma morphology during receptivity (G); Ovules in flower (H).

Table 2. Floristic features of *M. azedarach* L

Items	CL (mm)	CW (mm)	PL (mm)	PW (mm)	SL (mm)	SW (mm)	P _L (mm)	P _W (mm)
Mean	1.70 ± 0.05	1.00 ± 0.05	14.53 ± 0.18	3.52 ± 0.04	7.16 ± 0.05	2.79 ± 0.02	6.87 ± 0.02	0.83 ± 0.02
Range	1.67–1.73	0.8–0.94	12.42–15.63	3.17–4.15	6.97–7.39	2.66–2.91	6.72–7.01	0.64–0.95
N	150	150	150	150	150	150	150	150

Notes: CL, Calyx length; CW, Calyx width; PL, Petal length; PW, Petal width; SL, Stamen length; SW, Stamen width; P_L, Pistil length; P_W, Pistil width

**Figure 2.** Sequence of staminal tube development in *M. azedarach* flowers from bud to anthesis.**Table 3.** Floral characteristics of *M. azedarach* L

Items	I _t	F _t	P _a	P _f	P _i	P _t	Ovules	P/O
Mean	42.41 ± 5.26	15,927.73 ± 1470.82	636 ± 12.21	6370 ± 121.94	24.49 × 10 ⁵ ± 99,855	10.19 × 10 ⁷ ± 95.37 × 10 ⁵	5.81 ± 1.14	1096.38 ± 108.70
Range	25.98–69.92	10,936–23,912	594–663	5942–6634	21.04 × 10 ⁵ –30.17 × 10 ⁵	7.12 × 10 ⁷ –1.57 × 10 ⁸	3 to 8	894.72–1251.71
N	50	50	50	50	50	50	50	50

Notes: N, Sample size; I_t, Inflorescence per tree; F_t, Flowers per tree; P_a, Pollens per anther; P_f, Pollens per flower; P_i, Pollens per inflorescence; P_t, Pollens per tree; P/O, Pollens – ovule ratio.

**Figure 3.** Sequence of flower opening (anthesis) in *M. azedarach* L.




Figure 4. Anthesis in *Melia azedarach* in relation to time interval.

Notes: a: 06:00–08:00; b: 08:00–10:00; c: 10:00–12:00; d: 12:00–14:00; e: 14:00–16:00; A: 13–16 March; B: 17–20 March; C: 21–24 March; and D: 25–28 March.

and after that, i.e., 2 to 3 days before attaining anthesis, indicating receptivity of stigma. During its receptive phase, stigma was found to be wet, green and oily, while a dry and shrunken state confirms the end of stigmatic receptivity (Fig. 1G).

Floral visitors

Insect orders such as Hymenoptera, Diptera, Thysanoptera and Lepidoptera were noticed and identified from the flowers of *M. azedarach* (Fig. 7). The most common visitors to flowers were *Apis* spp. and syrphid flies. Attractive colour and subtle aroma were the prime reasons for a floral draw. Maximum visits in terms of numbers occurred between 8 am and 10 am, which coincided with anthesis period; however, infrequent visitors were observed during the evening (14:00 to 16:00), as illustrated in Fig. 8. In addition, observations of floral visitors are summarized in Table 4.

Breeding behaviour

One-way ANOVA result showed that fruit sets under natural pollination (NP) were significantly higher compared to open pollination (OP) or cross pollination ($F = 101.42$, $P = 0.00$). However, there was no substantial variation reported in fruit set percentage under natural pollination as well as self-pollination (SP) ($F = 2.08$, $P = 0.158$), which is strong evidence in favour of self-pollination. Interestingly, 1.65 per cent of fruit set was recorded in exposed emasculated flowers (open pollination), demonstrating the ability of cross-pollination. The proportion of fruit set was deficient; fruit set percentages of NP, SP and OP were 6.60, 5.77 and 1.65%, respectively (Fig. 9).

Discussion

Flowering phenology and reproductive biology are crucial for planning conservation strategies, efficient breeding approaches,




Figure 5. Sequence of floral bud development in *M. azedarach* L.




Figure 6. The sequence of pistil development in *M. azedarach* flowers from bud to anthesis.

and large-scale cultivation measures (Khanduri *et al.*, 2019). The current study provides in-depth details on reproductive biology of *M. azedarach* L. Floral phenology of *M. azedarach* under Bhubaneswar locality demonstrated that flowering began in the middle of March and continued until May, with a peak in late March (Table 1); However, Syamsuwida *et al.* (2012) observed flower initiation in August and this process progressed through generative buds to flower burst, typically from September to October. Subsequently, early fruits began to form in October to November, with fruits reaching physiological maturity during January to February. Similarly, Cavusoglu and Sulusoglu (2015) studied floral phenological growth of *M. azedarach* in Turkey,

and reported bud development during Mid-March, followed by onset of first flowering in late April coincided with leaf development. Full flowering was observed till the end of May concurrently with the presence of old fruits from the previous year still on the tree. Generally, flowering pattern in *M. azedarach*, was asynchronous, i.e., new flowers were developing at different times on the same individual, resulting in flowers with all stages of development apparent within an inflorescence. Therefore, it is challenging to determine particular climatic factors causing flowering. However, climatic factors like temperature, precipitation, elevation, soil water availability, and day length may influence and stimulate floral initiation (van Schaik, 1993).



Figure 7. a. *Apis* spp. visiting the flower of *M. azedarach*; **b:** Thrips (*Liothrips morulus*) identified from the *M. azedarach* flower.




Figure 8. Frequency of the numbers of floral visitors in *Melia azedarach* L.

M. azedarach produces cymose, most often axillary inflorescences bearing deep violet to pale violet, faintly scented flowers. The floral characteristics and inflorescence are almost identical to those of other closely related species in the genus, like *Melia dubia*, as reported by Johar *et al.*, 2015. There were minute variations in blossom size and colour among plants during a similar season. It appears that microclimatic changes and genetic makeup of plants caused such variations. Syamsuwida *et al.* (2012) in Maha Neem (*M. azedarach*), Johar *et al.* (2015) in Malabar Neem (*Melia composita*), and Dhillon *et al.* (2004) in Neem (*Azadirachta indica*) recorded similar observations. In addition, the current study, along with previous observations by Troup (1921) found no evidence supporting the andro-monoecious categorization of *M. azedarach*. However, Styles and Khosla (1976) had categorized it as an andro-monoecious species based on the production of two types of flowers (hermaphrodite and male), but contrary to these assertions, this study failed to uncover any distinct features among the flowers of the species.

Anthesis, anther dehiscence and receptivity of stigma are extremely important events in flower developmental process, as well as determining factors of pollination success. Maha Neem followed a diurnal pattern of anthesis, with a peak occurring between 8:00 and 10:00 am, coinciding with anther dehiscence, and rather no substantial flower opening after 12 pm (Fig. 4).

Our finding is consistent with the research of Johar *et al.*, 2015, who found similar diurnal anthesis in another species under Melioideae *i.e.* *M. dubia*. Our study also revealed that stigma starts receptivity before anthesis and continues its invitation up to 12 h after anthesis, whereas anther begins dehiscence during or shortly before anthesis. A similar report on stigma receptivity spans from 8:00 am to 11:00 am, exhibiting a receptivity rate of 66%, which subsequently decreases to 40% by noon on *M. azedarach* (Syamsuwida *et al.*, 2012). Pollen from freshly dehisced anther was highly viable, with a viability range of 96.67 ± 1.6 to 98.26 ± 1.2 per cent. However, its viability deteriorates further, and no viable pollen was noticed after 9 h of anthesis. Hence, Pollens from recently opened flowers and emasculation prior to reaching stigmatic receptivity, typically 4 to 5 days before anthesis, are very important for effective and successful hybridization in *M. azedarach*.

Pollen features *viz.* number of pollens per flower, pollen: ovule ratio, and viability were found to be sufficient for fertilization in *M. azedarach*. The number of flowers/trees and overall pollen output/tree varied among individuals. A higher number of primary and secondary branches maximizes total flower production, which in turn maximizes total pollen output, resulting in pollen competition and successful fertilization. A great load of pollen also increases competition among male gametophytes to fertilize

Table 4. Floral visitors of *Melia azedarach* L.

Species	Order	Time duration				
		06:00–08:00	08:00–10:00	10:00–12:00	12:00–14:00	14:00–16:00
<i>Thrips</i>	Thysanoptera	2	5	5	4	1
<i>Apis spp.</i>	Hymenoptera	10	21	13	12	3
<i>Vespa spp.</i> (Hornet Wasp)	Hymenoptera	8	18	11	0	0
<i>Lasius spp.</i> (Black ant)	Hymenoptera	0	12	0	0	0
<i>Sphaerophoria philanthus</i> (Syrphid fly)	Diptera	5	17	16	4	5
<i>Danaus genutia and others</i>	Lepidoptera	6	14	11	6	2




Figure 9. Effect of pollination treatments on the fruit sets of *Melia azedarach* L.

the available ovules and improves seed quality (Tangmitcharoen and Owens, 1997). Based on observations during the study, the total fruit set (reproductive success) from the available flowers was substantially low, because a notable number of flowers fell prematurely, impeding successful reproduction. Generally, increased number of primary and secondary branches maximizes overall flower production, consequently enhancing total pollen output and facilitating successful fertilization, which in turn promotes fruit set and reproductive success. Khanduri *et al.* (2019) similarly identified a robust positive correlation ($r = 0.989$) between lateral shoot production and total inflorescence yield in trees. There was a positive relationship between stigma morphological characters and receptivity. Stigma was found to be wet, green and oily during its receptive phase, while a dry and shrunken state confirms the end of stigmatic receptivity. Khanduri *et al.* (2019) also found a connection between stigmatic morphological traits and its receptivity with straight style, enlarged moist stigma, oily and green in *C. capitata*. Similarly, Pollen viability was negatively related to effective pollination period, i.e., time elapses between pollen shedding and fertilization success. During this time, pollen grains were subjected to various environmental stresses, particularly temperature and humidity, which may have hampered their ability to produce vigorous offspring. Pollen germination was attempted with various concentrations (50, 100 and 150 μ l) of IAA, IBA and NAA, but no pollen germination was found in any of the concentrations mentioned. It could be due to the refined morphological and functional characteristics of stigma (Arceo-Gomez *et al.*, 2011; Aronne *et al.*, 2012; Smitha and Thondaiman, 2016). The anatomical features of sexual organs, such as pollen and stigma and pollen-stigmatic interaction, should be explored in detail to comprehend pollen germination thoroughly.

Pollination experiments revealed that *M. azedarach* is primarily a self-pollinated species where the majority of fruits are produced by self and natural mechanism of pollination and only a few under open pollination. The non-significant difference in fruit set under natural and self-pollination suggests self-compatible and

spontaneous autogamy of *M. azedarach* (Fig. 9). Similar obligate reliance on autogamy in *M. azedarach* has also been supported by Syamsuwida *et al.* (2012) and Waites and Agren (2006) and they expect that the position of anther and stigma makes it possible. Self-pollination may be caused due to homogamy, which is occurrence of genital organs maturity, viz., stamen, and pistil at the same time. Further, temporal overlapping of stigmatic receptivity and dehiscent anther promotes self-pollination significantly. On the other hand, fruit set in exposed emasculated flowers demonstrates cross-pollination ability in Maha-Neem. Gituru *et al.* (2002) stated that densely crowded flowers attract floral visitors and pollinate open flowers present in inflorescence. Likewise, insects were predominantly visiting *M. azedarach* during anthesis, which may have aided fruit set in open emasculated flowers. Pollinator attraction, defined as the diversity and frequency of floral visits, was found to be limited in *M. azedarach*; however, subtle aroma and attractive flower colour encourage the probability of floral visits and subsequent fruit set through xenogamy. Peak visits observed during the period between 8 am and 10 am, coinciding with period of maximum anthesis and minimal during afternoon period (Fig. 8). Observation of floral visitors revealed that Maha-Neem is entomophilous, frequently visited by Apis and Syrphid flies. Nevertheless, lower pollen counts and a lack of nectar reward resulted in fewer butterfly sightings. These observations concur with Styles and Khosla (1976), who also reported frequent visiting of bees, flies, butterflies and thrips in *M. dubia* and *M. azedarach* respectively.

Conclusion

The floral biology and breeding system of *M. azedarach* study revealed that floral architecture of the species supports autogamy pollination. Therefore, seeds can be adopted to achieve true-to-type genotype or accession. However, species can also sustain hybridization with a systematic breeding technique, which involves early-stage emasculation, typically 4 to 5 days before anthesis, and pollens from freshly opened flowers since pollen loses its viability quickly. Our findings also demonstrated a

positive correlation between number of flowers, inflorescence numbers, pollen production and ultimately reproductive success. In addition, it was observed that abundant flowering and fruiting occur both years, consequently, seeds were available each year profusely. Information on reproductive biology of *M. azedarach* can be helpful for setting up conservation and effective management strategies, as well as controlled crossing programmes for producing interspecific hybrids and sustainable cultivation of lucrative commercial species.

Supplementary material. The supplementary material for this article can be found at <https://doi.org/10.1017/S1479262124000376>

Acknowledgments. The authors are grateful to the Head of the Department of Forest Biology and Tree Improvement, Odisha University of Agriculture and Technology (Odisha), India for providing the necessary facilities during the study. The authors also duly acknowledge the use of the facilities provided by the State Forest Department, Government of Odisha, India.

Authors contributions. Experimentation including conception and design - S.R.M; Data collection and analysis - S.K.D and M.R.K; Writing and original draft preparation - S.R.M and H.D; Final review and editing - S.R.M and S.K.D. Authors have read and approved the final manuscript.

Funding statement. This research received no specific grant from any funding agency in the public, commercial, or not-for-profit sectors.

Competing interests. The authors declare that they have no conflict of interest.

References

- Arceo-Gómez G, Martínez ML, Parra-Tabla V and García-Franco JG (2011) Anther and stigma morphology in mirror-image flowers of *Chamaecrista chamaecristoides* (Fabaceae): implications for buzz pollination. *Plant Biology* **13**, 19–24.
- Aronne G, Giovanetti M and De Micco V (2012) Morphofunctional traits and pollination mechanisms of *Coronilla emerus* L. flowers (Fabaceae). *The Scientific World Journal* **1**, 381575.
- Brewbaker JL and Kwack BH (1963) The essential role of calcium ion in pollen tube growth. *American Journal of Botany* **50**, 859–865.
- Cavusoglu A and Sulusoglu M (2015) Phenological and morphological characteristics of *Melia azedarach* L. in Kocaeli city in Turkey Horticulture. *Scientific Papers. Series B. Horticulture* **59**, 311–318.
- Cruden RW (1977) Pollen-Ovule ratios: a conservative indicator of breeding systems in flowering plants. *Evolution* **31**, 32–46.
- Dafni A (1992) *Pollination Ecology: A Practical Approach*. New York: Oxford University Press.
- Dhillon RS, Bisla SS and Hooda MS (2004) Phenology and breeding system of neem. *Indian Journal of Ecology* **31**, 30–32.
- Duong DV, Missanjo E and Matsumura J (2017) Variation in intrinsic wood properties of *Melia azedarach* L. planted in northern Vietnam. *Journal of Wood Science* **63**, 560–567.
- Gituru WR, Wang Q, Wang Y and Guo Y (2002) Pollination ecology, breeding system and conservation of *Caldesia grandis* (Alis mataceae), an endangered mash plant in China. *Botanical Bulletin of Academia Sinica* **43**, 230–231.
- Global Agricultural Information Network report. (No. IN9033) (2019) USDA Foreign Agricultural Service, USA.
- Johar V, Dhillon RS, Bangarwa KS and Ajit HA (2015) Phenological behavior and reproductive biology of *Melia composita* Benth. *Indian Journal of Agroforestry* **17**, 62–67.
- Khan GS and Chaudhry AQ (2007) Effect of spacing and plant density on the growth of poplar (*Populus deltoides*) trees under agroforestry system. *Pakistan Journal of Agricultural Science* **44**, 321–327.
- Khanduri VP, Kumar KS and Sharma CM (2016) Role of pollen production in mating success in some tropical tree species. *Brazilian Journal of Botany* **38**, 107–112.
- Khanduri VP, Sukumaran A and Sharma CM (2019) Reproductive biology of *Cornus capitata* Wall. ex Roxb.: a native species in East Asia. *Journal of Forestry Research* **30**, 2039–2050.
- Kuniyal CP, Bhadula SK and Prasad P (2003) Flowering, seed characteristics and seed germination behaviour in the populations of a threatened herb *Aconitum atrox* (bruhl) Muk. (Ranunculaceae). *International Journal of Environmental Science* **71**, 29–36.
- Lee SL, Ng KKS, Saw LG, Lee CT, Muhammad N, Naoki T, Yoshihiko T and Koskela J (2006) Linking the gaps between conservation research and conservation management of rare dipterocarps: a case study of *Shorea lumutensis*. *Biological Conservation* **131**, 72–92.
- Martén S and Quesada M (2001) Phenology, sexual expression, and reproductive success of the rare neotropical palm *Geonoma eptiolata*. *Biotropica* **33**, 596–605.
- Murugan R, Shivanna KR and Rao RR (2006) Pollination biology of *Aristolochia tagala*, a rare species of medicinal importance. *Current Science* **91**, 795–798.
- Newstrom LE, Frankie GW and Baker HG (1994) A new classification for plant phenology based on flowering patterns in lowland tropical rain forest trees at La Selva, Costa Rica. *Biotropica* **26**, 141.
- Nivedita M, Radha R, Megala S and Nithya S (2019) Review on *Melia azedarach* Linn. *World Journal of Pharmacy and Pharmaceutical Sciences* **8**, 363–391.
- Osborn NM, Kevan PG and Lane M (1988) Pollination biology of *Opuntia polyantha* and *Opuntia phaeacantha* (Cactaceae) in Southern Colorado. *Plant Systematics and Evolution* **159**, 85–94.
- Rahman MK, Asaduzzaman M, Rahman MM, Das AK and Biswas SK (2014) Physical and mechanical properties of Ghora neem (*Melia azedarach*) plywood. *Bangladesh Journal of Scientific and Industrial Research* **49**, 47–52.
- Smitha GR and Thondaiman V (2016) Reproductive biology and breeding system of *Saraca asoca* (Roxb.) De Wilde: a vulnerable medicinal plant. *Springer Plus* **5**, 2025.
- Sokal RE and Rohlf FJ (1995) *Biometry: The Principles and Practice of Statistics in Biological Research*. San Francisco: Freeman.
- Srinivasan M, Varadha Raj S and Eswari A (2018) Demand and supply analysis of roundwood in India. *Madras Agricultural Journal* **105**, 1–3.
- Styles BT and Khosla PE (1976) Cytology and reproductive biology of Meliaceae. In Burley J and Styles AT (eds), *Tropical Trees: Variation, Breeding and Conservation*. London, UK: Academic Press, pp. 61–67.
- Syamsuwida D, Palupi ER, Siregar IZ and Indrawan A (2012) Flower initiation, morphology, and developmental stages of flowering-fruiting of Mindi (*Melia azedarach* L.). *Journal Manajemen Hutan Tropika* **18**, 10–17.
- Tandon R, Shivanna KR and Mohan Ram HY (2003) Reproductive biology of *Butea monosperma* (Fabaceae). *Annals of Botany* **92**, 715–723.
- Tangmitcharoen S and Owens JN (1997) Floral biology, pollination, pistil receptivity, and pollen tube growth of teak (*Tectona grandis* Linn f.). *Annals of Botany* **79**, 227–241.
- Troup RS (1921) *The Silviculture of Indian Trees*. Dehradun: ICFRE press, Dehradun, India.
- van Schaik CP (1993) The phenology of tropical forests: adaptive significance and consequences for primary consumers. *Annual Review of Ecology and Systematics* **24**, 353–377.
- Venson I, Silva Guzmán JA, Fuentes Talavera FJ and Richter HG (2008) Biological, physical, and mechanical wood properties of Paraiso (*Melia azedarach*) from a roadside planting at huaxtla, Jalisco, Mexico. *Journal of Tropical Forest Science* **20**, 38–47.
- Waites AR and Agren JA (2006) Stigma receptivity and effects of prior self-pollination on seed set in Tristylous *Lythrum Salicaria* (Lytraceae). *American Journal of Botany* **93**, 142–147.
- Waller JM, Ritchie BJ and Holderness M (1998) *Plant Clinic Handbook*. New York, USA: CAB International.