

The 13th European Nutrition Conference, FENS 2019, was held at the Dublin Convention Centre, 15–18 October 2019

Dietary assessment in the German National Cohort (GNC)

Sven Knüppel¹, Matthias Clemens¹, Johanna Conrad², Sylvia Gastell¹, Karin Michels³, Michael Leitzmann⁴, Lilian Krist⁵, Tobias Pischon⁶, Gerard Krause⁷, Wolfgang Ahrens^{8,9}, Nina Ebert¹⁰, Karl-Heinz Jöckel¹¹, Alexander Kluttig¹², Nadia Obi¹³, Rudolf Kaaks¹⁴, Wolfgang Lieb¹⁵, Sabine Schipf^{16,17}, Hermann Brenner^{14,18}, Thorsten Heuer¹⁹, Ulrich Harttig¹, Jakob Linseisen^{20,21}, Ute Nöthlings² and Heiner Boeing¹

¹German Institute of Human Nutrition Potsdam-Rehbruecke (DIfE), Nuthetal, Germany,

²University of Bonn, Bonn, Germany,

³University of Freiburg, Freiburg, Germany,

⁴University of Regensburg, Regensburg, Germany,

⁵Charité-Universitätsmedizin Berlin, Berlin, Germany,

⁶Max-Delbrück Center for Molecular Medicine in the Helmholtz Association (MDC), Berlin, Germany,

⁷Helmholtz Centre for Infection Research, Braunschweig, Germany,

⁸Leibniz Institute for Prevention Research and Epidemiology - BIPS, Bremen, Germany,

⁹University Bremen, Bremen, Germany,

¹⁰Leibniz Institute for Diabetes Research at Heinrich Heine University Düsseldorf, Düsseldorf, Germany,

¹¹University of Duisburg-Essen, Essen, Germany,

¹²Martin Luther University Halle-Wittenberg, Halle (Saale), Germany,

¹³University Medical Center Hamburg-Eppendorf, Hamburg, Germany,

¹⁴German Cancer Research Center (DKFZ), Heidelberg, Germany,

¹⁵Christian-Albrechts University of Kiel, Kiel, Germany,

¹⁶University Medicine Greifswald, Greifswald, Germany,

¹⁷German Center for Diabetes Research (DZD), Greifswald, Germany,

¹⁸National Center for Tumor Diseases (NCT), Heidelberg, Germany,

¹⁹Max Rubner-Institut (MRI), Karlsruhe, Germany,

²⁰Helmholtz Centre Munich, Neuherberg, Germany and

²¹UNIKA-T Augsburg, Augsburg, Germany

Abstract

We describe a novel dietary assessment strategy to estimate usual food intake in the ongoing large-scale multi-center German National Cohort (GNC). The dietary assessment is based on three 24 h food lists (24h-FL) and a food frequency questionnaire (FFQ) enriched by information from the representative German National Nutrition Survey II (NVS II). The novelty of this dietary assessment strategy is based on separating the probability of food intake from daily consumption amounts. The probability of consumption is estimated from 24h-FLs used in the GNC. To estimate daily consumption amounts, the already collected data of the NVS II are used. The 24h-FL simplifies the question on food consumption for all foods asked to consumption or not and so the questionnaire can be completed in about 10 minutes, reducing the burden on study participants. As proof of concept, we applied the assessment strategy to pretest data collected in 2012 to 2013 to assess the feasibility of the instruments. In brief, the novel dietary assessment strategy comprises three steps. First, the individuals' consumption probability is estimated by three 24h-FLs and one FFQ applying a logistic linear mixed model adjusted for characteristics of the participants. Second, person-specific daily consumption amounts are estimated from the NVS II applying a linear mixed model taking the characteristics of the participants into account. Third, usual food intake is estimated by the consumption probability multiplied by person-specific daily amounts. Usual intake of 41 food groups in 318 men and 377 women were estimated. Of those participants who completed the first 24h-FL, 84.4, and 68.5% completed the second and third 24h-FL, respectively. No associations were observed between probability to participate and lifestyle factors. The estimated usual food intake distributions were in a plausible range as shown by comparing the estimated energy intake to the energy needs approximated by estimated total energy expenditure. Total energy was estimated to be 2,707 kcal/day for men and 2,103 kcal/day for women. With a few exceptions, the estimated food-based consumption probabilities did not differ considerably between men and women. The differences in energy intake between men and women were mainly due to their differences in the estimated person-specific daily amounts. As a conclusion, plausible but not validated values for usual food intake were derived in the pretest study, so that the combination of three repeated 24h-FLs, an FFQ and person-specific daily amounts from an external source is a feasible strategy for dietary assessment.

Conflict of Interest

There is no conflict of interest.