



# On Deformations of 1-motives

A. Bertapelle and N. Mazzari

*Abstract.* According to a well-known theorem of Serre and Tate, the infinitesimal deformation theory of an abelian variety in positive characteristic is equivalent to the infinitesimal deformation theory of its Barsotti–Tate group. We extend this result to 1-motives.

## 1 Introduction

Let  $R$  be an artinian local ring with maximal ideal  $\mathfrak{m}$  and perfect residue field  $k$  of positive characteristic  $p$ . Let  $\mathcal{M}_1(R)$  denote the category of (smooth) 1-motives over  $R$ . For any 1-motive  $M$  (resp. Barsotti–Tate group  $\mathcal{B}$ ) over  $R$ , let  $M_0$  (resp.  $\mathcal{B}_0$ ) denote its base change to  $k = R/\mathfrak{m}$ . Let  $\text{Def}(R, k)$  denote the category whose objects are triples  $(M_0, \mathcal{B}, \varepsilon_0)$  where  $M_0$  is a 1-motive over  $k$ ,  $\mathcal{B}$  is a Barsotti–Tate group over  $R$  and  $\varepsilon_0: \mathcal{B}_0 \rightarrow M_0[p^\infty]$  is an isomorphism of  $\mathcal{B}_0$  with the Barsotti–Tate group of  $M_0$ . The aim of this paper is to prove the following theorem.

*Theorem 1.1* *The functor*

$$(1.1) \quad \begin{aligned} \Delta_R: \mathcal{M}_1(R) &\longrightarrow \text{Def}(R, k) \\ M &\longmapsto (M_0, M[p^\infty], \text{natural } \varepsilon_0), \end{aligned}$$

*is an equivalence of categories.*

This result generalises the well-known Serre–Tate equivalence for abelian schemes over artinian local rings  $R$  as above (cf. [6, Theorem 1.2.1], [8, V, Theorem 2.3]). Arguments in [6, 8] do not extend directly to the case of 1-motives but are used to get some intermediate results. Note that the proof of fullness (Proposition 4.3) and essential surjectivity (Proposition 4.5) uses weights, a careful study of the case of 1-motives of the form  $[\mathbb{Z} \rightarrow \mathbb{G}_m]$  (Lemma 2.4), which is orthogonal to the classical case of abelian schemes, and Galois descent arguments (Lemma 4.4). Another essential ingredient is that tori and étale group schemes over  $k$  lift uniquely to  $R$ .

As an application, in Section 4.3 we extend the description of the formal moduli space of an ordinary abelian variety over an algebraically closed field and its Serre–Tate coordinates to 1-motives. Consequently, canonical liftings of (ordinary) 1-motives over  $k$  to 1-motives over  $W(k)$  exist (Proposition 4.10).

---

Received by the editors October 10, 2017; revised November 17, 2017.

Published electronically March 22, 2018.

AMS subject classification: 14L15, 14C15, 14L05.

Keywords: 1-motive, Barsotti–Tate group.

**Notation**

Let  $p^s, s \geq 1$ , denote the characteristic of  $R$ . Then  $R$  is canonically endowed with the structure of a finite  $W_s(k)$ -algebra, where  $W_s(k) = W(k)/(p^s)$  denotes the ring of Witt vectors of length  $s$ . We also fix a positive integer  $n$  such that  $(1 + \mathfrak{m})^{p^n} = \{1\}$ . For any  $R$ -group scheme  $G$ , we identify  $G(R) = \text{Hom}_R(\text{Spec } R, G)$  with  $\text{Hom}_{R\text{-gr}}(\mathbb{Z}, G)$  by mapping  $a \in G(R)$  to the morphism  $u: \mathbb{Z} \rightarrow G$  such that  $u(1) = a$ .

**2 Barsotti–Tate Groups and 1-motives**

Let  $\mathcal{M}_1(R)$  be the category of (smooth) 1-motives over  $R$ . Its objects are two term complexes of commutative  $R$ -group schemes  $M = [u: L \rightarrow G]$  in degrees  $-1$  and  $0$ , where  $G$  is extension of an abelian scheme  $A$  by a torus  $T$ , and  $L$  is locally for the étale topology on  $R$  isomorphic to  $\mathbb{Z}^r$  for some non-negative integer  $r$ . Recall that a 1-motive has a natural weight filtration

$$0 \subseteq W_{-2}M = [0 \rightarrow T] \subseteq W_{-1}M = [0 \rightarrow G] \subseteq W_0M = M.$$

Since morphisms of 1-motives respect filtrations,  $\mathcal{M}_1(R)$  is a filtered category. We will denote by  $\mathcal{M}_1(R)_{\leq i}$  the full subcategory of  $\mathcal{M}_1(R)$  consisting of 1-motives such that  $M = W_iM$ . Given a 1-motive  $M$ , let  $M_{\text{ab}} = [u_{\text{ab}}: L \rightarrow A]$ , where  $u_{\text{ab}}$  is the composition of  $u$  with the canonical morphism  $G \rightarrow A$ . Let  $\mathcal{M}_1(R)_{\geq -1}$  be the full subcategory of  $\mathcal{M}_1(R)$  consisting of 1-motives  $M = M_{\text{ab}}$ . Similarly  $\text{Def}(R, k)_{\geq -1}$  (resp.  $\text{Def}(R, k)_{\leq -1}$ ) denotes the full subcategory of  $\text{Def}(R, k)$  consisting of objects such that  $M_0$  is in  $\mathcal{M}_1(k)_{\geq -1}$  (resp. in  $\mathcal{M}_1(k)_{\leq -1}$ ).

For any  $m \in \mathbb{N}$ , consider the cone of the multiplication by  $m$  on  $M$ :

$$M/mM: L \xrightarrow{\begin{pmatrix} -u \\ -m \end{pmatrix}} G \oplus L \xrightarrow{(-m, u)} G,$$

where  $L$  is in degree  $-2$ , and let  $M[m] = H^{-1}(M/mM)$  (see [4, 10.1.4] or [2, §1.3] with a different sign convention); it is a finite and flat  $R$ -group scheme and it fits in the diagram

$$(2.1) \quad \begin{array}{ccccccc} & & & L & \xlongequal{\quad} & L & \\ & & & \downarrow \begin{pmatrix} -u \\ -m \end{pmatrix} & & \downarrow -m & \\ \tilde{\eta}_{M[m]}: & 0 & \longrightarrow & G[m] & \longrightarrow & \text{Ker}((-m, u)) & \longrightarrow & L & \longrightarrow & 0 \\ & & & \parallel & & \downarrow & & \downarrow & & \\ \eta_{M[m]}: & 0 & \longrightarrow & G[m] & \longrightarrow & M[m] & \longrightarrow & L/mL & \longrightarrow & 0. \end{array}$$

**Remark 2.1** Let us consider

$$\xi_{G[m]}: 0 \longrightarrow G[m] \longrightarrow G \xrightarrow{m} G \longrightarrow 0.$$

By direct computations one gets  $\tilde{\eta}_{M[m]} = -u^* \xi_{G[m]}$  as elements of  $\text{Ext}_R(L, G[m])$ .

Since the push-out of the extension  $\tilde{\eta}_{M[p^r]}$  along the canonical morphism  $G[p^r] \rightarrow G[p^{r+1}]$  is isomorphic to  $p \cdot \tilde{\eta}_{M[p^{r+1}]}$ , there is a morphism of complexes

$\eta_{M[p^r]} \rightarrow \eta_{M[p^{r+1}]}$  for any  $r \geq 1$ . One then gets a short exact sequence of Barsotti–Tate groups (BT groups for short)

$$(2.2) \quad \eta_{M[p^\infty]}: 0 \longrightarrow G[p^\infty] \longrightarrow M[p^\infty] \longrightarrow L[p^\infty] \longrightarrow 0,$$

where  $L[p^\infty] := L \otimes_{\mathbb{Z}} \mathbb{Q}_p / \mathbb{Z}_p$ .

**Lemma 2.2** *Let  $M_0 = [u_0: L_0 \rightarrow G_0]$ , let  $M'_0$  be 1-motives over  $k$ , and let  $\varphi_0: M_0 \rightarrow M'_0$  be a morphism of 1-motives. Denote by  $T_0$  the maximal subtorus of  $G_0$ .*

- (i) *Any finite and flat  $R$ -group scheme  $F$  that lifts  $M_0[m]$  is endowed with a unique weight filtration that lifts the weight filtration  $T_0[m] \subseteq G_0[m] \subseteq M_0[m]$ .*
- (ii) *Any morphism of finite flat  $R$ -group schemes  $F \rightarrow F'$  that lifts  $\varphi_0[m]$  respects filtrations.*
- (iii) *Any BT group over  $R$  that lifts  $M_0[p^\infty]$  admits a unique filtration that lifts the weight filtration on  $M_0[p^\infty]$  and any morphism of BT groups over  $R$  which lifts  $\varphi_0[p^\infty]$  respects filtrations.*
- (iv) *If  $M_0$  lifts to a 1-motive  $M$  over  $R$ , then the natural weight filtration on  $M[m]$  agrees with the one induced by the weight filtration on  $M_0$ .*

**Proof** Let  $F$  be a finite flat  $R$ -group scheme lifting  $M_0[m]$ . Let  $L$  be the unique étale lifting of  $L_0$  over  $R$ . We claim that  $F$  fits into a short exact sequence

$$0 \longrightarrow E \longrightarrow F \longrightarrow L/mL \longrightarrow 0,$$

which lifts the sequence  $\eta_{M_0[m]}$  for  $M_0$  in diagram (2.1). Indeed, there exists a unique lifting  $f: F \rightarrow L/mL$  of the morphism  $M_0[m] \rightarrow L_0/mL_0$ , since  $L/mL$  is étale, and we set  $E = \text{Ker } f$ . Note further that  $f$  factors through the epimorphism  $\pi_F: F \rightarrow F^{\text{ét}} = F/F^\circ$  with  $F^\circ$  the identity component of  $F$ . Let  $f^{\text{ét}}: F^{\text{ét}} \rightarrow L/mL$  satisfy  $f^{\text{ét}} \circ \pi_F = f$ . The morphism  $f_0^{\text{ét}}$  is an epimorphism of finite étale  $k$ -group schemes, hence so is  $f^{\text{ét}}$ . It follows that  $f$  is an fppf epimorphism, hence the short exact sequence as claimed.

Let  $L_0^*$  and  $F^*$  be the Cartier duals of  $L_0$  and  $F$ , respectively, and let  $T$  be the unique torus lifting  $T_0$ . We have seen that the canonical morphism  $M_0[m]^* \rightarrow L_0^*/mL_0^*$  lifts uniquely to a morphism  $F^* \rightarrow L^*/mL^*$  over  $R$  and hence, by Cartier duality, the immersion  $T_0[m] \rightarrow M_0[m]$  lifts uniquely to an immersion  $T[m] \rightarrow F$  over  $R$ . The composition  $T[m] \rightarrow F \rightarrow L/mL$  is the zero map, since its reduction modulo  $\mathfrak{m}$  is the zero map and  $L/mL$  is étale. The closed immersion  $T[m] \rightarrow F$  factors thus through  $E$ . By construction, the filtration

$$0 \subseteq T[m] \subseteq E \subseteq F$$

of  $F$  lifts the weight filtration on  $M_0[m]$ , so the existence of a filtration as in (i) is proved.

With similar arguments, one shows statement (ii). Statement (iv) and the uniqueness of the filtration in (i) follow from (ii), and statement (iii) is a formal consequence of (i) and (ii). ■

The classical Serre–Tate theorem states that deformations of an abelian variety over  $k$  only depend on deformations of its BT group. As we will now explain, the analogous result for 1-motives of the type  $[\mathbb{Z} \rightarrow \mathbb{G}_{m,k}]$  can explicitly be deduced from the

canonical isomorphism

$$(2.3) \quad R^* \xrightarrow{\sim} k^* \times (1 + \mathfrak{m}), \quad x \longmapsto (x_0, x/[x_0]),$$

where  $x_0$  is the reduction of  $x$  modulo  $\mathfrak{m}$  and  $[x_0] \in W_s(k)$  is the multiplicative representative of  $x_0$ . Recall that  $(1 + \mathfrak{m})^{p^n} = \{1\}$ , and hence  $1 + \mathfrak{m} = \mu_{p^n}(R) = \mu_{p^\infty}(R)$ .

We now consider BT groups that are extensions of  $\mathbb{Q}_p/\mathbb{Z}_p$  by  $\mu_{p^\infty}$ . By [8, Proposition 2.5, p. 180], the map

$$\Psi: 1 + \mathfrak{m} \longrightarrow \text{Ext}_R(\mathbb{Q}_p/\mathbb{Z}_p, \mu_{p^\infty}),$$

which associates with  $u \in 1 + \mathfrak{m}$  the push-out along  $u: \mathbb{Z} \rightarrow \mu_{p^\infty}$  of the sequence

$$0 \longrightarrow \mathbb{Z} \longrightarrow \mathbb{Z}[p^{-1}] \longrightarrow \mathbb{Q}_p/\mathbb{Z}_p \longrightarrow 0$$

is an isomorphism. In particular,  $p^n$  kills  $\text{Ext}_R(\mathbb{Q}_p/\mathbb{Z}_p, \mu_{p^\infty})$ . For any  $u \in R^*$ , let  $M^u = [u: \mathbb{Z} \rightarrow \mathbb{G}_{m,R}]$  and consider the homomorphism

$$(2.4) \quad \Phi: 1 + \mathfrak{m} \longrightarrow \text{Ext}_R(\mathbb{Q}_p/\mathbb{Z}_p, \mu_{p^\infty}),$$

which maps  $u \in 1 + \mathfrak{m} \subset R^*$  to the extension  $\eta_{M^u[p^\infty]}$  as in (2.2).

**Lemma 2.3** *We have  $\Phi = -\Psi$ ; hence,  $\Phi$  is an isomorphism.*

**Proof** Let  $\text{Ext}_{\mathbb{Z}/p^n\mathbb{Z}}(\cdot, \cdot)$  denote classes of extensions of  $p^n$ -torsion  $R$ -group schemes. By [8, p. 183], we have isomorphisms

$$(2.5) \quad \begin{aligned} \text{Ext}_R(\mathbb{Q}_p/\mathbb{Z}_p, \mu_{p^\infty}) &\simeq \varprojlim_m \text{Ext}_R(\mathbb{Z}/p^m\mathbb{Z}, \mu_{p^\infty}) \\ &\simeq \text{Ext}_R(\mathbb{Z}/p^n\mathbb{Z}, \mu_{p^\infty}) \simeq \text{Ext}_{\mathbb{Z}/p^n\mathbb{Z}}(\mathbb{Z}/p^n\mathbb{Z}, \mu_{p^n}), \end{aligned}$$

where the first isomorphism is induced by pull-back along  $\mathbb{Z}/p^m\mathbb{Z} \rightarrow \mathbb{Q}_p/\mathbb{Z}_p$ , the second isomorphism follows, since  $p^n$  kills  $\text{Ext}_R(\mathbb{Q}_p/\mathbb{Z}_p, \mu_{p^\infty})$ , and the third isomorphism is obtained by passing to kernels of the multiplication by  $p^n$ , since  $\mu_{p^\infty}$  is divisible. Note that the composition of the isomorphisms in (2.5) associates with an extension of  $\mathbb{Q}_p/\mathbb{Z}_p$  by  $\mu_{p^\infty}$  the corresponding sequence of kernels of the multiplication by  $p^n$ . Let

$$\Psi_n, \Phi_n : 1 + \mathfrak{m} \longrightarrow \text{Ext}_{\mathbb{Z}/p^n\mathbb{Z}}(\mathbb{Z}/p^n\mathbb{Z}, \mu_{p^n})$$

denote the composition of  $\Psi$  (resp.  $\Phi$ ) with the isomorphisms in (2.5). We are left to prove that  $\Phi_n = -\Psi_n$ .

Note that  $\Psi_n(u)$  can also be constructed taking first the push-out along  $u: \mathbb{Z} \rightarrow \mathbb{G}_{m,R}$  of  $\zeta: 0 \rightarrow \mathbb{Z} \rightarrow \mathbb{Z} \rightarrow \mathbb{Z}/p^n\mathbb{Z} \rightarrow 0$  and then considering the sequence of kernels of the multiplication by  $p^n$ . Indeed,  $u_*\zeta = \iota_n^* \iota_{n*} \Psi(u)$  with  $\iota_n: \mathbb{Z}/p^n\mathbb{Z} \rightarrow \mathbb{Q}_p/\mathbb{Z}_p$  and  $\iota: \mu_{p^\infty} \rightarrow \mathbb{G}_{m,R}$  the usual morphisms. Since Cartier duality induces the identity both on  $1 + \mathfrak{m} = \text{Hom}_{R\text{-gr}}(\mathbb{Z}, \mathbb{G}_m)$  and on  $\text{Ext}_{\mathbb{Z}/p^n\mathbb{Z}}(\mathbb{Z}/p^n\mathbb{Z}, \mu_{p^n})$ ,  $\Psi_n(u)$  is also represented by the sequence of cokernels of the multiplication by  $p^n$  of the sequence obtained via pull-back along  $u: \mathbb{Z} \rightarrow \mathbb{G}_{m,R}$  from the sequence  $\xi_{\mu_{p^n}}: 0 \rightarrow \mu_{p^n} \rightarrow \mathbb{G}_{m,R} \rightarrow \mathbb{G}_{m,R} \rightarrow 0$ . On the other hand,  $\Phi_n(u)$  is the sequence  $\eta_{M^u[p^n]}$  in diagram (2.1). By Remark 2.1 and diagram (2.1), we conclude that  $\Phi_n(u) = -\Psi_n(u)$ . ■

We now see how to interpret (2.3) in terms of deformations of 1-motives.

**Lemma 2.4** *Given a 1-motive  $M_0 = [u_0: \mathbb{Z} \rightarrow \mathbb{G}_{m,k}]$  and a  $\mathcal{B} \in \text{Ext}_R(\mathbb{Q}_p/\mathbb{Z}_p, \mu_{p^\infty})$  there is a unique 1-motive  $M = [u: \mathbb{Z} \rightarrow \mathbb{G}_{m,R}]$  that lifts  $M_0$  and whose BT group is isomorphic to  $\mathcal{B}$  as extension of  $\mu_{p^\infty}$  by  $\mathbb{Q}_p/\mathbb{Z}_p$ .*

**Proof** Note that if  $\mathcal{B}, \mathcal{B}'$  are two liftings of  $M_0[p^\infty]$ , there is at most one morphism  $\mathcal{B} \rightarrow \mathcal{B}'$  as extensions of  $\mathbb{Q}_p/\mathbb{Z}_p$  by  $\mu_{p^\infty}$ . We have to prove that the homomorphism

$$R^* \longrightarrow k^* \times \text{Ext}_R(\mathbb{Q}_p/\mathbb{Z}_p, \mu_{p^\infty}), \quad u \longmapsto (u_0, \eta_{M^u[p^\infty]})$$

is an isomorphism. Note that if  $u = [u_0], u_0 \in k^*$ , then  $u$  admits  $p^r$ -th roots for any  $r \geq 1$ , since  $k$  is perfect; hence,  $\eta_{M^u[p^\infty]}$  is split, since  $\text{Ext}_R(\mathbb{Q}_p/\mathbb{Z}_p, \mu_{p^\infty})$  is killed by  $p^n$ . We are then reduced to checking that the homomorphism

$$1 + \mathfrak{m} \longrightarrow \text{Ext}_R(\mathbb{Q}_p/\mathbb{Z}_p, \mu_{p^\infty}), \quad u \longmapsto \eta_{M^u[p^\infty]},$$

is an isomorphism. But this homomorphism equals  $\Phi$  in (2.4), and the latter is an isomorphism by Lemma 2.3. ■

### 3 Auxiliary Results

#### 3.1 Galois Actions

Let  $k'/k$  be a finite Galois extension of  $\Gamma = \text{Gal}(k'/k)$  and set  $R' = R \otimes_{W(k)} W(k')$ . Note that  $R'$  is an artinian local ring with residue field  $k'$  and  $\text{Spec } R' \rightarrow \text{Spec } R$  is a Galois covering of group  $\Gamma$ . Then  $\Gamma$  naturally acts on  $\mathcal{M}_1(R')$  and  $\text{Def}(R', k')$  and the Serre–Tate functor  $\Delta_{R'}$  (1.1) commutes with the Galois action.

Note that the datum of a 1-motive  $M$  over  $R$  is equivalent to the datum of a 1-motive  $M' = [u': L' \rightarrow G']$  over  $R'$  with a  $\Gamma$ -action compatible with the  $\Gamma$ -action on  $\text{Spec } R'$ . Indeed,  $L'$  descends to a lattice over  $R$ . Further, since the topological space underlying  $G'$  coincides with the topological space underlying the semi-abelian  $k'$ -variety  $G'_0$ , it can be covered by affine open subschemes which are stable under the  $\Gamma$ -action and hence it descends to an  $R$ -group scheme. The maximal subtorus  $T'$  of  $G'$  descends to a subtorus  $T$  of  $G$  and  $G/T$  is an abelian scheme, since it is an abelian scheme after base change to  $R'$ . Similarly, the datum of an object  $(G_0, \mathcal{B}, \varepsilon_0)$  in  $\text{Def}(R, k)$  is equivalent to the datum of a  $(G'_0, \mathcal{B}', \varepsilon'_0)$  in  $\text{Def}(R', k')$  together with a  $\Gamma$ -action compatible with the  $\Gamma$ -action on  $k'$  (in the first and third entries) and the  $\Gamma$ -action on  $R'$  (on the second entry).

#### 3.2 Refinements

Recall that any 1-motive  $M = [u: L \rightarrow G]$  over  $R$  admits a so-called universal extension  $M^{\text{h}} = [u^{\text{h}}: L \rightarrow G^{\text{h}}]$  that fits into a short exact sequence of complexes of  $R$ -group schemes

$$(3.1) \quad 0 \longrightarrow [0 \longrightarrow \mathbb{V}(M)] \longrightarrow M^{\text{h}} \longrightarrow M \longrightarrow 0,$$

where  $\mathbb{V}(M)$  is the vector group associated to the dual sheaf of  $\text{Ext}_{\text{Zar}}^1(M, \mathbb{G}_{a,R})$ . Note that  $\mathbb{V}(M)$  is killed by  $p^s$ , since the multiplication by  $p^s$  morphism is the 0 map on  $\mathbb{G}_{a,R}$ . Note that  $M^{\text{h}}$  is denoted by  $\mathbb{E}(M) = [L \rightarrow \mathbb{E}(M)_G]$  in [1, 2].

**Remark 3.1** We can refine some preliminary results in [6]. Let  $s' \geq s$  be the minimal integer such that  $\mathfrak{m}^{s'} = 0$ .

(i) Let  $G$  be any smooth commutative  $R$ -group scheme. By [6, Lemma 1.1.2] the kernel of the reduction map  $G(R) \rightarrow G(k)$  is killed by  $p^{s(s'-1)}$ . In our hypothesis one can prove that it is killed by  $p^{s'-1}$ . Indeed, by the theory of Greenberg functor the sections  $G(R/\mathfrak{m}^i)$ ,  $1 \leq i \leq s'$ , can be identified with the group of  $k$ -rational sections of a smooth  $k$ -group scheme  $\text{Gr}_i(G)$ . Further, there are so-called change of level morphisms  $\rho_i^1: \text{Gr}_{i+1}(G) \rightarrow \text{Gr}_i(G)$  such that  $\rho_i^1(k)$  coincides with the reduction map  $G(R/\mathfrak{m}^{i+1}) \rightarrow G(R/\mathfrak{m}^i)$ . By Greenberg's structure theorem ([5],[3, Thm. A.17]) the kernel of  $\rho_i^1$  is a  $k$ -vector group, thus killed by  $p$ . It follows then by induction that  $\text{Ker}(G(R) \rightarrow G(k))$  is killed by  $p^{s'-1}$ .

(ii) By [6, Lemma 1.1.3 (3)] a morphism of semi-abelian varieties  $\varphi_0: G_0 \rightarrow H_0$  lifts over  $R$  up to multiplication by  $p^{s(s'-1)}$ . We can also improve this estimate if  $p > 2$ . Let  $M, N$  be 1-motives over  $R$ . By [1, Thm. 2.1] there exists a canonical morphism between universal extensions (3.1)  $\varphi^{\text{h}}: M^{\text{h}} \rightarrow N^{\text{h}}$  that lifts  $\varphi_0^{\text{h}}$ . If  $\varphi^{\text{h}}(\mathbb{V}(M)) \subseteq \mathbb{V}(N)$ , then  $\varphi^{\text{h}}$  induces a morphism  $\varphi: M \rightarrow N$  that lifts  $\varphi_0$ . In general, since the multiplication by  $p^s$  morphism kills  $\mathbb{V}(M)$ , the morphism  $p^s \varphi^{\text{h}}$  maps  $\mathbb{V}(M)$  to 0. Hence,  $p^s \varphi_0$  lifts to a morphism " $p^s \varphi$ ":  $M \rightarrow N$ .

## 4 Proof of the Main Theorem

### 4.1 Full Faithfulness

**Proposition 4.1** *Let  $M, N$  be two 1-motives over  $R$ . Then the reduction map*

$$\text{Hom}_{\mathcal{M}_1(R)}(M, N) \longrightarrow \text{Hom}_{\mathcal{M}_1(k)}(M_0, N_0)$$

*is injective.*

**Proof** Consider a morphism  $\varphi: M \rightarrow N$  and assume that its reduction modulo  $\mathfrak{m}$  is the 0 morphism. If  $p > 2$ , then  $\varphi = 0$ , since the induced morphism between universal extensions  $\varphi^{\text{h}}: M^{\text{h}} \rightarrow N^{\text{h}}$  is the zero map [1, Thm. 2.1]. In general, one has  $\varphi = 0$  in degree  $-1$  by the equivalence of the étale sites over  $R$  and over  $k$ , and  $\varphi = 0$  in degree 0 by [6, Lemma 1.1.3 2)]. ■

**Corollary 4.2** *The functor (1.1) is faithful.*

**Proposition 4.3** *The functor (1.1) is full.*

**Proof** Let  $M = [u: L \rightarrow G], N = [v: F \rightarrow H]$  be two 1-motives over  $R$ ,  $\varphi_0: M_0 \rightarrow N_0$  a morphism between their reduction modulo  $\mathfrak{m}$ , and  $\psi: M[p^\infty] \rightarrow N[p^\infty]$  a lifting of  $\varphi_0[p^\infty]: M_0[p^\infty] \rightarrow N_0[p^\infty]$ . We have to prove that there exists a lifting  $\varphi: M \rightarrow N$  of  $\varphi_0$  over  $R$  such that  $\varphi[p^\infty] = \psi$ .

As a first step, consider the case  $L = F = 0$ . We assume that  $p > 2$ ; if  $p = 2$ , the same proof works replacing  $s$  with  $s(s' - 1)$ , where  $m^{s'} = 0$ , and it coincides with the one in [6, p. 144]. Given a  $\varphi_0: G_0 \rightarrow H_0$  and a morphism  $\psi: G[p^\infty] \rightarrow H[p^\infty]$  lifting  $\varphi_0[p^\infty]$ , by Remark 3.1(ii) there exists a morphism “ $p^s \varphi$ ”:  $G \rightarrow H$  lifting  $p^s \varphi_0$ . Further,  $p^s \psi =$  “ $p^s \varphi$ ” $[p^\infty]$  by [6, Lemma 1.1.3(2)], since both morphisms lift  $p^s \varphi_0[p^\infty]$ . Hence, “ $p^s \varphi$ ” kills  $G[p^s]$  and thus “ $p^s \varphi$ ” =  $p^s \varphi$  for a morphism  $\varphi: G \rightarrow H$  (necessarily unique since  $\text{Hom}_{R\text{-gr}}(G, H)$  has no  $p$ -torsion). Thus, the restriction of the functor (1.1) to  $\mathcal{M}_1(R)_{\leq -1}$  is full.

As a second step, consider the case where  $G$  and  $H$  are abelian varieties. Fullness of the restriction of the functor (1.1) to  $\mathcal{M}_1(R)_{\geq -1}$  follows from the previous step via Cartier duality.

For the general case, let  $M = [u: L \rightarrow G], N = [v: F \rightarrow H]$  be two 1-motives over  $R$ ,  $\varphi_0: M_0 \rightarrow N_0$  a morphism between their reduction modulo  $\mathfrak{m}$  and  $\psi: M[p^\infty] \rightarrow N[p^\infty]$  a lifting of  $\varphi_0[p^\infty]: M_0[p^\infty] \rightarrow N_0[p^\infty]$ . Let  $\varphi_0 = (f_0, g_0)$ , i.e.,  $f_0: L_0 \rightarrow F_0, g_0: G_0 \rightarrow H_0$  and  $g_0 \circ u_0 = v_0 \circ f_0$ . Recall that any lifting  $\psi$  of  $\varphi_0[p^\infty]$  respects weight filtrations by Lemma 2.2. Hence, by the first step of the proof there exists a unique morphism  $g: G \rightarrow H$  lifting  $g_0$ . Further, by the equivalence between the category of étale group schemes over  $k$  and the category of étale group schemes over  $R$ , there exists a unique  $f: L \rightarrow F$  lifting  $f_0$ . We are left to prove that  $g \circ u = v \circ f$ , so that  $\varphi = (f, g)$  is a morphism of 1-motives, and that  $\varphi[p^\infty] = \psi$ . The latter equality follows from [6, Lemma 1.1.3 2)], since both morphisms are liftings of  $\varphi_0[p^\infty]$ .

Let  $Z = [w: L \rightarrow H]$  with  $w = g \circ u - v \circ f$ . We claim that  $\eta_{Z[p^\infty]}$  in (2.2) is split. For proving this claim, it is sufficient to check that  $\eta_{Z[p^r]}$  (and hence  $\tilde{\eta}_{Z[p^r]}$  in (2.1)) is split for any  $r$ . Since  $\psi[p^r]$  restricts to the morphism  $g[p^r]: G[p^r] \rightarrow H[p^r]$  on weight  $\leq -1$  subgroups and induces  $f/p^r f: L/p^r L \rightarrow F/p^r F$  in weight 0, it is

$$(f/p^r f)^* \eta_{N[p^r]} = g[p^r]_* \eta_{M[p^r]},$$

and hence

$$f^* \tilde{\eta}_{N[p^r]} = g[p^r]_* \tilde{\eta}_{M[p^r]}.$$

By Remark 2.1, we have

$$\begin{aligned} \tilde{\eta}_{Z[p^r]} &= (-w)^* \xi_{H[p^r]} = f^*(v^* \xi_{H[p^r]}) - u^*(g^* \xi_{H[p^r]}) = f^* \tilde{\eta}_{N[p^r]} - u^* g[p^r]_* \xi_{G[p^r]} \\ &= f^* \tilde{\eta}_{N[p^r]} - g[p^r]_* \tilde{\eta}_{M[p^r]} = 0. \end{aligned}$$

Hence,  $Z$  is a 1-motive such that  $\eta_{Z[p^\infty]}$  is split extension of  $L[p^\infty]$  by  $H[p^\infty]$  and  $w_0 = 0$ . We are left to prove that  $w = 0$ . We can work étale locally on  $R$  and then assume that  $L$  is constant and the maximal subtorus  $S$  of  $H$  is split. By the second step of this proof,  $w_{\text{ab}}: L \rightarrow B$  is the 0 morphism. Hence,  $w$  factors through  $S$ . Let  $M_t = [w: L \rightarrow S]$  and note that  $M_t[p^\infty]$  is split extension of  $L[p^\infty]$  by  $S[p^\infty]$ . Hence,  $w = 0$  by Lemma 2.4. ■

### 4.2 Essential Surjectivity

The strategy of the proof of essential surjectivity is first to construct the desired 1-motive étale locally and then to apply descent. We then need the following result.

**Lemma 4.4** *Let  $k'/k$  be a finite Galois extension of Galois group  $\Gamma$  and set  $R' = R \otimes_{W(k)} W(k')$ . A 1-motive  $M'$  over  $R'$  descends to  $R$  if and only if its image in  $\text{Def}(R', k')$  via (1.1) descends to  $\text{Def}(R, k)$ .*

**Proof** The necessity is clear. For sufficiency, let  $M' = [u': L' \rightarrow G']$  and assume that  $(M'_0, \mathcal{B}' = M'[p^\infty], \text{can}: \mathcal{B}'_0 \xrightarrow{\sim} M'_0[p^\infty])$  descends to an object

$$(M_0 = [L_0 \rightarrow G_0], \mathcal{B}, \varepsilon_0: \mathcal{B}_0 \xrightarrow{\sim} M_0[p^\infty])$$

in  $\text{Def}(R, k)$ . For any  $\sigma \in \Gamma$  we then have an isomorphism in  $\text{Def}(R', k')$

$$(\varphi_{\sigma,0}, \psi_\sigma): (M'_0, \mathcal{B}', \text{can}) \xrightarrow{\sim} (\sigma^* M'_0, \sigma^* \mathcal{B}', \sigma^* \text{can}),$$

where

$$\varphi_{\sigma,0}: M'_0 \xrightarrow{\sim} \sigma^* M'_0, \quad \psi_\sigma: \mathcal{B}' \xrightarrow{\sim} \sigma^* \mathcal{B}'$$

make the diagram

$$\begin{array}{ccc} \mathcal{B}'_0 & \xrightarrow{\psi_{\sigma,0}} & \sigma^* \mathcal{B}'_0 \\ \text{can} \downarrow & & \downarrow \sigma^* \text{can} \\ M'_0[p^\infty] & \xrightarrow{\varphi_{\sigma,0}[p^\infty]} & \sigma^* M'_0[p^\infty] \end{array}$$

commute. By the full faithfulness of (1.1) proved in Corollary 4.2 and Proposition 4.3, the isomorphism  $(\varphi_{\sigma,0}, \psi_\sigma)$  gives a unique isomorphism  $\varphi_\sigma: M' \xrightarrow{\sim} \sigma^* M'$  that lifts  $\varphi_{\sigma,0}$  and restricts to  $\psi_\sigma$  on BT groups. Hence, we have defined an action of  $\Gamma$  on  $M'$  compatible with the  $\Gamma$ -action on  $R'$  and thus  $M'$  descends over  $R$ , as explained in Section 3.1. ■

**Proposition 4.5** *The functor (1.1) is essentially surjective.*

**Proof** Let  $(M_0 = [L_0 \rightarrow G_0], \mathcal{B}, \varepsilon_0)$  be an object of  $\text{Def}(R, k)$ .

As a first step, consider the case where  $L_0 = 0$ . Thanks to Lemma 4.4 we can assume that the maximal subtorus  $T_0$  of  $G_0$  is split of dimension  $d$ . The Cartier dual of  $G_0$  is a 1-motive  $[w_0: \mathbb{Z}^d \rightarrow A_0^*]$ , where  $A_0^*$  is the dual of the abelian quotient  $A_0$  of  $G_0$ . The abelian variety  $A_0$  lifts to an abelian scheme  $A$  over  $R$ , and since the reduction map  $A(R) \rightarrow A(k)$  is surjective, the morphism  $w_0$  lifts to a morphism  $w: \mathbb{Z}^d \rightarrow A^*$ . Passing to Cartier duals, we obtain an  $R$ -group scheme  $G$  that is an extension of  $A$  by  $\mathbb{G}_{m,R}^d$  and lifts  $G_0$ . Now, the BT group  $G[p^\infty]$  might not be isomorphic to  $\mathcal{B}$ . Repeating word by word the proof of the classical Serre-Tate theorem [6, Thm. 1.2.1, pp. 145–146], one finds a finite flat subgroup scheme  $K$  of  $G[p^{2^s}]$  such that  $G/K$  is a lifting of  $G_0$  with BT group isomorphic to  $\mathcal{B}$ .

The case when  $M_0 = M_{0,\text{ab}}$  follows immediately by Cartier duality from the previous step.

For the general case, we can assume that  $L_0$  is constant and the maximal torus in  $G_0$  is split, again by Lemma 4.4. Recall that by Lemma 2.2 the BT group  $\mathcal{B}$  is naturally filtered so that  $W_{-1}\mathcal{B}$  is a lifting of  $G_0[p^\infty]$  and  $\mathcal{B}/W_{-2}\mathcal{B}$  is a lifting of  $M_{0,\text{ab}}[p^\infty]$ . By the previous steps we know that  $G_0$  lifts to an  $R$ -scheme  $G$ , which is an extension of an abelian scheme  $A$  by  $\mathbb{G}_{m,R}^d$ , and  $G[p^\infty] = W_{-1}\mathcal{B}$ ; further  $M_{0,\text{ab}}$  lifts to a 1-motive  $M_A = [u_A: \mathbb{Z}^m \rightarrow A]$  whose BT group is isomorphic to  $\mathcal{B}/W_{-2}\mathcal{B}$ .



Let  $M' = [u': \mathbb{Z}^m \rightarrow G]$  be any extension of  $M_A$  by  $T$ ; it exists, since  $H^1(R, \mathbb{G}_{m,R}) = 0$ . Since  $T(R) \rightarrow T(k)$  is surjective, we can assume that  $u'$  is also a lifting of  $u_0$ . We are then left to alter  $u'$  so that  $M'[p^\infty] \simeq \mathcal{B}[p^\infty]$  in  $\text{Ext}_R((\mathbb{Q}_p/\mathbb{Z}_p)^m, G[p^\infty])$ .

Let  $\mathcal{E} = \mathcal{B}[p^\infty] - M'[p^\infty]$ , as extension of  $(\mathbb{Q}_p/\mathbb{Z}_p)^m$  by  $G[p^\infty]$ . Since the push-out along  $G[p^\infty] \rightarrow A[p^\infty]$  maps  $\mathcal{E}$  to the trivial extension of  $(\mathbb{Q}_p/\mathbb{Z}_p)^m$  by  $A[p^\infty]$ , there exists by Lemma 2.4 a 1-motive  $N = [v: \mathbb{Z}^m \rightarrow T]$  such that  $v_0 = 0$  and the push-out of  $N[p^\infty]$  along  $T[p^\infty] \rightarrow G[p^\infty]$  is  $\mathcal{E}$ . Then  $M = [u = u' + v: \mathbb{Z}^m \rightarrow G]$  is a lifting of  $M_0$  and  $M[p^\infty]$  is isomorphic to  $\mathcal{B}[p^\infty]$ . ■

With this proposition, the proof of Theorem 1.1 is completed.

**Remark 4.6** Note that some results on deformations of 1-motives (but not Theorem 1.1) were proved with other methods in Madapusi's thesis [7]; however, they are not included in the preprint written by the author under the name K. Madapusi Pera and bearing the same title of the thesis.

Since any extension of an étale BT group by a toroidal BT group is split over  $k$ , we deduce from Theorem 1.1 the following generalization of Lemma 2.4.

**Corollary 4.7** *Suppose  $T$  is an  $R$ -torus and  $L$  is a lattice. Given a 1-motive  $M_0 = [u_0: L_0 \rightarrow T_0]$  and any BT group  $\mathcal{B}$  that is an extension of  $L[p^\infty]$  by  $T[p^\infty]$  there is a unique 1-motive  $M = [u: L \rightarrow T]$  that lifts  $M_0$  and whose BT group is isomorphic to  $\mathcal{B}$ .*

### 4.3 Formal Moduli and Serre–Tate Coordinates

Let  $k$  be an algebraically closed field of characteristic  $p > 0$ . We say that a 1-motive  $M_0$  over  $k$  is ordinary if  $\text{gr}_{-1}M_0 = A_0$  is an ordinary abelian variety (possibly trivial). Thanks to Theorem 1.1 one can easily extend Serre–Tate theory for ordinary abelian varieties to ordinary 1-motives. In this section, we will illustrate some results in this direction. Proofs are only sketched, since no new phenomena appear.

Following [6, § 2] one can define the formal moduli of a given 1-motive  $M_0$  over  $k$ . Namely, let  $\widehat{\mathcal{M}}_{M_0} = \widehat{\mathcal{M}}$  be the functor

$$\widehat{\mathcal{M}}(R) := \{R\text{-liftings of } M_0\}/\text{iso},$$

where  $R$  is an artinian local ring with residue field  $k$ . By Theorem 1.1 we get a bijection

$$\widehat{\mathcal{M}}(R) = \{R\text{-liftings of } M_0[p^\infty]\}/\text{iso}.$$

**Theorem 4.8** *Let  $M_0, N_0$  be ordinary 1-motives over  $k$ .*

- (i) *Let  $R$  be an artinian local ring as above. For any 1-motive  $M$  over  $R$  lifting  $M_0$ , there exists a canonical  $\mathbb{Z}_p$ -bilinear form (the Serre–Tate coordinates)*

$$q(M/R, \cdot, \cdot): T_p M_0(k) \otimes T_p M_0^*(k) \longrightarrow \widehat{\mathbb{G}}_m(R),$$

where  $M_0^*$  denotes the Cartier dual of  $M_0$  and  $T_p M_0(k) := \varprojlim_n M_0[p^n](k)$ . It induces an isomorphism of functors

$$\widehat{\mathcal{M}}(\cdot) \simeq \text{Hom}_{\mathbb{Z}_p}(T_p M_0(k) \otimes T_p M_0^*(k), \widehat{\mathbb{G}}_m(\cdot)).$$

- (ii) Let  $\varphi_0: M_0 \rightarrow N_0$  be a morphism of 1-motives and let  $M, N$  be liftings over  $R$  of  $M_0, N_0$ , respectively. Then  $\varphi_0$  lifts to an  $R$ -morphism  $\varphi: M \rightarrow N$  if and only if

$$q(M/R, \alpha, \varphi_0^*(\beta)) = q(N/R, \varphi_0(\alpha), \beta)$$

for every  $\alpha \in T_p M_0(k)$  and  $\beta \in T_p M_0^*(k)$ . Further, if a lifting exists, it is unique.

**Proof** The proof goes exactly as in the classical case. For the convenience of the reader we point out the main steps (cf. [6, p. 152]).

- Since  $k$  is algebraically closed and  $M_0$  is ordinary, the BT groups  $M_0[p^\infty]^\circ$  is split multiplicative and  $M_0[p^\infty]^{\acute{e}t}$  is split étale. Hence, the same are true for  $M[p^\infty]^\circ$  and  $M[p^\infty]^{\acute{e}t}$  for any lifting  $M$  of  $M_0$  over  $R$ . For this reason we may write  $M_0[p^\infty]^\circ$  (resp.  $M_0[p^\infty]^{\acute{e}t}$ ) for the corresponding functor on the category of artinian local rings with residue field  $k$ .
- By Cartier duality, there is a perfect pairing  $M_0[p^n] \times M_0^*[p^n] \rightarrow \mu_{p^n}$  [4, 10.2.5], inducing an isomorphism of functors

$$M_0[p^\infty]^\circ \xrightarrow{\sim} \text{Hom}_{\mathbb{Z}_p}(T_p M_0^*(k), \widehat{\mathbb{G}}_m),$$

on the category of artinian local rings with residue field  $k$ . We denote the corresponding pairing by  $E_M: M_0[p^\infty]^\circ \times T_p M_0^*(k) \rightarrow \widehat{\mathbb{G}}_m$ .

- By [8, Proposition 2.5 p. 180] and the previous steps, for any  $R$  as above and any lifting  $M$  of  $M_0$  over  $R$ , we have isomorphisms

$$\begin{aligned} \text{Ext}_R(M[p^\infty]^{\acute{e}t}, M[p^\infty]^\circ) &= \text{Ext}_R(T_p M_0(k) \otimes \mathbb{Q}_p/\mathbb{Z}_p, M[p^\infty]^\circ) \\ &\simeq \text{Hom}(T_p M_0(k), M[p^\infty]^\circ(R)); \end{aligned}$$

thus, there exists a unique  $\phi_M \in \text{Hom}_R(T_p M_0(k), M[p^\infty]^\circ(R))$  associated with the class of the BT group  $M[p^\infty]$ , viewed as extension of  $M[p^\infty]^{\acute{e}t}$  by  $M[p^\infty]^\circ$ .

Then we can define  $q(M/R, \alpha, \beta) := E_M(\phi_M(\alpha), \beta)$  in  $\widehat{\mathbb{G}}_m(R)$  and follow word by word the proof in [6]. ■

**Remark 4.9** Let  $M_0$  be an ordinary 1-motive over  $K$  and denote its graded quotients (by the weight filtration) by  $L_0, A_0$  and  $T_0$ . Fix basis:  $\alpha_1, \dots, \alpha_g$  of  $T_p A_0(k)$  with  $g = \dim A_0$ ;  $\alpha_{g+1}, \dots, \alpha_{g+\ell}$  of  $T_p L_0(k)$  with  $\ell = \text{rank } L_0$ ;  $\beta_1, \dots, \beta_g$  of  $T_p A_0^*(k)$ ;  $\beta_{g+1}, \dots, \beta_{g+t}$  of  $T_p T_0^*(k)$  with  $t = \dim T_0$ . Then the coordinate ring of  $\widehat{\mathcal{M}}$  is isomorphic to

$$W(k)[[T_{i,j}]] \quad \text{where } T_{i,j} = q(\alpha_i, \beta_j) - 1,$$

where  $q(\cdot, \cdot)$  is the bilinear form associated with the universal formal deformation. In particular  $\widehat{\mathcal{M}}$  is represented by a  $(g + \ell) \times (g + t)$ -dimensional formal torus.

**Proposition 4.10** Let  $M_0$  be an ordinary 1-motive over  $k$ . Then there exists a 1-motive  $M$  over  $W(k)$  lifting  $M_0$  such that  $\text{End}_{\mathcal{M}_1(W(k))}(M) \rightarrow \text{End}_{\mathcal{M}_1(k)}(M_0)$  is bijective.

**Proof** By Theorem 1.1 there exists a unique 1-motive  $M_n = [L_n \rightarrow G_n]$  over  $W_{n+1}(k)$  lifting  $M_0$  such that  $M_n[p^\infty]$  has split connected-étale sequence. As  $n$  goes to infinity, these liftings form a compatible system. In order to see that the limit is algebraizable it

is enough to check that  $\varinjlim_n G_n$  is algebraizable, since all  $L_n$  are torsion-free constant abelian groups of the same rank, and for any group scheme  $J$  of finite type over  $W(k)$

$$(4.1) \quad J(W(k)) = \varprojlim_m J(W_m(k)).$$

Now, by Cartier duality, the system  $(G_n)_n$  corresponds to a compatible system of 1-motives  $[T_n^* \rightarrow A_n^*]$ . The latter is algebraizable if the system  $(A_n^*)_n$  is. Since  $M_n[p^\infty]$  has split connected-étale sequence, the same are true for  $A_n[p^\infty]$  and  $A_n^*[p^\infty]$ . So  $(A_n^*)_n$  is algebraizable by the proof of [8, Ch. 5, Thm 3.3, p. 173].

Let us denote by  $M = [u: L \rightarrow G]$  (resp.  $A$ ) the lift of  $M_0$  (resp  $A_0$ ) over  $W(k)$  constructed in the previous paragraph; then  $G$  is extension of  $A$  by the unique torus  $T$  lifting  $T_0$ . By construction and Theorem 1.1, any reduction map  $\text{End}_{\mathcal{M}_1(W_{n+1}(k))}(M_n) \rightarrow \text{End}_{\mathcal{M}_1(W(k))}(M_0)$  is bijective. We are left to check that the map  $\text{End}_{\mathcal{M}_1(W(k))}(M) \rightarrow \varprojlim \text{End}_{\mathcal{M}_1(W_{n+1}(k))}(M_n) = \text{End}_{\mathcal{M}_1(W(k))}(M_0)$  is bijective. Let  $\varphi = (f, g)$  be an endomorphism of  $M$ , i.e., we have morphisms  $f: L \rightarrow L, g: G \rightarrow G$  such that  $u \circ f = g \circ u$ . If  $\varphi_0 = 0$ , then  $\varphi = 0$  by devissage, since

$$(4.2) \quad \text{End}_{W(k)}(L) = \text{End}_k(L_0), \quad \text{End}_{W(k)}(T) = \text{End}_k(T_0),$$

and  $\text{End}_{W(k)}(A) = \text{End}_k(A_0)$  by [8, Ch. 5, Thm 3.3]. For surjectivity, let  $\varphi_n = (f_n, g_n): M_n \rightarrow M_n$  be a compatible system of endomorphisms. By (4.2) and (4.1) it suffices to show the existence of  $g: G \rightarrow G$  lifting the morphisms  $(g_n)_n$ . We can work with the Cartier dual  $[T^* \rightarrow A^*]$  of  $G$  instead, and lift the Cartier duals of  $(g_n)_n$ ; as above, we can reduce to the case  $T^* = 0$  and then conclude by [8, Ch. 5, Thm 3.3]. ■

**Acknowledgements** This work was done while the second author was visiting the University of Padua supported by a “délégation CNRS”. The first author thanks the project PRIN 2015 “Number Theory and Arithmetic Geometry” for financial support. Both authors thank the referee, whose detailed comments and helpful suggestions improved the exposition of this paper.

## References

- [1] F. Andreatta and A. Bertapelle, *Universal extension crystals of 1-motives and applications*. J. Pure Appl. Algebra 215(2011), no. 8, 1919–1944. <http://dx.doi.org/10.1016/j.jpaa.2010.11.004>
- [2] F. Andreatta and L. Barbieri-Viale, *Crystalline realizations of 1-motives*. Math. Ann. 331(2005), 111–172. <http://dx.doi.org/10.1007/s00208-004-0576-4>
- [3] A. Bertapelle and C. D. González-Avilés, *The Greenberg functor revisited*. Eur. J. Math. (2018). <http://dx.doi.org/10.1007/s40879-017-0210-0>
- [4] P. Deligne, *Théorie de Hodge. III*. Inst. Hautes Études Sci. Publ. Math. 44(1974), 5–77.
- [5] M. J. Greenberg, *Schemata over local rings. II*. Ann. of Math. 78(1963), 256–266. <http://dx.doi.org/10.2307/1970342>
- [6] N. Katz, *Serre–Tate local moduli*. In: Algebraic surfaces (Orsay, 1976–78), Lecture Notes in Math., 868, Springer, Berlin–New York, 1981, pp. 138–202.
- [7] K. S. Madapusi Sampath, *Toroidal compactifications of integral models of Shimura varieties of Hodge type*. PhD thesis, Chicago, 2011.
- [8] W. Messing, *The crystals associated to Barsotti–Tate groups with applications to abelian schemes*. Lecture Notes in Mathematics, 264, Springer-Verlag, Berlin–New York, 1972.

*Dipartimento di Matematica "Tullio Levi-Civita", Università degli Studi di Padova, via Trieste, 63, I-35121 Padova, Italy*

*e-mail:* [alessandra.bertapelle@unipd.it](mailto:alessandra.bertapelle@unipd.it)

*Institut de Mathématiques de Bordeaux, University of Bordeaux, F-33405 Talence cedex, France*

*e-mail:* [nicola.mazzari@math.u-bordeaux.fr](mailto:nicola.mazzari@math.u-bordeaux.fr)