

Methods: This patient will be presented, along with systematic bibliography review of the topic.

Results: The following results were extracted upon the attention given to this patient which was admitted to the Psychiatric Unit. First of all, the mouth stitches were removed, along with a petition for toxicological analysis. The results gave positive for cannabis and benzodiazepines. The patient was also brought previously this year with another autolytic attempt, this time on cocaine consumption too. Furthermore, a thorough review was made of the other autolytic attempts, including those which happened in his country of origin. The patient has hundreds of small cuts among his arms, from previous cuts made in the past. Furthermore, subcutaneous wounds were auto inflicted in the ER, with a small blade.

Among the whole interview, it was clear he had a personality disorder, with high impulsivity levels and lack of control once the situation overflows.

We also tried to understand the outcome of suturing his mouth. The patient referred his acts of impulsiveness due to his overwhelming situation of both having no job at this moment and the pain he was suffering due to his ankle procedure.

The patient was admitted to our Unit due to the high risk he could repeat this act. Upon arrival, the same day he was admitted, the patient asked if he had to stay at the unit. When explaining the following already told event, furthermore insisting in the possibility of been evaluated by the Traumatology team, he proceeded to try and hang himself with his medical-hospital clothing.

The patient was treated with antipsychotics. Along with Lormetazepam at night. At the end of the hospitalization, and after been evaluated by the Psychiatrist of this Unit, the patient was also treated with Lithium due to its effectiveness in the treatment of autolytic attempts.

Conclusions: Personality disorders are one of the psychiatric pathologies that prevail with greater frequency in autolytic attempts ¹. Additionally, it should be taken into account the possible ongoing consumption of psychoactive drugs that could also derive in psychopathological decompensation. On top of the following, the use of antipsychotic treatment is indicated for the managing of conduction altercations ², besides Lithium being a great option in managing suicidal temptations ³.

Disclosure of Interest: None Declared

EPV0716

Personality disorders and addiction A study of 54 patients

Y. Bensalah^{1*}, L. Tbatou¹, M. Sabir¹ and F. Elomari¹

¹Psychiatric hospital Ar Razi, salé, Morocco

*Corresponding author.

doi: 10.1192/j.eurpsy.2024.1359

Introduction: Personality disorders are very often comorbid with addictions and are known to have a negative impact on the development of substance use disorder.

Objectives: Evaluate the prevalence of personality disorder in patients with problematic use of psychoactive substances followed at Ar Razi hospital Determine the relationship between different personality disorders and the clinical aspects of psychoactive substance use

Methods: This is a descriptive and analytical cross-sectional study carried out among 54 patients followed at Ar-Razi hospital Salé in Morocco for problematic use of psychoactive substances from June 1 to September 15, 2023 (Diagnosis assessed by DSM 5) Data collection was done using a questionnaire including clinical and socio-demographic characteristics and data on addiction to psychoactive and behavioral substances. The psychometric scale used to assess personality disorder: Personality Disorder Questionnaire (PDQ-4+)

Results: We recruited 54 patients with age ranges from 18 to 45 years, with a male predominance. The average age at the start of psychoactive substance use was 15 years. Tobacco is the most used substance followed by cannabis Antisocial, histrionic and borderline personality disorders were the most common in our population. There was a statistically significant difference between specific personality disorder and the presence of severe psychoactive substance use disorder

Conclusions: The frequency of personality disorders is high among subjects with problematic use of psychoactive substances. It is necessary to take care of them simultaneously (integrated care) in order to improve the prognosis

Disclosure of Interest: None Declared

EPV0718

Avoidant personality disorder through the lens of ICD 11

A. Bogdanovska Toskic^{1*} and S. Arsova Hadzi-Angjelkovska²

¹Psychiatry, General Hospital Kumanovo, Kumanovo and ²Psychiatry, University Clinic of Psychiatry, Skopje, North Macedonia

*Corresponding author.

doi: 10.1192/j.eurpsy.2024.1360

Introduction: With the new dimensional diagnosis of personality disorders in ICD 11, the categorical model has been abandoned. The types of personality disorders in the new dimensional model should show certain common characteristics. According to the recognition of the common characteristics of individual types of personality disorders, as well as determining the severity, a transition from the categorical to the dimensional diagnostic system can be made.

Objectives: To analyze and present the trait domains specifiers in persons with avoidant personality disorder and to facilitate the adoption of the new diagnostic criteria.

Methods: An unsystematized literature review was made, with key words: avoidant personality disorder, ICD 11, ICD 10, traits; and a case was presented.

Results: This is the case of a 26-year-old student who has had no friends since his school days. During his secondary education, on the initiative of another person, he got together with several other people, but he was not fully accepted. During the studies, the communication with the colleagues took place only at the university and around the responsibilities. About a year ago, he had reduced willpower and suicidal thoughts, when he took antidepressant and adjuvant anti-psychotic therapy for some time. He is now being examined due to severe tension, dissatisfaction, lack of friends, repeated suicidal ideation. According to researches, people with avoidant disorder have prominent trait domains – negative affectivity, detachment and