

γ -ADMISSIBILITY IN FIRST-ORDER RELEVANT LOGICS: PROOF USING NORMAL MODELS IN THE MARES–GOLDBLATT SETTING

NICHOLAS FERENZ 

Center of Philosophy, School of Arts and Humanities, University of Lisbon
and

THOMAS MACAULAY FERGUSON 

Department of Cognitive Science, Rensselaer Polytechnic Institute

Abstract. For relevant logics, the admissibility of the rule of proof γ has played a significant historical role in the development of relevant logics. For first-order logics, however, there have been only a handful of γ -admissibility proofs for a select few logics. Here we show that, for each logic $L^{\circ t}$ of a wide range of propositional relevant logics for which excluded middle is valid (with fusion and the Ackermann truth constant), the first-order extensions $QL^{\circ t}$ and $LQ^{\circ t}$ admit γ . Specifically, these are particular “conventionally normal” extensions of the logic $G^{g,d}$, which is the least propositional relevant logic (with the usual relational semantics) that admits γ by the method of normal models. We also note the circumstances in which our results apply to logics without fusion and the Ackermann truth constant.

§1. Introduction. The admissibility of γ in relevant logics remains an important question. With recent interest in quantified (modal) relevant logics, attention is turned to γ -admissibility in these logics. A particular boon in this research is the semantic framework for the quantified relevant logics **QR** and **RQ** introduced by Mares & Goldblatt [16], which has been extended more generally to first-order and first-order modal relevant logics sound and complete for ternary relational frames by Ferenz [8].¹ Here we employ this Mares–Goldblatt semantics as generalized by Ferenz to investigate which first-order relevant logics admit γ using the method of *normal models*, which was introduced by Sylvan (né Routley) and Meyer [25]. This provides a foundation to explore γ -admissibility in first-order modal relevant logics, taking advantage of the general framework provided by the Mares–Goldblatt interpretation of quantifiers.

Ackermann’s rule γ , originally given in [1], is a form of *disjunctive syllogism* (or, classically, *detachment/modus ponens*). In the tradition of relevant logic, γ ought to be rejected when formulated as a rule under which theories are closed. This would entail that from inconsistent theories anything is derivable, which is anathema to the

Received: July 4, 2023.

2020 *Mathematics Subject Classification*: Primary 03B47, 03Bxx, 03Cxx.

Key words and phrases: relevant logic, gamma admissibility, first-order relevant logic.

¹ Additionally, Tedder & Ferenz [34] have extended this framework to logics sound and complete for neighbourhood semantics, and the framework has been applied to conditional logics extending **FDE** [7].



relevantist project. However, it is another story if we take γ to close the set of theorems. Thus, γ is presented follows:

$$\neg A \vee B, A \Rightarrow B.$$

The symbol ' \Rightarrow ' here indicates a *rule of proof* in the sense of Humberstone [14]; Smiley [33]: That is, $A_1, \dots, A_n \Rightarrow B$ means that if each A_i is a theorem of the logic in question, then so is B . The rule γ is typically not given in the definition of a relevant logic; it is, however, sometimes an admissible rule. That is, in particular, the addition of the rule does not result in new theorems. The question of which relevant logics admit γ is an interesting and significant question, and has been in the crosshairs of relevantists since the beginning.

We present logics as Hilbert-style axiom systems with axiom and rule schemes defining a set of theorems. We use the usual notion of a proof (of A) in this setting as a sequence of formulas (ending with A) where each formula is either an instance of an axiom scheme or follows from previous formulas by application of an instance of a rule scheme. As our interest is in the set of theorems and not a consequence relation, we set our focus on *rules of proof* (defined above). In defining a logic, the rules explicitly given in the statement of the logic we call the *primitive* rules of the systems. Similar to primitive rules are the *derived* rules, whose form can be captured exactly by a series of axioms and primitive rule applications. For example, a rule of reiteration of the form $A \Rightarrow A$, if not primitive, can be shown to be derivable using modus ponens and the axiom $A \rightarrow A$. Finally, a rule is *admissible* if there exists a proof of the conclusion of the rule whenever there are proofs of the premises.²

In this paper we will give all primitive rules using the \Rightarrow notation, but the distinction between primitive and admissible rules is worth noting. One way to expand relevant logics is by the addition of proper axioms, with the notion of derivation suitable altered to include the additional proper axioms. These need not be schematic, and can allow relevant logic to represent reasoning from theories such as set theory, Peano Arithmetic, or even just a small set of beliefs. Part and parcel to the relevant approach is that inconsistent theories do not imply every formula, and so γ is not desirable as a primitive rule in these cases. On the other hand, the rules taken as primitive, although presented as rules of proof, are taken to apply in these extended cases.³

Meyer et al. [21, p. 120] note, “the cut theorem...is for classical theories simply γ is peculiar notation.” This is emphasized in Urquhart [35], summarizing the history and importance of γ , where he suggests additional similarities. Notably he conjectures a speed-up theorem for γ is relevant logics, in analogy to speedup theorems for the rule *cut* in classical logics (as shown in, e.g., Pudlák [23]).⁴ The admissibility of γ ,

² A logic which has an admissible rule that is not derivable lacks the property of being *structurally complete* (see Raftery & Świrydowicz [24] for an exploration of structural completeness in some relevant logics).

³ Note that there is another way to extend relevant logics to deal with theories: taking theories to be sets of sentences closed under Modus Ponens, Adjunction, and provable implications. This approach highlights the focus on logics as sets of theorems, and the admissibility of γ on the set of theorems retains its importance.

⁴ A speedup theorem is essentially a complexity result on the size of proofs. For γ this would mean there are theorems that are relatively small using γ , but whose derivations without γ have a much larger lower bound on their size.

if Urquhart's conjecture is proven, has significant consequences for proof-theoretic presentations of first-order relevant logics.

While some logics admit γ , such as **R** and **E**, several do not. In particular, several contraction-less relevant logics extended by Boolean negation do not admit γ Meyer et al. [22]. The naïve set theory of Brady [3] also fails to admit γ , but in this case the failure of γ is a feature and not a bug; on the other hand, the proof in Friedman & Meyer [13] showing the failure of γ for Meyer's relevant arithmetic **R**[#] was a catastrophic event for the development of **R**[#]. Meyer & Dunn [20] first showed that **R**, **E**, and **T**—the favorite children of Anderson and Belnap—admit γ using algebraic techniques. Later, additional techniques were found, such as normal models [26], metavaluations [19], and reduced frames [32]. Although the technique of normal models is the basis of the techniques of this paper, metavaluations involve defining valuations mapping formulas to the values 0 or 1 based on the relationship they bear to a regular theory. For the interested reader, the method is elegantly described in [6]. Note that the normal models method is restricted by the requirement of the principle of excluded middle as an axiom. For relevant logics without this requirement, one turns to the other methods referenced in this paragraph.

For a detailed account of the history of γ in propositional relevant logics, the reader is directed to Urquhart [35] and the references therein. Many modal propositional logics have also been shown to admit γ ; e.g., see Mares & Meyer [17]; Routley & Meyer [25]; Seki [29–31]. The Mares–Goldblatt style semantics of Ferenz [8] for first-order modal relevant logics combines the Mares–Goldblatt machinery with the general frames of Seki [28], the latter of which is used to obtain γ -admissibility results for a wide class of modal relevant logics. Thus, Ferenz [8] has laid the groundwork for using the method of normal models in both the first-order and the first-order modal settings. Here we pursue the former.

For quantified relevant logics, as far as we know, only a handful have been shown to admit γ . The first, and most relevant to this paper, is the proof of γ -admissibility in **RQ** in [21, Theorem 6]. The method of proof is by an algebraic semantics for **RQ**. The Mares–Goldblatt interpretation of the quantifiers introduces a natural semantics for **RQ**, defined as a Hilbert style axiom system. The genesis of such a semantics was due to the fact that the most straightforward way of defining a constant domain, ternary relational semantics extending the semantics for **R** validated formulas which were not theorems of **RQ**. This is the incompleteness result of Fine [12].⁵ The set of validities of the class of the most straightforward constant domain, ternary relational models (semantically) determines a logic, and we will call this logic **RQ**. This properly contains the theorems of **RQ**, and was shown by Weiss [36] to admit γ . However, giving an axiomatization (finite or otherwise) of this set of validities is still an open question.

In Kripke [15], a method of proving γ -admissibility for first-order extensions of **R** and **E** using semantic tableaux is stated; however, no proof is given in detail. Kripke, however, fails to establish which of **RQ**/**RQ** and **EQ**/**EQ** his proof is applicable to.

The paper is divided as follows. We begin by introducing preliminaries such as the definitions of languages, logics, semantics, and key notions. Then we tackle proving γ -admissibility by generalizing the normal models method. Here we use the

⁵ Note that Fine [11] also gives an adequate semantics for **RQ**, but that many relevantists have nevertheless been searching for simpler, more natural semantics. The author claims that the Mares–Goldblatt semantics is exactly what was sought.

Mares–Goldblatt style semantics. Finally we make concluding remarks concerning future directions.

§2. Preliminaries. We jump straight into a presentation of first-order relevant logics. For the reader not familiar with propositional relevant logics, one may consult one of Dunn & Restall [6]; Bimbó [2] for an excellent overview. Some familiarity with propositional relevant logics and ternary relational semantics is assumed, but such knowledge is not required.

2.1. First-order relevant logics.

2.1.1. Language. A first-order language (with constants and without function symbols) is built up from a set of symbols divided as follows:

1. A denumerable set of variables $Var = \{x_0, x_1, \dots\}$. Here we assume a fixed but arbitrary ordering of the elements of Var , which is tracked by variable subscripts.
2. An at most denumerable *signature* \mathbb{S} consisting of
 - (a) a set of constant symbols $Con^{\mathbb{S}} = \{c_0, c_1, \dots\}$,
 - (b) a non-empty set of predicate symbols $Pred^{\mathbb{S}}$, where $P^n \in Pred^{\mathbb{S}}$ is an n -ary predicate. The set of n -ary predicates shall be written as $Pred^n \subseteq Pred^{\mathbb{S}}$,
3. A constant symbol \mathbf{t}
4. Binary operators $\wedge, \vee, \rightarrow, \circ$
5. Unary operator \neg
6. Quantifier symbols \forall, \exists .

The notion of being a *term*, relative to a signature, is defined as usual, and we will use τ with decorations varying over terms. A signature will henceforth be assumed fixed, and we will cease to mention signatures unless such a remark is required.⁶ Given a set U of individuals, a *variable assignment* is a denumerable sequence of individuals, $f \in U^\omega$, such that the n -th element in the sequence (written as fn) is the individual assigned to the n -th variable x_n given by the assumed fixed ordering. Given a variable assignment f , an x_n -variant of f differs from f in at most the assignment to the variable x_n . We write $f \sim_n f'$ (or $f \sim_{x_n} f'$) to denote that f and f' are x_n variants of one another. We will write $f[j/n]$ (or $f[j/x_n]$), with $j \in U$ to denote the result of replacing the n -th element of f with the individual j .

DEFINITION 2.1 (The First-Order Relevant Language). *The basic first-order relevant language \mathcal{L} , or well-formed formulas (hereby wff) is defined in Backus-Naur form as follows:*

$$\varphi ::= P^n(\tau_1, \dots, \tau_n) | \mathbf{t} | \neg\varphi | \varphi \wedge \varphi | \varphi \vee \varphi | \varphi \rightarrow \varphi | \varphi \circ \varphi | \forall x_n \varphi | \exists x_n \varphi.$$

Implicit is the use of parentheses around each construction with a binary connective. That is, we assume that all unary operators (including quantifiers) bind more strongly than binary operators. Moreover, we assume the right arrow binds weaker than fusion, which itself binds weaker than the extensional conjunction and disjunction.

⁶ In the canonical model constructions, the signature is assumed to contain sufficiently many constant symbols.

We write $\mathcal{A}[\tau/x]$ to denote the result of substituting every free occurrence of x in \mathcal{A} with the term τ . Similarly, we will use $\mathcal{A}[\tau_0/v_0, \dots, \tau_n/v_n]$ for the result of simultaneously replacing v_0 through v_n with τ_0 through τ_n respectively. The usual definitions of *bound* and *free* variables are assumed. A term τ is *free for* x (or *freely substitutable for* x) in \mathcal{A} if τ does not become bound in the resulting formula $\mathcal{A}[\tau/x]$.⁷

When we write a formula with a variable superscript, such as \mathcal{A}^x , this means that x does not occur free in \mathcal{A} .

2.1.2. Axiomatic presentations. Although no propositional language was defined, we first axiomatize a wide class of propositional relevant logics.⁸ Although alternative axiom systems can define several of the logics we present, a singular modular system extending a base logic is used.

DEFINITION 2.2 (Propositional Logics). *The base propositional logic $\mathbf{B}^{\circ t}$ is defined by the following axioms and rules:*⁹

- (ID) $\mathcal{A} \rightarrow \mathcal{A}$
- ($\wedge E$) $\mathcal{A} \wedge \mathcal{B} \rightarrow \mathcal{A}$
- ($\wedge E$) $\mathcal{A} \wedge \mathcal{B} \rightarrow \mathcal{B}$
- ($\vee I$) $\mathcal{A} \rightarrow \mathcal{A} \vee \mathcal{B}$
- ($\vee I$) $\mathcal{B} \rightarrow \mathcal{A} \vee \mathcal{B}$
- ($\wedge I$) $((\mathcal{A} \rightarrow \mathcal{B}) \wedge (\mathcal{A} \rightarrow \mathcal{C})) \rightarrow (\mathcal{A} \rightarrow (\mathcal{B} \wedge \mathcal{C}))$
- ($\vee E$) $((\mathcal{A} \rightarrow \mathcal{C}) \wedge (\mathcal{B} \rightarrow \mathcal{C})) \rightarrow ((\mathcal{A} \vee \mathcal{B}) \rightarrow \mathcal{C})$
- (\wedge - \vee) $\mathcal{A} \wedge (\mathcal{B} \vee \mathcal{C}) \rightarrow (\mathcal{A} \wedge \mathcal{B}) \vee (\mathcal{A} \wedge \mathcal{C})$
- (DNE) $\neg\neg\mathcal{A} \leftrightarrow \mathcal{A}$
- (MP) $\mathcal{A}, \mathcal{A} \rightarrow \mathcal{B} \Rightarrow \mathcal{B}$
- (ADJ) $\mathcal{A}, \mathcal{B} \Rightarrow \mathcal{A} \wedge \mathcal{B}$
- (Prefix) $\mathcal{A} \rightarrow \mathcal{B} \Rightarrow (\mathcal{C} \rightarrow \mathcal{A}) \rightarrow (\mathcal{C} \rightarrow \mathcal{B})$
- (Suffix) $\mathcal{A} \rightarrow \mathcal{B} \Rightarrow (\mathcal{B} \rightarrow \mathcal{C}) \rightarrow (\mathcal{A} \rightarrow \mathcal{C})$
- (RCont) $\mathcal{A} \rightarrow \mathcal{B} \Rightarrow \neg\mathcal{B} \rightarrow \neg\mathcal{A}$
- ($R\circ$) $\mathcal{A} \rightarrow (\mathcal{B} \rightarrow \mathcal{C}) \Leftrightarrow (\mathcal{A} \circ \mathcal{B}) \rightarrow \mathcal{C}$
- (Rt) $t \rightarrow \mathcal{A} \Leftrightarrow \mathcal{A}$

Each of the logics of interest is a first-order extension of some propositional extension of $\mathbf{B}^{\circ t}$. The propositional extensions are defined using the following list of axioms and rules:

- (A1) $\mathcal{A} \vee \neg\mathcal{A}$
- (A2) $\mathcal{A} \wedge (\mathcal{A} \rightarrow \mathcal{B}) \rightarrow \mathcal{B}$

⁷ N.b. that we lack function symbols in the signatures with which we are concerned; had the terms been built up with function symbols, this condition may have required also that no variable occurring in τ becomes bound in $\mathcal{A}[\tau/x]$.

⁸ A propositional language can be approximated by eliminating the quantifiers and taking only 0-ary predicates.

⁹ Note that “DNE” here stands for “double negation equivalence.”

- (A3) $(\mathcal{A} \rightarrow \mathcal{B}) \wedge (\mathcal{B} \rightarrow \mathcal{C}) \rightarrow (\mathcal{A} \rightarrow \mathcal{C})$
 (A4) $(\mathcal{A} \rightarrow (\mathcal{A} \rightarrow \mathcal{B})) \rightarrow (\mathcal{A} \rightarrow \mathcal{B})$
 (A5) $\mathcal{A} \rightarrow ((\mathcal{A} \rightarrow \mathcal{B}) \rightarrow \mathcal{B})$
 (A6) $\mathcal{A} \rightarrow (\mathcal{B} \rightarrow \mathcal{B})$
 (A7) $\mathcal{A} \rightarrow (\mathcal{B} \rightarrow \mathcal{A})$
 (A8) $(\mathcal{A} \rightarrow \mathcal{B}) \rightarrow ((\mathcal{A} \rightarrow \mathcal{C}) \rightarrow (\mathcal{A} \rightarrow \mathcal{B} \wedge \mathcal{C}))$
 (A9) $\mathcal{A} \rightarrow (\mathcal{A} \rightarrow \mathcal{A})$
 (A10) $\mathcal{A} \vee \mathcal{B} \rightarrow ((\mathcal{A} \rightarrow \mathcal{B}) \rightarrow \mathcal{B})$
 (A11) $(\mathcal{A} \wedge \mathcal{B} \rightarrow \mathcal{C}) \rightarrow (\mathcal{A} \wedge \neg \mathcal{C} \rightarrow \neg \mathcal{B})$
 (A12) $\mathcal{A} \rightarrow \neg(\mathcal{A} \rightarrow \neg \mathcal{A})$
 (A13) $(\mathcal{A} \rightarrow \neg \mathcal{A}) \rightarrow \neg \mathcal{A}$
 (A14) $(\mathcal{A} \rightarrow \mathcal{B}) \rightarrow (\neg \mathcal{B} \rightarrow \neg \mathcal{A})$
 (A15) $\mathcal{A} \rightarrow \mathcal{B} \vee \neg \mathcal{B}$
 (A16) $\mathcal{A} \rightarrow (\neg \mathcal{A} \rightarrow \mathcal{B})$
 (A17) $(\mathcal{A} \rightarrow \mathcal{B}) \rightarrow ((\mathcal{B} \rightarrow \mathcal{C}) \rightarrow (\mathcal{A} \rightarrow \mathcal{C}))$
 (A18) $(\mathcal{A} \rightarrow \mathcal{B}) \rightarrow ((\mathcal{C} \rightarrow \mathcal{A}) \rightarrow (\mathcal{C} \rightarrow \mathcal{B}))$
 (A19) $(\mathcal{A} \rightarrow (\mathcal{B} \rightarrow \mathcal{C})) \rightarrow (\mathcal{B} \rightarrow (\mathcal{A} \rightarrow \mathcal{C}))$
 (A20) $(\mathcal{A} \rightarrow (\mathcal{B} \rightarrow \mathcal{C})) \rightarrow ((\mathcal{A} \rightarrow \mathcal{B}) \rightarrow (\mathcal{A} \rightarrow \mathcal{C}))$
 (A21) $(\mathcal{A} \rightarrow \mathcal{B}) \rightarrow ((\mathcal{A} \rightarrow (\mathcal{B} \rightarrow \mathcal{C})) \rightarrow (\mathcal{A} \rightarrow \mathcal{C}))$
 (A22) $(\mathcal{A} \wedge \mathcal{B} \rightarrow \mathcal{C}) \rightarrow (\mathcal{A} \rightarrow (\mathcal{B} \rightarrow \mathcal{C}))$
- (R1) $\mathcal{C} \vee \mathcal{A} \Rightarrow \mathcal{C} \vee \neg(\mathcal{A} \rightarrow \neg \mathcal{A})$
 (R2) $\mathcal{C} \vee (\neg \mathcal{A} \rightarrow \mathcal{A}) \Rightarrow \mathcal{C} \vee \mathcal{A}$
 (R3) $\mathcal{C} \vee \mathcal{A}, \mathcal{C} \vee (\mathcal{A} \rightarrow \mathcal{B}) \Rightarrow \mathcal{C} \vee \mathcal{B}$
 (R4) $\mathcal{C} \vee (\mathcal{A} \rightarrow \mathcal{B}) \Rightarrow \mathcal{C} \vee (\neg \mathcal{B} \rightarrow \neg \mathcal{A})$
 (R5) $\mathcal{A} \Rightarrow (\mathcal{A} \rightarrow \mathcal{B}) \rightarrow \mathcal{B}$
 (R6) $\mathcal{C} \vee (\mathcal{A} \rightarrow \mathcal{B}), \mathcal{C} \vee (\mathcal{D} \rightarrow \mathcal{E}) \Rightarrow \mathcal{C} \vee ((\mathcal{B} \rightarrow \mathcal{D}) \rightarrow (\mathcal{A} \rightarrow \mathcal{E}))$

Some familiar and noteworthy logics are defined as follows:

$$\begin{array}{ll}
 \mathbf{G}^{\circ t} =_{df} \mathbf{B}^{\circ t} + (A1) & \mathbf{EW}^{\circ t} =_{df} \mathbf{TW}^{\circ t} + (R5) \\
 \mathbf{G}^{g \circ t} =_{df} \mathbf{G}^{\circ t} + (R1) & \mathbf{E}^{\circ t} =_{df} \mathbf{T}^{\circ t} + (R5) \\
 \mathbf{G}^{g, d \circ t} =_{df} \mathbf{G}^{g \circ t} + (R3) + (R4) & \mathbf{RW}^{\circ t} =_{df} \mathbf{EW}^{\circ t} + (A5) \\
 \mathbf{TW}^{\circ t} =_{df} \mathbf{B}^{\circ t} + (A14) + (A17) + (A18) & \mathbf{R}^{\circ t} =_{df} \mathbf{RW}^{\circ t} + (A4) \\
 \mathbf{T}^{\circ t} =_{df} \mathbf{TW}^{\circ t} + (A4) + (A13) & \mathbf{RM}^{\circ t} =_{df} \mathbf{R}^{\circ t} + (A9)
 \end{array}$$

Because all of the results given in this paper are for logics with fusion and the Ackermann truth constant, we save on notation by dropping the superscript. Except for when otherwise stated, we take \mathbf{L} to denote $\mathbf{L}^{\circ t}$. Importantly, we have to extend known γ -admissibility results for propositional results to include \circ and t . We do so in the appendix, where we drop this notational convention.

Note that the base logic for the paper is $\mathbf{G}^{g, d \circ t}$, which contains the two disjunctive rules (R3) and (R4). It is pointed out in Seki [29] that these two rules are required for

the normal models method to work for logics with both fusion and left implication. These logics, in Seki's terminology are L3 logics. As far as the authors can tell, both rules are required for each of fusion and left implication.

PROPOSITION 2.3 (*Conventionally Normal Propositional Logics*). *The propositional relevant logic \mathbf{G}^s , any extension of \mathbf{G}^s by axioms and rules with frame postulates being conjunctions of R and N statements (or implications with a single R statement in the antecedent), and several extensions of \mathbf{G}^s with implicational frame conditions with two R statements in which (R3) and (R1) are derivable admit γ [27, Section 5.6]. (See Seki [29] for the refinement from \mathbf{G} to \mathbf{G}^s .)*

Such logics are also called *Conventionally Normal* in Routley et al. [27].

THEOREM 2.4. *For any conventionally normal logic \mathbf{L} with (R3) and (R4) admitting γ , the logic $\mathbf{L}^{\circ t}$ also admits γ .*

Proof. See the appendix. □

This last theorem is needed because we include both fusion and little \mathbf{t} in our first-order formulations. We will say more about this fact later.

DEFINITION 2.5 (*First-Order Logics*). *Let \mathbf{L} be a propositional relevant logic (with fusion and \mathbf{t}) defined above. The logic \mathbf{LQ} is defined by adding the following axioms and rule schemes, in the first-order language:¹⁰*

- ($\forall E$) $\forall x.A \rightarrow A[\tau/x]$, where τ is free for x in A
- ($\exists I$) $A[\tau/x] \rightarrow \exists x.A$, where τ is free for x in A
- (EC) $\forall x.(A \vee B^x) \rightarrow \forall x.A \vee B^x$
- (dEC) $A^x \wedge \exists x.B \rightarrow \exists x.(A^x \wedge B)$
- ($A\exists E$) $\forall x.(A \rightarrow B^x) \rightarrow (\exists x.A \rightarrow B^x)$
- ($R\forall I$) $A^x \rightarrow B \Rightarrow A^x \rightarrow \forall x.B$
- ($R\exists E$) $A \rightarrow B^x \Rightarrow \exists x.A \rightarrow B^x$

Moreover, the logic \mathbf{QL} is defined similarly, but without (EC) and (dEC).

The principle (EC) is often described as the *extensional confinement axiom*; together with (dEC), we consider these both to be the extensional confinement axioms.

An important restriction for us is primitive inclusion of \circ in each of the defined logics. In some first-order logics, an extension with either \circ or even \leftarrow (the inclusion of which makes ($A\exists E$) derivable) is not a conservative extension [34]. With fusion, the following formula become a theorem (scheme):

$$(A\forall I) \forall x.(A^x \rightarrow B) \rightarrow (A^x \rightarrow \forall x.B).$$

Moreover, with (EC) and (dEC) as the only 'toggle' considered in this paper, there are many first-order relevant logics that we do not consider. In a neighbourhood setting, e.g., we can drop ($A\exists E$) and ($A\forall I$). We may also be able, though not in this paper, to

¹⁰ The reader is reminded that a super-scripted x means that x does not occur free is the decorated (sub)formula.

define and explore γ -admissibility in additional first-order relevant logics. We briefly discuss some of this freedom.

The following is a list of further rules and meta-rules to consider.

$$({}^d\mathbf{R}\forall\mathbf{G}) \mathcal{C}^x \vee \mathcal{B} \Rightarrow \mathcal{C}^x \vee \forall x\mathcal{B}$$

$$({}^d\mathbf{R}\exists\mathbf{G}) \mathcal{C}^x \vee \mathcal{B} \Rightarrow \mathcal{C}^x \vee \exists x\mathcal{B}$$

$$({}^d\mathbf{R}\forall\mathbf{I}) \mathcal{C}^x \vee (\mathcal{A}^x \rightarrow \mathcal{B}) \Rightarrow \mathcal{C}^x \vee (\mathcal{A}^x \rightarrow \forall x\mathcal{B})$$

$$({}^d\mathbf{R}\exists\mathbf{E}) \mathcal{C}^x \vee (\mathcal{A} \rightarrow \mathcal{B}^x) \Rightarrow \mathcal{C}^x \vee (\exists x\mathcal{A} \rightarrow \mathcal{B}^x)$$

MR1 If $\mathcal{A} \Rightarrow \mathcal{B}$, then $\mathcal{C} \vee \mathcal{A} \Rightarrow \mathcal{C} \vee \mathcal{B}$, where universal generalization (here, $(\mathbf{R}\forall\mathbf{I})$ ¹¹) is not used on a free variable in \mathcal{A} to obtain \mathcal{B} .

MR2 If $\mathcal{A} \Rightarrow \mathcal{B}$, then $\exists x\mathcal{A} \Rightarrow \exists x\mathcal{B}$, where universal generalization is not used on a free variable in \mathcal{A} to obtain \mathcal{B} .

It is easy to show that $({}^d\mathbf{R}\forall\mathbf{G})$ is a derivable rule in every first order logic \mathbf{LQ} , due to the presence of (EC). (A quick proof uses (EC), (MP), and the derivable $(\mathbf{R}\forall\mathbf{G})$.) In a logic \mathbf{QL} , it remains an open question whether or not this rule is derivable/admissible. The rule $({}^d\mathbf{R}\exists\mathbf{G})$ is derivable in every first-order logic, due to $(\exists\mathbf{I})$. Moreover, since $\forall x(\mathcal{A}^x \rightarrow \mathcal{B}) \rightarrow (\mathcal{A}^x \rightarrow \forall x\mathcal{B})$ is a theorem of any first-order logic (as defined in this paper), it is easy to show that $({}^d\mathbf{R}\forall\mathbf{I})$ is also a derivable rule with (R3). The case is similar for $({}^d\mathbf{R}\exists\mathbf{E})$, requiring (R3) again. Note, however, that (EC) and (dEC) are not needed for $({}^d\mathbf{R}\forall\mathbf{I})$ and $({}^d\mathbf{R}\exists\mathbf{E})$.

We therefore know then that \mathbf{LQ} will have each of the four disjunctive rules above, and that \mathbf{QL} sometimes has some of these rules. In 3, we show γ -admissibility for \mathbf{QL} and \mathbf{LQ} , provided that \mathbf{L} admits γ by normal models.

Finally, note that, as shown in Ferenz [8], $(\mathbf{A}\exists\mathbf{E})$ and $(\mathbf{A}\forall\mathbf{I})$ are valid in the Mares–Goldblatt semantics for relational semantics. In contrast, in Brady’s presentation of the content semantics of Brady [4], the base first-order relevant logics do not include these axioms (but do include the extensional confinement axiom(s)), although the later presentation in Brady [5] incorporates $(\mathbf{A}\exists\mathbf{E})$ and $(\mathbf{A}\forall\mathbf{I})$ into the setting of content semantics. While we get $(\mathbf{A}\forall\mathbf{I})$ for free using fusion, $(\mathbf{A}\exists\mathbf{E})$ is nonetheless valid on the Mares–Goldblatt semantics. (This also means that fusion does not conservatively extend Brady’s base system.) This contrast shows that the Mares–Goldblatt semantics and Brady’s content semantics diverge for first-order logics based on weak propositional relevant logics (at least for those captured by a relational semantics).

2.2. Mares–Goldblatt semantics. The target semantics is based on ternary relational semantics.

DEFINITION 2.6 (Ternary Relational Frames). *A ternary relational frame for \mathbf{B} is a tuple $\mathfrak{F} = \langle W, N, R, * \rangle$ where $\emptyset \neq N \subseteq W$, $R \subseteq W^3$, $* : W \rightarrow W$, and we further define, for each $a, b \in W$, $a \leq b =_{df} \exists x \in N(Rxab)$. Moreover, the following conditions are satisfied:*

¹¹ Brady [4, 5] has the restriction using the typical universal generalization rule $\mathcal{A} \Rightarrow \forall x\mathcal{A}$. We use $(\mathbf{R}\forall\mathbf{I})$ because it is equivalent in the presence of \mathbf{t} . MR1 and MR2 are used to define extensions of Brady’s \mathbf{BBQ} , where Brady’s \mathbf{BBQ} is strictly weaker than what we here would call \mathbf{BBQ} . Notably, $(\mathbf{A}\exists\mathbf{E})$ and $(\mathbf{A}\forall\mathbf{I})$ are not included.

- (c1) \leq is a preorder on W ;
- (c2) N is an (\leq) -upset;¹²
- (c3) If $a \leq a', b \leq b', c' \leq c$, and $Ra'b'c'$, then $Rabc$ ($R \downarrow \downarrow \uparrow$)
- (c4) $b \leq c$ implies $c^* \leq b^*$;
- (c5) $a^{**} = a$.

A ternary relational model for **B** is a frame with a valuation function $||-||$ that assigns an upset $||p|| \subseteq W$ to each propositional variable p . This assignment is extended to all formulas by the following:¹³

$$\begin{array}{ll} ||\mathbf{t}|| = N & ||\neg \mathcal{A}|| = \neg ||\mathcal{A}|| \\ ||\mathcal{A} \wedge \mathcal{B}|| = ||\mathcal{A}|| \cap ||\mathcal{B}|| & ||\mathcal{A} \vee \mathcal{B}|| = ||\mathcal{A}|| \cup ||\mathcal{B}|| \\ ||\mathcal{A} \rightarrow \mathcal{B}|| = |\mathcal{A}| \rightarrow ||\mathcal{B}|| & ||\mathcal{A} \circ \mathcal{B}|| = ||\mathcal{A}|| \circ ||\mathcal{B}|| \end{array}$$

For models for logics extending **B**: we provide the following list of frame conditions (saving space by writing an axiom's name instead of the entire axiom):

- (cA1) $a \in N \Rightarrow a^* \leq a$
- (cA2) $Raaa$
- (cA3) $Rabc \Rightarrow \exists x \in W(Rabx \ \& \ Raxc)$
- (cA4) $Rabc \Rightarrow \exists x \in W(Rabx \ \& \ Rxbc)$
- (cA5) $Rabc \Rightarrow Rbac$
- (cA6) $Rabc \Rightarrow b \leq c$
- (cA7) $Rabc \Rightarrow a \leq c$
- (cA8) $Rabc \ \& \ Rcdf \Rightarrow Radf \ \& \ Rbdf$
- (cA9) $Rabc \Rightarrow a \leq c \text{ or } b \leq c$
- (cA10) $Rabc \Rightarrow Rbac \ \& \ a \leq c$
- (cA11) $Rabc \Rightarrow \exists x \in W(b \leq x \ \& \ c^* \leq x \ \& \ Raxb^*)$
- (cA12) Ra^*aa^*
- (cA13) Raa^*a
- (cA14) $Rabc \Rightarrow Rac^*b^*$
- (cA15) $a^* \leq a$
- (cA16) $Rabc \Rightarrow a \leq b^*$
- (cA17) $Rabc \ \& \ Rcdf \Rightarrow \exists x \in W(Radx \ \& \ Rbxf)$
- (cA18) $Rabc \ \& \ Rcdf \Rightarrow \exists x \in W(Rbdx \ \& \ Raxf)$

¹² We define the 'upsets' as $\wp(W)^\uparrow = \{X \in \wp(W) : \forall a, b \in W (a \in X \ \& \ a \leq b) \Rightarrow b \in X\}$.

¹³ The operations \neg , \rightarrow , and \circ on subsets of W , on the right-hand side, are defined as follows:

1. $\neg X =_{df} \{a \in W : a^* \notin X\}$
2. $X \rightarrow Y =_{df} \{a \in W : \forall b, c \in W (Rabc \ \& \ b \in X \Rightarrow c \in Y)\}$
3. $X \circ Y =_{df} \{a \in W : \exists b, c \in W (Rbca \ \& \ b \in X \ \& \ c \in Y)\}$.

- (cA19) $Rabc \ \& \ Rcdf \Rightarrow \exists x \in W(Radx \ \& \ Rxbf)$
(cA20) $Rabc \ \& \ Rcdf \Rightarrow \exists x, y \in W(Radx \ \& \ Rbdy \ \& \ Rxyf)$
(cA21) $Rabc \ \& \ Rcdf \Rightarrow \exists x, y \in W(Radx \ \& \ Rbdy \ \& \ Rxyf)$
(cA22) $Rabc \ \& \ Rcdf \Rightarrow \exists x \in W(b \leq x \ \& \ d \leq x \ \& \ Raxf)$
(cR1) $a \in N \Rightarrow Ra^*aa^*$
(cR2) $a \in N \Rightarrow Raa^*a$
(cR3) $a \in N \Rightarrow Raaa$
(cR4) $a \in N \ \& \ Rabc \Rightarrow Rac^*b^*$
(cR5) $\exists x \in N(Raxa)$
(cR6) $a \in N \ \& \ Rabc \ \& \ Rcdf \Rightarrow \exists x, y \in W(Radx \ \& \ Rbxy \ \& \ Rayf)$

(This list is to be read as (cX) is the frame condition corresponding to the axiom of rule scheme (X).)

DEFINITION 2.7 (Models for \mathbf{LQ}). A Mares–Goldblatt frame for \mathbf{LQ} (an \mathbf{LQ} -frame), for a propositional relevant logic \mathbf{L} is a tuple $\mathfrak{F} = \langle W, N, R, *, U, Prop, PropFun \rangle$, where $\langle W, N, R, * \rangle$ is an \mathbf{L} -frame, U is a non-empty set, and we have that $Prop \subseteq \wp(W)^\uparrow$, $PropFun \subseteq \{\varphi : U^\omega \longrightarrow Prop\}$. Moreover, the following conditions are satisfied:

- (cq1) $Prop$ contains N , and is closed under $\cap, \cup, \neg, \rightarrow, \circ$;
(cq2) $PropFun$ contains a constant function φ_N ($\varphi_N f = N$), and is closed under $\cap, \cup, \neg, \rightarrow, \circ, \forall_n$ and \exists_n , for every $n \in \omega$, where
(a) $(\neg\varphi)f = \neg(\varphi f)$
(b) $(\varphi \otimes \psi)f = \varphi f \otimes \psi f$, for each $\otimes \in \{\cap, \cup, \rightarrow, \circ\}$
(c) $(\forall_n \varphi)f = \bigcap_{g \sim_{x_n} f} \varphi g = \bigcup \{X \in Prop \mid X \subseteq \bigcap_{g \sim_{x_n} f} \varphi g\}$
(d) $(\exists_n \varphi)f = \bigcup_{g \sim_{x_n} f} \varphi g = \bigcap \{X \in Prop \mid \bigcup_{g \sim_{x_n} f} \varphi g \subseteq X\}$.
(cq3) For every $\varphi \in PropFun$, $X, Y \in Prop$, $n \in \omega$, and $f \in U^\omega$ ¹⁴
(cEC) $X - Y \subseteq \bigcap_{j \in U} \varphi(f[j/n])$ only if $X - Y \subseteq (\forall_n \varphi)f$
(cdEC) $\bigcup_{j \in U} \varphi(f[j/n]) \subseteq X \cup \overline{Y}$ only if $(\exists_n \varphi)f \subseteq X \cup \overline{Y}$.

A pre-model for \mathbf{LQ} is a tuple $\mathfrak{M} = \langle F, |\cdot| \rangle$ such that F is a Mares–Goldblatt frame for \mathbf{LQ} and $|\cdot|$ is a valuation function that assigns:

1. an individual $|c| \in U$ to each constant symbol c ;
2. a function $|P^n| : U^n \longrightarrow \wp(W)$ to each n -ary predicate symbol P^n ; and
3. a propositional function $|A| : U^\omega \longrightarrow \wp(W)$ to each formula A such that, when A is atomic, for every $f \in U^\omega$:

$$|P^n \tau_1, \dots, \tau_n|f = |P^n|(|\tau_1|f, \dots, |\tau_n|f)$$

¹⁴ Note that $X - Y$ and \overline{Y} are defined in the usual set-theoretic sense, but that $Prop$ is not necessarily closed under either of these operations.

where “ $|\tau|f$ ” is fn when τ is the variable x_n , and $|c|$ when τ is constant symbol c . Moreover, when \mathcal{A} is not atomic (or \mathbf{t}), the valuation is extended as follows:

$$\begin{array}{ll} |\mathbf{t}| = \phi_N & |\neg\mathcal{A}| = \neg|\mathcal{A}| \\ |\mathcal{A} \wedge \mathcal{B}| = |\mathcal{A}| \cap |\mathcal{B}| & |\mathcal{A} \circ \mathcal{B}| = |\mathcal{A}| \circ |\mathcal{B}| \\ |\mathcal{A} \vee \mathcal{B}| = |\mathcal{A}| \cup |\mathcal{B}| & |\mathcal{A} \rightarrow \mathcal{B}| = |\mathcal{A}| \rightarrow |\mathcal{B}| \\ |\forall x_n \mathcal{A}| = \forall_n |\mathcal{A}| & |\exists x_n \mathcal{A}| = \exists_n |\mathcal{A}| \end{array}$$

A model for \mathbf{LQ} is a pre-model for \mathbf{LQ} that assigns an element of *Prop* to each atomic formula.

A formula \mathcal{A} is *satisfied* by a variable assignment f in a model \mathfrak{M} , written $\mathfrak{M}, f \models \mathcal{A}$, when $N \subseteq |\mathcal{A}|f$. A formula is *valid in a model* \mathfrak{M} ($\mathfrak{M} \models \mathcal{A}$) when it is satisfied by every variable assignment in that model; *valid in a frame* \mathfrak{F} ($\mathfrak{F} \models \mathcal{A}$) when it is valid in every model based on that frame; *valid in a class of frames* \mathbb{C} ($\mathbb{C} \models \mathcal{A}$) when it is valid in every frame in that class. Admissibility of a rule in the semantic context is understood as the preservation of validity.

PROPOSITION 2.8 (Soundness and Completeness for \mathbf{LQ}). *For a wide class of logics including \mathbf{B} and its usual extensions, Ferenz [8] has shown that \mathbf{LQ} is sound and complete w.r.t. the class of \mathbf{LQ} -frames.*

To prepare the reader for the proof of γ -admissibility, we will briefly sketch the technique of normalization and describe its history. Although the problem of γ -admissibility was tackled by Meyer & Dunn [20], the algebraic techniques employed there are relatively involved. Routley & Meyer [26] offered a new technique of *normalization* as a simpler and more elegant form of argument of γ -admissibility whose development would continue in Routley & Meyer [25] and Routley et al. [27]. The shape of the technique is straightforward: A Routley–Meyer model is described as *normal* if for some normal point a , $a = a^*$, i.e., some normal point is its own star point.¹⁵

Not all \mathbf{R} models are normal (and *a fortiori* for weaker relevant logics), but surprisingly Routley & Meyer [26] describe a recipe through which one can pick an arbitrary point $o \in N$ and *normalize* a model to include a point $0 \in N$ such that $o^* \leq 0 = 0^* \leq o$. In particular, the condition for the excluded middle entails that $o^* \leq o$, and we ensure that the new 0 is a negation-consistent point. We may think of 0 as a consistitized version of o . If the logic is sound and complete with respect to normal models with appropriate frame conditions, then, if $\neg\mathcal{A} \vee \mathcal{B}$ and \mathcal{A} are theorems, they will be true in 0 . Since $0 = 0^*$, we can conclude that \mathcal{A} is not true at 0 , whence by the truth conditions for disjunction, \mathcal{B} will be true at 0 and also at o . As o was arbitrarily selected, this means that \mathcal{B} is semantically valid and thus a theorem by completeness.

This sketch, of course, has a great many subtleties and nuances. It is the task of the next section to fill in the gaps and transform this sketch into a proof.

¹⁵ *N.b.* that the definition in Routley & Meyer [26] is that this property holds for a *distinguished point* but the effect is the same.

§3. γ -Admissibility for QL and LQ logics. While explicitly stated in some definitions and lemmata, in this section we assume \mathbf{L} to denote a propositional relevant logic that admits γ via the normal models method and contains (R3) and (R4): that is, a *conventionally normal L3* extension of $\mathbf{G}^{g,d}$. The keystone of this paper is the method of normal models, so we present the crucial definition.

DEFINITION 3.9 (Normal Models). *For any logic \mathbf{LQ} or \mathbf{QL} (based on a propositional relevant logic \mathbf{L} with (R3) and (R4) admitting γ by the normal models method) defined above, an \mathbf{LQ} -(\mathbf{QL})-model (-frame) is normal if it satisfies the following:*

(Norm) $a = a^*$, for some $a \in N$.

DEFINITION 3.10 (Normalization of a frame \mathfrak{F}). *Where \mathbf{L} is a conventionally normal L3 logic, $0 \notin W$ and $o \in N$, the normalization of a \mathbf{QL} -frame or \mathbf{LQ} -frame $\mathfrak{F} = \langle W, N, R, *, U, Prop, PropFun \rangle$ is a frame $\langle W', N', R', *, U, Prop', PropFun' \rangle$, defined by:¹⁶*

1. $W' = W \cup \{0\}$;
2. $N' = N \cup \{0\}$;
3. R' is given by $R'abc$ iff $Rabc$, whenever $a, b, c \in W$, and when $a, b \in W$, R' satisfies the following
 - (a) $R'000$;
 - (b) $R'00a$ iff $R'0oa$;
 - (c) $R'0a0$ iff $R'0ao^*$;
 - (d) $R'a00$ iff $R'aoo^*$;
 - (e) $R'0ab$ iff $R'oab$;
 - (f) $R'a0b$ iff $R'aob$;
 - (g) $R'ab0$ iff $R'abo^*$;
4. $*$ ' is defined by:
 - (a) $a^{*'} = a^*$ when $a \in W$;
 - (b) $0^{*'} = 0$;
5. For each $X \in Prop$, add $X \in Prop'$ when $o^* \notin X$ and, when $o \in X$, $X \cup \{0\} \in Prop'$;
6. For each $\varphi \in PropFun$, add φ and φ' to $PropFun'$, where φ' is defined by:¹⁷

$$\forall f \in U^\omega, o \in \varphi f \text{ implies } \varphi' f = \varphi f \cup \{0\}.$$

We called this normalized frame the normalization of \mathfrak{F} at 0 for $o \in N$.

The proof of the following Lemma and Corollary are standard. That is, note that their statement and proof (as in, e.g., Seki [29]) relies only on the *propositional machinery* of a frame: $U, Prop$, and $PropFun$ are irrelevant.

LEMMA 3.11. *If \mathfrak{F} is an \mathbf{QL} - or \mathbf{LQ} -frame (for conventionally normal L3 \mathbf{L}) and \mathfrak{F}' is a normalization of \mathfrak{F} at 0 for o :*

¹⁶ The o and 0 are used as in Seki [29], and correspond to T and T' of Routley et al. [27, p. 387].

¹⁷ The reader is to especially note that $PropFun'$ is not that much of an extension of $PropFun$. That is, the reader is to note the scope of the universal quantifier in the definition. There is no 'mixing' in the image of a propositional function — that is, a propositional function in $PropFun'$ either returns the new elements of $Prop'$ with 0 , or else the old elements of $Prop$ that do not contain 0 .

1. the relation R' is well-defined;
2. the ordering \leq' is such that, for all $a, b \in W$:
 - (a) $a \leq' b$ iff $a \leq b$;
 - (b) $0 \leq' b$ iff $0 \leq b$;
 - (c) $a \leq' 0$ if $a \leq o^*$.

COROLLARY 3.12. *If \mathfrak{F} is an **QL**- or **LQ**-frame (for conventionally normal $L3$ **L**) and \mathfrak{F}' is a normalization of \mathfrak{F} at 0 for o , then $o^* \leq 0 \leq o$.*

The next lemma does require extra verification of the first-order machinery. Of course, U is unaffected, but we must show that $Prop$ and $PropFun$ are well-defined and closed under the required operations.

LEMMA 3.13. *Suppose that L is a propositional relevant logic (extending $G^{g,d}$) which admits γ (by the method of normalizing models). Let \mathfrak{F} be a **QL**- or **LQ**-frame and*

$$\mathfrak{F}' = \langle W', N', R', *, U, Prop', PropFun' \rangle,$$

*the normalization of \mathfrak{F} at 0 for o . Then \mathfrak{F}' is also a **QL**- or **LQ**-frame, respectively.*

Proof. Here we must check every frame postulate. By the supposition that L admits γ by the method of normalization (see the appendix), we can use the arguments of Routley et al. [27, pp. 389–390] Seki [29, pp. 214–216] to cover to show that the frame postulates corresponding to **L** (including those of $G^{g,d}$) are satisfied (with respect to W', N', R' and $*$).

What remains to show is that $Prop'$ and $PropFun'$ are well defined and that conditions (cq1) and (cq2) hold for **QL**, and additionally that (cq3) holds for **LQ**.

$Prop'$ is well-defined: We need to check that each element of $Prop'$ is an \leq -upset. For each such X not containing 0, we need to show that $o^* \notin X$, but this is so by definition. Suppose that $0 \in X' \in Prop'$, and that $\exists y \in X' (y \leq z \ \& \ z \notin X')$. As X' comes from an element, say X , of $Prop$, such a y cannot exist in X' . So let $y = 0$. This means that $0 \leq z$ and $z \notin X'$. However, by Lemma 3.11(2).(b), $0 \leq z$ and thus $z \in X$, a contradiction. Hence $Prop$ is well-defined.

$PropFun'$ is well-defined: It is easy to see that each element $\varphi \in PropFun'$ is a function that produces a unique element of $Prop'$ as output, given a particular input.

(cq1): Suppose that $X', Y' \in Prop'$ (and that $X' = X \cup \{0\}$, if $0 \in X'$, for some $X \in Prop$, a similarly for Y/Y'). We give the cases as follows:

- ⊂ If $0 \notin X', Y'$, then $0 \notin X' \cap Y' \in Prop$, and hence in $Prop'$. If $0 \in X'$ but $0 \notin Y'$, then $X' \cap Y' = X \cap Y \in Prop'$. If $0 \in X', Y'$, then $X' \cap Y' = \{0\} \cup (X \cap Y)$, and $X \cap Y \in Prop$. (The remaining sub-cases are symmetrical to the second sub-case.)
- ⊃ This case is similar to the previous case. One sub-case is shown. Suppose that $0 \in X', Y'$. Then $X' \cup Y' = \{0\} \cup (X \cup Y)$, where $X \cup Y \in Prop$.
- Either $0 = o^* \in X'$ or not. If it is, then $o \in X$. Then $o^* \notin \neg X$, and $o \notin \neg X$. From the latter, $\neg X' = \neg X \in Prop$. On the other hand, the assumption that $0 \notin X' (= X)$ implies that $o \notin X$. Thus $o^* \in \neg X$, and by Lemma 3.12 we have $o \in \neg X$, and so $0 \in \neg X' = \{0\} \cup \neg X$.
- The case where $X' = X$ and $Y' = Y$ is trivial. Suppose that $X' = X \cup \{0\}$ and $Y' = Y$. We know that $Ra0c$ iff $Raoc$, and $X' \rightarrow Y = \{a \in W' : \forall b, c ((Rabc \ \& \ b \in X') \Rightarrow c \in Y)\}$, which means that $a \in X' \rightarrow Y$ iff $a \in X \rightarrow Y$, which means that $X' \rightarrow Y = X \rightarrow Y \in Prop'$.

On the other hand, if $X' = X$ and $Y' = Y \cup \{0\}$, then we note that $R'ab0$ iff $Rabo^*$. So $X' \rightarrow Y' = X \rightarrow Y'$. If $a \in X \rightarrow Y'$ and $R'ab0$, then $R'abo^*$, and so $a \in X \rightarrow Y$. Finally, suppose that $X' = X \cup \{0\}$ and $Y' = Y \cup \{0\}$. We combine the reasoning of the previous two cases to show $X' \rightarrow Y' = X \rightarrow Y$. (Using $R'a00$ iff $R'ao0^*$ where needed.)

◦ Similar to the \rightarrow case.

(cq2): Suppose that $\varphi', \psi' \in PropFun$: The cases are as follows:

φ'_N : It is easy to check that $(\varphi_N)'$ is our desired φ'_N .

\cap For the remaining cases, we show only the subcases where one of φ' or ψ' is not in $PropFun$. It is easy to check that, for each $f \in U^\omega$, $\varphi'f \cap \psi'f = (\varphi \cap \psi)'f$, which shows that $PropFun'$ is closed under (the lifted operator) \cap .

\cup Similar to the previous case.

\neg We show that $\neg(\varphi'f) = (\neg\varphi)'f$, for each $f \in U^\omega$. If $0 \notin \neg(\varphi'f) \cup (\neg\varphi)'f$, then $\neg(\varphi'f) = \neg(\varphi f) = (\neg\varphi)f = (\neg\varphi)'f$. On the other hand, $0(0^*) \in \neg(\varphi'f)$ iff $0^*(0) \notin (\varphi'f)$ iff $o^* \notin (\varphi)f$ iff $o \in \neg(\varphi f) = (\neg\varphi)f$ iff $0 \in (\neg\varphi)'f$.

\rightarrow Fixing an f , we must show that $(\varphi' \rightarrow \psi')f = (\varphi \rightarrow \psi)'f$. To handle multiple cases in parallel, let the notation $\llbracket a/0 \rrbracket$ denote a in case $a \neq 0$ and 0 otherwise. Then for left-to-right, we prove the contrapositive. If $\llbracket a/0 \rrbracket \notin (\varphi \rightarrow \psi)'f$ then $\llbracket a/o \rrbracket \notin (\varphi \rightarrow \psi)f$, meaning that there exist $b, c \in W$ such that $R\llbracket a/o \rrbracket bc$ and $b \in \varphi f$ and $c \notin \psi f$, i.e., there is a counterexample. Counterexamples will lift to the new model, i.e., $R'\llbracket a/0 \rrbracket bc$ will hold. Also as $\varphi f \subseteq \varphi'f$, $b \in \varphi'f$ and as $\psi'f \cap W = \psi f$, that $c \notin \psi f$ entails that $c \notin \psi'f$. Together, $\llbracket a/0 \rrbracket \notin (\varphi'f \rightarrow \psi'f)$. For right-to-left, if $\llbracket a/0 \rrbracket \in (\varphi \rightarrow \psi)'f$ then $\llbracket a/o \rrbracket \in (\varphi \rightarrow \psi)f$, meaning that in the original model for all $b, c \in W$ such that $R\llbracket a/o \rrbracket bc$, if $b \in \varphi f$ then $c \in \psi f$. Suppose for contradiction that $\llbracket a/0 \rrbracket \notin (\varphi'f \rightarrow \psi'f)$. Then there are $d, e \in W'$ for which $R'\llbracket a/0 \rrbracket de$ while $d \in \varphi'f$ and $e \notin \psi'f$. But (using the same notation) this means that in the original model $R\llbracket a/o \rrbracket \llbracket d/o \rrbracket \llbracket e/o^* \rrbracket$ with $\llbracket d/o \rrbracket \in \varphi f$ and $\llbracket e/o^* \rrbracket \notin \psi f$. But since this takes place in the original model, $\llbracket e/o^* \rrbracket$ would be forced to be a member of ψf . Consequently, $\llbracket a/0 \rrbracket \in (\varphi'f \rightarrow \psi'f)$.

◦ Similar to the \rightarrow case.

\forall_n We show that $(\forall_n \varphi')f = (\forall_n \varphi)'f$, for every $f \in U^\omega$. Fix an arbitrary $f \in U^\omega$. There are two distinct cases to consider $0 \in (\forall_n \varphi)'f$ and $0 \notin (\forall_n \varphi)'f$. For the latter, that is $0 \notin (\forall_n \varphi)'f$, this entails that $o \notin (\forall_n \varphi)f$. That is, for all $X \in Prop$, $X \subseteq \bigcap_{g \sim_x f} (\varphi g)$ entails $o \notin X$. By definition, it follows that each $\varphi'g \in \bigcap_{g \sim_x f} (\varphi'g)$ does not contain 0 . And so, $0 \notin \bigcap_{g \sim_x f} (\varphi'g) = (\forall_n \varphi')f$. That $(\forall_n \varphi')f = (\forall_n \varphi)'f$ follows from the fact that each set belongs to the original model, where the identity holds.

Now suppose that $0 \in (\forall_n \varphi)'f$. Then $o \in (\forall_n \varphi)f$ and thus $o \in X \subseteq \bigcap_{g \sim_x f} (\varphi g)$ for some $X \in Prop$. Then $o \in \varphi g$ for every g . Let us denote $X \cup \{0\}$ by X' . Then $X' \subseteq \bigcap_{g \sim_x f} (\varphi'g)$, as by definition $\varphi'g$ must include 0 . But then $0 \in (\forall_n \varphi')f$. Using the fact that $0 \in (\forall_n \varphi')f$ iff $0 \in (\forall_n \varphi)'f$, we now show the identity $(\forall_n \varphi')f = (\forall_n \varphi)'f$. The $a = 0$ case is covered by what we have already shown. If $a \neq 0$, then the corresponding identity of the original model is sufficient.

\exists_n Similar but dual to the previous case.

This completes the **QL** portion of the lemma. The remaining case concerns **LQ**.

(cq3): We show only the sub-case for (cEC). Assume that $X', Y' \in Prop'$, $\varphi' \in PropFun'$, $n \in \omega$, and $f \in U^\omega$. Further suppose that $X' - Y' \subseteq \bigcap_{j \in U} \varphi'(f[j/n])$. There are two cases.

Case 1: $0 \in X' - Y'$, which is $0 \in X'$ and $0 \notin Y'$. Then $0 \in \bigcap_{j \in U} \varphi'(f[j/n])$. Thus, $0 \in$

$\varphi'(f[j/n])$ for each $j \in U$. We want to show that there is an $X \in Prop'$ such that $0 \in X \subseteq \varphi'(f[j/n])$ for each $j \in U$. Consider the set $\forall_n \varphi' f \cup \{0\}$. This is indeed an element of $Prop'$: $0 \in \varphi'(f[j/n])$ entails $o \in \varphi'(f[j/n])$, which forces in turn both $o \in \forall_n \varphi' f$ and $0 \in \forall_n \varphi' f \cup \{0\} \in Prop'$. This completes the case with $X = \forall_n \varphi' f \cup \{0\}$. Because the original frame satisfied (cEC), we have shown that every element of $X' - Y'$ is an element of $(\forall_n \varphi')f$, as required.

Case 2: $0 \notin X' - Y'$. We assume that $X' - Y' \subseteq \bigcap_{j \in U} \varphi'(f[j/n])$. As $0 \notin X' - Y'$, the

result follows from the original model satisfying (cEC) and the fact that both $\bigcap_{j \in U} \varphi'(f[j/n])$ and $(\forall_n \varphi')f$ are either equal to their corresponding sets in the original model, or additionally contain 0. \square

As each normalized frame is a frame of the right kind, soundness for the corresponding logic is straightforward. We record this fact.

LEMMA 3.14 (Normal Soundness). *For any formula A , if A is a theorem of **QL** or **LQ** on **L3 L**, then A is valid in every normal **QL**-frame or **LQ**-frame, respectively.*

Given a **LQ**-model's valuation, we define the standard valuation for the normalization of the model's frame in the following definition. Note that other valuations are possible, but that this standard normalization valuation plays a key role in what's to come.

DEFINITION 3.15. *If $\mathfrak{M} = \langle \mathfrak{F}, |- \rangle$ is an **QL**- or **LQ**-model (for conventionally normal **L3** logic **L**), we take as the standard normalization of model \mathfrak{M} at 0 (for $0 \in N$) to be the tuple $\mathfrak{M}' = \langle \mathfrak{F}', |-'| \rangle$, where \mathfrak{F}' is the normalization of \mathfrak{F} (at 0), and $|-'|$ is defined as follows:¹⁸*

1. $|c|' = |c|$;
2. for all $\vec{j} \in \mathcal{U}^n$: $|P^n|'(\vec{j}) = |P^n|(\vec{j})$, if $o \notin |P^n|(\vec{j})$, and $(|P^n|(\vec{j})) \cup \{0\}$ if $o \in |P^n|(\vec{j})$;
3. A propositional function $|A|'$ is given to each formula in the usual way, given the previous two clauses.

LEMMA 3.16. *Let **L** be a conventionally normal **L3** propositional relevant logic. Given a **QL**- or **LQ**-model $\mathfrak{M} = \langle \mathfrak{F}, |- \rangle$, the standard normalization $\mathfrak{M}' = \langle \mathfrak{F}', |-'| \rangle$ of \mathfrak{M} is a **QL**- or **LQ**-model, respectively.*

Proof. The underlying frame is an **QL**- or **LQ**-frame, as per Lemma 3.13. It remains to be shown that the valuation assigns an element of $PropFun'$ to each atomic

¹⁸ Note that this terminology is new, but reflects the usual method of normalization.

proposition. That every formula is assigned an element of $PropFun'$ follows from each atomic formula being assigned an element of $PropFun'$ together with the fact that $PropFun'$ is closed under the required operations.

It is straightforward to check that each atomic formula is mapped to an element of $PropFun'$. As $PropFun'$ is closed under the appropriate operators, it follows in the usual way that each formula is mapped to an element of $PropFun'$. \square

LEMMA 3.17. *Let \mathfrak{M} be a QL - or LQ -model with set W (for conventionally normal $L3$ logic L). Further let \mathfrak{M}' be the standard normalization of \mathfrak{M} . For all $a \in W$, for every formula A and $f \in U^\omega$, $a \in |A|f$ iff $a \in |A'|f$.*

Proof. The proof is by induction on the complexity of A . If A is the atomic $P\tau_1, \dots, \tau_n$, then $|P\tau_1, \dots, \tau_n|f = |P|'(|\tau_1|'f, \dots, |\tau_n|'f)$. For each $|\tau_i|'f$, $|\tau_i|'f = |\tau_i|f$. Thus, by definition, $|P\tau_1, \dots, \tau_n|f$ restricted to W is $|P\tau_1, \dots, \tau_n|f$, as required. The case for t is straightforward. For the inductive cases, we only show a couple.

Case $A = B \vee C$: Suppose that $a \in W$ and $f \in U^\omega$. If $a \in |B \vee C|f$, then by (a couple suppressed steps) the inductive hypothesis either $a \in |B|'f$ or $a \in |C|'f$, which entails that $a \in |B \vee C|'f$, as required. The other direction is similarly straightforward.

Case $A = B \rightarrow C$: Right-to-left is trivial as R' and R agree on all arguments from W . For left-to-right, suppose that $a \in |B \rightarrow C|f$. Then for all $b, c \in W$ such that $Rabc$ and $b \in |B|f$, also $c \in |C|f$. By induction hypothesis, this entails that for all $b, c \in W$ such that $R'abc$ and $b \in |B|'f$, also $c \in |C|'f$. This is *nearly* sufficient to establish that $a \in |B \rightarrow C|'f$; it could go away only in case $R'ab'c'$, $b' \in |B|'f$, and $c' \in |C|'f$ when either $b' = 0$ or $c' = 0$. But in such cases, one could select appropriate $b'' = 0$ or $c'' = 0^*$ such that $R'ab''c''$ while $b'' \in |B|'f$ and $c'' \notin |C|'f$. As $a, b'', c'' \in W$, though, this is impossible, as it would entail that $c'' \in |C|'f$. Thus $a \in |B \rightarrow C|'f$, as required.

Case $A = \forall x_n B$: Suppose that $a \in W$, $f \in U^\omega$ and that $a \in |\forall x_n B|f$. Then $a \in X \in Prop$ and $X \subseteq \bigcap_{g \sim_{x_n} f} |B|g$. Then $a \in |B|g$, for each $g \sim_{x_n} f$. By the induction hypothesis, for all $b \in W$, $b \in |B|'g$ iff $b \in |B|g$, for each such g . Thus (i) $a \in \bigcap_{g \sim_{x_n} f} |B|'g$ and (ii) $X \subseteq \bigcap_{g \sim_{x_n} f} |B|'g$ (because $X \subseteq W$). Then $a \in X \in Prop'$ and $X \subseteq \bigcap_{g \sim_{x_n} f} |B|g$, which is that $a \in |\forall x_n B|'f$, as required.

For the other direction, assume that $a \in W$, $f \in U^\omega$ and that $a \in |\forall x_n B|'f$. Then $a \in X' \in Prop'$ and $X' \subseteq \bigcap_{g \sim_{x_n} f} |B|'g$. Consider X , which is equal to X' if $X' \subseteq W$, and is $X' - \{0\}$ otherwise. Clearly $X \in Prop$. Moreover, by the transitivity of the subset relation, $X \subseteq \bigcap_{g \sim_{x_n} f} |B|'g$, and also we have $a \in X$. Now, by the induction hypothesis, for every $b \in W$, $b \in |B|'g$ iff $b \in |B|g$, for each such g . Thus we infer that $X \subseteq \bigcap_{g \sim_{x_n} f} |B|g$, and so $a \in |\forall x_n B|f$. \square

THEOREM 3.18. *For any formula A and any $L3$ logic L admitting γ by normal models, A is a theorem of LQ (QL) iff A is valid in every normal LQ -frame (QL -frame).*

Proof. The *only if* direction is soundness, and is covered by Lemma 3.14. For the *if* direction, suppose that A is not a theorem of LQ (QL). Then there is a canonical LQ -model (QL -model) with frame $\mathfrak{F} = \langle W, N, R, 0, U, Prop, PropFun \rangle$, (canonical) valuation $|\cdot|$, and $o \in N$ such that $o \notin |A|f$, for some $f \in U^\omega$.

For a new 0 , take the standard normalization of \mathfrak{M} (at 0 for o), denoted $\mathfrak{M}' = \langle \mathfrak{F}', |\cdot|' \rangle$. By Lemma 3.16, this \mathfrak{M}' is a LQ -model (QL -model). By Lemma 3.17,

$o \notin |\mathcal{A}|'f$. But $0 \leq o$ then entails that $0 \notin |\mathcal{A}|'f$, and since $0 \in N'$, we have that \mathcal{A} is not valid on \mathfrak{F} . \square

In particular, this proof shows that, for every invalid formula \mathcal{A} , there is a normal point $0 = 0^*$ in a normal model at which the invalidity of \mathcal{A} is witnessed. In the proof we took at arbitrary $o \in N$ at which \mathcal{A} fails and introduced a new point 0 which, because of the excluded middle axiom and out construction, is such that $o^* \leq 0 \leq o$. The latter entails that 0 is also a point at which \mathcal{A} fails.

COROLLARY 3.19. *For the logics \mathbf{LQ} and \mathbf{QL} , where $\mathbf{L3}$ logic \mathbf{L} admits γ by normal models, the rule γ is admissible.*

Proof. Suppose that $\neg\mathcal{A} \vee \mathcal{B}$ and \mathcal{A} are both theorems of \mathbf{LQ} . Then by Theorem 3.18, these formulas are valid on every normal model. Consider an arbitrary normal model $\mathfrak{M} = \langle W, N, R, *, U, Prop, PropFun, |\cdot| \rangle$ with normal point 0 ($0 = 0^*$). Since $0 \in N$, $0 \in |\neg\mathcal{A} \vee \mathcal{B}|f \cap |\mathcal{A}|f$ for every $f \in U^\omega$. Since $0 \in |\mathcal{A}|f$ and $0 = 0^*$, we have that $0 \notin |\neg\mathcal{A}|f$. But then given the definition of $|\neg\mathcal{A} \vee \mathcal{B}|f$, $0 \in |\mathcal{B}|f$, as required. \square

We reintroduce the \circ and \mathfrak{t} notation in a logic's name for the next corollary, which gives a sufficient condition for γ -admissibility in logics without \circ and \mathfrak{t} .

COROLLARY 3.20. *For every conventionally normal propositional logic \mathbf{L} , if $\mathbf{LQ}^{\circ\mathfrak{t}}$ admits γ and conservatively extends \mathbf{LQ} , then \mathbf{LQ} admits γ .*

Proof. Suppose that $\vdash_{\mathbf{LQ}} \mathcal{A}$ and $\vdash_{\mathbf{LQ}} \neg\mathcal{A} \vee \mathcal{B}$ and that \mathcal{A} and \mathcal{B} do not contain fusion or \mathfrak{t} . Then we have $\vdash_{\mathbf{LQ}^{\circ\mathfrak{t}}} \mathcal{B}$ since $\mathbf{LQ}^{\circ\mathfrak{t}}$ admits γ . Moreover, since \mathcal{B} is in the language of \mathbf{LQ} , by the conservative extension assumption we have $\vdash_{\mathbf{LQ}} \mathcal{B}$. \square

As a result, whenever we can show the conservative extension by \circ and \mathfrak{t} in the first-order case, we can extend our admissibility results to the weaker logics.

§4. Concluding remarks. We have shown γ is admissible in a wide range of first-order relevant logics. A major upshot is that we can conceive of many logics \mathbf{QL} and \mathbf{LQ} as having a well-behaved semantics. Well behaved, that is, in the sense that the machinery for interpreting quantified formulas is sufficiently independent from the propositional machinery required for γ -admissibility. Thus, we have essentially shown that the \mathbf{QL} and \mathbf{LQ} extensions of \mathbf{L} conserve the property of γ -admissibility (w.r.t. the normal models method).¹⁹ From this point of view, and from the fact that γ -admissibility ensures the set of theorems of a logic is negation consistent (and that the logic contains the set of theorems of classical first-order logic in \neg, \vee, \forall), we can take the logics that we have shown to admit γ as well-behaved in yet another sense: that the constant domain extensions \mathbf{QL} and \mathbf{LQ} preserve normal models γ -admissibility, negation consistency of the set of theorems, and the containment of (the theorems of) their classical counterparts.

There are, however, many other ways to axiomatize first-order extensions of relevant logics. From this work, we plan to extend these results to first-order modal relevant logics and first-order logics that require neighbourhood semantics. (The neighbourhood semantics given in Tedder & Ferenz [34] is apt for a normal model

¹⁹ We thank an anonymous reviewer for the viewpoint that our results are a kind of conservative extension result.

approach, as negation is treated in a similar way using a star function. Similarly, the framework of Ferenz & Tedder [10] may be used for weaker modal relevant logics.) For the modal cases, the work of Seki [29] will again be invaluable. In the neighbourhood cases, we are able to better model axiomatizations of first-order logics that, e.g., drop the axiom forms of universal introduction and existential elimination.

A further avenue for future work is to extend the investigation in the paper Ferenz & Ferguson [9] in which γ admissibility holds in the case of the varying-domain version of **QR** described in Mares [18] to examine varying-domain analogues of the weaker relevant logics considered in this paper. Notably, Mares' system includes an existence predicate that requires some additional consideration when effecting a normalization. The present paper's emphasis on considering the consequences of additional connectives—like fusion and t —for the technique of normal models makes this an especially attractive area to explore.

Finally, as a referee has kindly suggested to us, there appears to be some kinship between Slaney's work on reduced models for weak relevant logics in Slaney [32] and the techniques employed in this paper. Slaney's method does not require that a logic has the principle of excluded middle, and so extending his results to first-order relevant logics is a natural step to expand on the results of this paper. We plan on revisiting the connections between normalization and Slaney's results, as well as exploring the metavaluation method, for first-order relevant logics in future work.

§5. Appendix: γ -Admissibility for propositional logics with fusion and t . In this appendix, we prove Theorem 2.4 which states that, for any conventionally normal logic **L** admitting γ , the logic **L** ^{$\circ t$} also admits γ . The proof will largely follow Seki's presentation in Seki [29] (minus modalities but tending to the cases of \circ and t).

LEMMA 5.21. *The normalization of an **L** ^{$\circ t$} frame is an **L** ^{$\circ t$} frame.*

Proof. Let $\mathfrak{F}' = \langle W', N', R', *' \rangle$ be the normalization of an **L** ^{$\circ t$} frame $\mathfrak{F} = \langle W, N, R, * \rangle$. That $'$ is a well-defined unary function is immediate; that R' is a well-defined ternary relation follows from a similar argument to that found in Routley et al. [27, p. 387]. \square

In particular, we note the following corollaries follow from Lemma 5.21

COROLLARY 5.22. *In a normalization \mathfrak{F}' based on \mathfrak{F} , the following hold for all $a, b \in W$*

$$a \leq' b \text{ iff } a \leq b \quad 0 \leq' b \text{ iff } 0 \leq b.$$

COROLLARY 5.23. *Let $\mathfrak{F}' = \langle W', N', R', *' \rangle$ be the normalization of an **L** ^{$\circ t$} frame $\mathfrak{F} = \langle W, N, R, * \rangle$ at 0 for $o \in W$. Then $o*' \leq' 0 \leq' o$.*

From the soundness of **L** ^{$\circ t$} , the following follows immediately.

LEMMA 5.24. *If \mathcal{A} is provable in **L** ^{$\circ t$} then \mathcal{A} is valid in every normal **L** ^{$\circ t$} frame.*

At this point, we lift the notion of normalization of a frame to define the normalization of a model.

DEFINITION 5.25. Let $\langle \mathfrak{F}, \|\cdot\| \rangle$ be an $\mathbf{L}^{\circ t}$ model. Then let its normalization be $\langle \mathfrak{F}', \|\cdot\|' \rangle$ where \mathfrak{F}' is the normalization of \mathfrak{F} and $\|\cdot\|'$ is defined so that for atoms p :

$$\|p\|' = \begin{cases} \|p\| & \text{if } o \notin \|p\| \\ \|p\| \cup \{0\} & \text{if } o \in \|p\|. \end{cases}$$

Now, we introduce two lemmata establishing that the normalization of a model will enjoy appropriate properties.

LEMMA 5.26. In a normalized $\mathbf{L}^{\circ t}$ model, $\|p\|'$ is an upset with respect to \leq' for every atom p .

Proof. By appeal to the model of which the model is a normalization. If $o \notin \|p\|$, then this follows. Otherwise, $\|p\|' = \|p\| \cup \{0\}$. Take an arbitrary $a \in \|p\|'$ and b such that $a \leq' b$. We prove that $b \in \|p\|'$.

Note again that both $o \in \|p\|$ and $0 \in \|p\|'$. If $b = 0$, the result is immediate, so assume that $b \in W$. Now, if $a = 0$, because $0 \leq' b$ holds precisely when $o \leq b$ holds in the original model. If $a \neq 0$, then because $b \neq 0$, $a \leq' b$ holds when $a \leq b$ holds in the original model. In both cases, as $\|p\|$ is an upset, $b \in \|p\|$ and *a fortiori* $b \in \|p\|'$ \square

LEMMA 5.27. In a normalized $\mathbf{L}^{\circ t}$ model, $\|\mathcal{A}\|'$ is an upset with respect to \leq' for every formula \mathcal{A} .

Proof. As $\langle \mathfrak{F}, \|\cdot\|' \rangle$ is a model, closure of propositions from the basis of Lemma 5.26 follows through a standard inductive argument. \square

This brings us to the key fact, namely, that a normalized model will agree with the model from which it was constructed. We ensure especially that the semantic clauses for \circ and t do not somehow prove problematic.

LEMMA 5.28. In a normalized $\mathbf{L}^{\circ t}$ model and point $a \in W$, $a \in \|\mathcal{A}\|$ iff $a \in \|\mathcal{A}\|'$.

Proof. By induction on complexity of \mathcal{A} . The basis step is provided by definition of the normalized model and most cases are covered as in Routley et al. [27, p. 391]. Given our interest in t and \circ , we provide these steps:

- If $\mathcal{A} = t$ then $\|t\|' = \|t\| \cup \{0\}$, so this holds for every $a \in W$.
- If $\mathcal{A} = \mathcal{B} \circ \mathcal{C}$ then to show left-to-right, pick an $a \in \|\mathcal{B} \circ \mathcal{C}\|$; this holds precisely when $b \in \|\mathcal{B}\|$ and $c \in \|\mathcal{C}\|$ such that $Rbca$. By induction hypothesis, this entails that $b \in \|\mathcal{B}\|'$ and $c \in \|\mathcal{C}\|'$. Additionally, by construction of the normalized frame, $R'bca$ holds, whence $a \in \|\mathcal{B} \circ \mathcal{C}\|'$. For right-to-left, on the other hand, pick an $a \in \|\mathcal{B} \circ \mathcal{C}\|'$ such that $a \in W$. Then there are $b \in \|\mathcal{B}\|'$ and $c \in \|\mathcal{C}\|'$ such that $R'bca$. Now, fix the following:

$$b' = \begin{cases} b & \text{if } b \neq 0 \\ o & \text{if } b = 0 \end{cases} \quad \text{and } c' = \begin{cases} c & \text{if } c \neq 0 \\ o & \text{if } c = 0. \end{cases}$$

Notably, b' and c' are elements of W . Then either trivially or by construction of R' , we have that $Rb'c'a$ and either trivially or by construction of \leq' , also $b \leq b'$ and $c \leq c'$, whence by Corollary 5.27, $b' \in \|\mathcal{B}\|'$ and $c' \in \|\mathcal{C}\|'$. In other words, there are $b' \in \|\mathcal{B}\|$ and $c' \in \|\mathcal{C}\|$ such that $Rb'c'a$, i.e., $a \in \|\mathcal{B} \circ \mathcal{C}\|$. \square

One final lemma will suffice to prove Theorem 2.4

LEMMA 5.29. $\mathbf{L}^{\circ t}$ proves \mathcal{A} iff \mathcal{A} is valid in every normal $\mathbf{L}^{\circ t}$ frame.

Proof. We prove the right-to-left direction via contraposition. Suppose that \mathcal{A} is not provable in $\mathbf{L}^{\circ f}$. Then there exists a model $\langle \mathfrak{F}, \|\cdot\| \rangle$ and a normal point $o \in N$ such that $o \notin \|\mathcal{A}\|$. By Lemma 5.28, in the normalization $\langle \mathfrak{F}, \|\cdot\|' \rangle$, $o \notin \|\mathcal{A}\|'$. By Corollary 5.23, $0 \leq' 0$, entailing that $0 \notin \|\mathcal{A}\|'$. Because $0 \in N'$, \mathfrak{F}' witnesses that \mathcal{A} is not valid in every normal $\mathbf{L}^{\circ f}$ frame. \square

This brings us to the main theorem concerning $\mathbf{L}^{\circ f}$.

THEOREM 2.4. γ is admissible for $\mathbf{L}^{\circ f}$.

Proof. Suppose for contradiction that γ fails, i.e., that there are $\mathbf{L}^{\circ f}$ theorems $\neg\mathcal{A} \vee \mathcal{B}$ and \mathcal{A} such that \mathcal{B} is not provable. By Lemma 5.29, there exists an $\mathbf{L}^{\circ f}$ model $\langle \mathfrak{F}', \|\cdot\|' \rangle$ with \mathfrak{F}' normal such that $0 \notin \|\mathcal{B}\|'$. As $0 \in N'$, however, $0 \in \|\neg\mathcal{A} \vee \mathcal{B}\|'$ and $0 \in \|\mathcal{A}\|'$. Consequently, either $0 \in \|\neg\mathcal{A}\|'$ or $0 \in \|\mathcal{B}\|'$. Because $0 = 0^{*'}$, the requirement of the former case that $0^{*'} \notin \|\mathcal{A}\|'$ translates to $0 \notin \|\mathcal{A}\|'$, contradicting the hypothesis that $0 \in \|\mathcal{A}\|'$. The latter case is ruled out insofar as 0 was assumed not to be a member of $\|\mathcal{B}\|'$. As both disjuncts lead to contradiction, we conclude that γ holds. \square

Acknowledgments. We would like to thank many comments from participants of several conferences, including Logica 2023, the Second Third Workshop, and the Seminar for Applied Mathematical Logic at the Institute of Computer Science, Czech Academy of Science. We also thank three anonymous reviewers for their comments, which lead to a much better version of the paper.

Funding. Nicholas Ferenz gratefully acknowledged funding from by RVO 67985807 and that this work is also financed by national funds through FCT — Fundação para a Ciência e a Tecnologia, I.P., under the scope of UIDB/00310/2020 project, identified as DOI 10.54499/UIDB/00310/2020.

BIBLIOGRAPHY

- [1] Ackermann, W. (1956). Begründung einer Strengen Implikation. *The Journal of Symbolic Logic*, **21**, 113–128.
- [2] Bimbó, K. (2007). Relevance logics. In Jacquette, D., editor, *Philosophy of Logic*, vol 5. of Handbook of the Philosophy of Science (D. Gabbay, P. Thagard and J. Woods, eds.). North Holland: Elsevier, pp. 723–789.
- [3] Brady, R. (1983). The Non-Triviality of Dialectical Set Theory. In Priest, G., Routley, R., and Norman, J., editors, *Paraconsistent Logics: Essays in the Inconsistent*. Munich: Philosophia Verlag, pp. 437–471.
- [4] ———. (1988). A content semantics for quantified relevant logics I. *Studia Logica*, **47**, 111–127.
- [5] ———. (1989). A content semantics for quantified relevant logics II. *Studia Logica*, **48**, 243–257.
- [6] Dunn, J. M., & Restall, G. (2002). Relevance logic. In Gabbay, D. M., and Guenther, F., editors, *Handbook of Philosophical Logic* (second edition). Netherlands: Springer, pp. 1–128.
- [7] Ferenz, N. (2022). Conditional **FDE** logics. In Bimbó, K., editor, *Relevance Logics and Other Tools for Reasoning*. Essays in Honour of J. Michael Dunn, London: College Publications, pp. 182–214.

- [8] ———. (2023). Quantified modal relevant logics. *Review of Symbolic Logic*, **16**, 210–240.
- [9] Ferenz, N., & Ferguson, T. M. (2024). γ -admissibility for Mares' varying domain **QR**. In Sedlár, I., editor, *The Logica Yearbook 2023*, London: College Publications. To appear.
- [10] Ferenz, N., & Tedder, A. (2023). Neighbourhood semantics for modal relevant logics. *Journal of Philosophical Logic*, **52**, 145–181.
- [11] Fine, K. (1988). Semantics for quantified relevance logics. *Journal of Philosophical Logic*, **17**, 22–59.
- [12] ———. (1989). Incompleteness for quantified relevant logics. In Norman, J. & Sylvan, R., editors. *Directions in Relevant Logic*. New York: Kluwer Academic Publishers, pp. 205–225, Reprinted in *Entailment* vol. 2, §52, Anderson, Belnap, and Dunn, 1992.
- [13] Friedman, H., & Meyer, R. K. (1992). Whither relevant arithmetic? *Journal of Symbolic Logic*, **57**(3), 824–831.
- [14] Humberstone, L. (2011). *The Connectives*. Cambridge: MA.
- [15] Kripke, S. (2022). A proof of gamma. In Bimbó, K., editor. *Relevance Logics and Other Tools for Reasoning. Essays in Honor of J. Michael Dunn*, London: College Publications, pp. 261–265.
- [16] Mares, E., & Goldblatt, R. (2006). An alternative semantics for quantified relevant logic. *The Journal of Symbolic Logic*, **71**, 163–187.
- [17] Mares, E., & Meyer, R. K. (1992). The admissibility of γ in **R4**. *Notre Dame Journal of Formal Logic*, **33**, 197–206.
- [18] Mares, E. D. (2009). General information in relevant logic. *Synthese*, **167**(2), 421–440.
- [19] Meyer, R. K. (1976). Metacompleteness. *Notre Dame Journal of Formal Logic*, **17**, 501–516.
- [20] Meyer, R. K., & Dunn, J. M. (1969). E, R, and γ . *Journal of Symbolic Logic*, **34**, 460–474.
- [21] Meyer, R. K., Dunn, J. M., & Leblanc, H. (1974). Completeness of relevant quantification theories. *Notre Dame Journal of Formal Logic*, **15**, 97–121.
- [22] Meyer, R. K., Giambrone, S., & Brady, R. (1984). Where gamma fails. *Studia Logica*, **43**, 247–256.
- [23] Pudlák, P. (1998). The lengths of proofs. In Buss, S. R., editor. *Handbook of Proof Theory*. Amsterdam: North Holland, pp. 547–637.
- [24] Raftery, J. G., & Świrydowicz, K. (2016). Structural completeness in relevance logics. *Studia Logica*, **104**, 381–387.
- [25] Routley, R., & Meyer, R. K. (1972). The semantics of entailment (2). *The Journal of Philosophical Logic*, **1**, 53–73.
- [26] ———. (1973). The semantics of entailment (1). In Leblanc, H., editor. *Truth, Syntax, and Modality*. Amsterdam: North-Holland, pp. 199–243.
- [27] Routley, R., Plumwood, V., Meyer, R. K., & Brady, R. T. (1982). *Relevant Logics and Their Rivals: Part 1 The basic Philosophical and Semantical Theory*. Ridgeview: Atascadero, CA.
- [28] Seki, T. (2003). General frames for relevant modal logics. *Notre Dame Journal of Formal Logic*, **44**, 93–109.
- [29] ———. (2011a). The γ -admissibility of relevant modal logics I—the method of normal models. *Studia Logica*, **97**, 199–231.

- [30] ———. (2011b). The γ -admissibility of relevant modal logics II—the method using metavaluations. *Studia Logica*, **97**, 351–383.
- [31] ———. (2012). An algebraic proof of the admissibility of γ in relevant modal logics. *Studia Logica*, **100**, 1149–1174.
- [32] Slaney, J. K. (1987). Reduced models for relevant logics without WI. *Notre Dame Journal of Formal Logic*, **28**(3), 395–407.
- [33] Smiley, T. (1973). Relative necessity. *Journal of Symbolic Logic*, **28**, 113–134.
- [34] Tedder, A., & Ferenz, N. (2022). Neighbourhood semantics for quantified relevant logics. *Journal of Philosophical Logic*, **51**, 457–484.
- [35] Urquhart, A. (2016). The story of γ . In Bimbo, K., editor, *J. Michael Dunn on Information Based Logics* (Outstanding Contributions to Logic, vol. 8). Cham: Springer, pp. 93–106.
- [36] Weiss, Y. (2020). Cut and gamma I: Propositional and constant domain **R**. *The Review of Symbolic Logic*, **13**, 887–909.

CENTER OF PHILOSOPHY, SCHOOL OF ARTS AND HUMANITIES
 UNIVERSITY OF LISBON
 CIDADE UNIVERSITÁRIA, ALAMEDA DA UNIVERSIDADE
 1649-004, LISBON
 PORTUGAL

E-mail: nicholas.ferenz@gmail.com

DEPARTMENT OF COGNITIVE SCIENCE
 RENSSELAER POLYTECHNIC INSTITUTE
 110 8TH STREET, TROY, NY 12180
 USA

E-mail: tferguson@gradcenter.cuny.edu