

**Conclusions:** Psychoactive substance use has become a growing problem among residents in Tunisia. The associated factors should attract attention to identify these subjects.

**Disclosure:** No significant relationships.

**Keywords:** substance use; medical residents; Tunisia

## Personality Disorders / Depressive Disorders 05

### EPP0779

#### Psychopathological and clinical-typological aspects of youth chronic endogenous depression.

V. Kaleda\* and V. Migalina

FSBSI «Mental Health Research Centre», Department Of Youth Psychiatry, Moscow, Russian Federation

\*Corresponding author.

doi: 10.1192/j.eurpsy.2022.940

**Introduction:** Youth ontogenesis contributes significantly to depressive disorders, causing pronounced atypia, a high level of comorbid pathology. A long-term depressive state lead can to persistent, adverse consequences.

**Objectives:** To study the clinical, psychopathological and psychometric features of youth chronic endogenous depression (UCED).

**Methods:** 62 patients of the age 16-25 were examined clinically and psychopathologically; the patients were first hospitalized from 2017 to 2020 for a chronic depressive state with non-psychotic mental disorders (ICD-10: F31, F32, F33, F34, F21 keys) lasting more than two years. Psychometric assessment was done by HDRS, SOPS, and SANS.

**Results:** UCED are characterized by a pronounced atypia with a predominance of symptoms for negative affectivity with apathy, anhedonia, physical and mental asthenia, depressive devitalization. In contrast with non-chronic youth depressions, cognitive disorders, motor inhibition, a large proportion of comorbid pathology are presented in the chronic ones. Depending on the prevalence of additional psychopathological disorders, 2 types were distinguished: Type I – depression with a clear-cut affective psychopathological structure (54.8%, 34 patients); Type II - depression with the symptoms of other than affective registers (45.2%, 28 patients). Psychometric assessment on the HDRS scale, in the sub-scale “negative symptoms” of the SOPS scale, in the sub-scale “anhedonia-associality” of the SANS scale showed a greater severity of psychopathological symptoms in type II depression ( $p < 0.05$ ).

**Conclusions:** The obtained data confirm the differences between UCED and non-chronic youth depressions and demonstrate the aggravating effect of symptoms of the non-affective spectrum on the severity of UCED and the level of negative affectivity.

**Disclosure:** No significant relationships.

**Keywords:** chronic depression; youth; persistent depressive; dysthymia

### EPP0780

#### Resistance or pseudo-resistance?

R. Zanardi<sup>1</sup>, F. Attanasio<sup>2\*</sup>, C. De Cesare<sup>2</sup>, V. Fazio<sup>2</sup> and C. Colombo<sup>1,2</sup>

<sup>1</sup>IRCCS San Raffaele Scientific Institute, Psychiatry - Mood Disorders, Milano, Italy and <sup>2</sup>Università Vita-Salute San Raffaele, Psychiatry, Milano, Italy

\*Corresponding author.

doi: 10.1192/j.eurpsy.2022.941

**Introduction:** Treatment-Resistant Depression continues to represent a great challenge for clinicians.

**Objectives:** We investigated patients with history of resistance, assessing prognostic factors, response to treatments, and remission over time.

**Methods:** We recruited 202 unipolar and bipolar depressed inpatients. According to anamnestic backgrounds, patients were assigned to: A) *Non-resistant*: responders, with no characteristics of resistance in the current episode. B) *Resistant*: resistant to two antidepressant trials of adequate doses and duration. C) *Pseudo-resistant*: non-responders, not classifiable as *Resistant* because of inadequate trials. During hospitalization, patients were treated by clinical judgment, following a rehabilitation program.

**Results:**

