

It became impossible, so by and by Jean and he were married”.

The sudden onset of blindness in the war and his recovery on the passage home are suggestive of combat hysteria. It is likely that the symptom of blindness was an unconscious attempt to 'shut out' the sights of mutilation he had seen while assisting the Army Surgeon.

M. A. PATTON

*Department of Medicine,
University Hospital of Wales,
Cardiff CF4 4XW*

References

- HURST, A. F. (1940) *Medical Diseases of War*. London: Edward Arnold.
MOTT, F. W. (1919) *War Neuroses and Shell Shock*. London: Oxford Medical Publications.
YEALLAND, L. R. (1918) *Hysterical Disorders of Warfare*. London: Macmillan.

ANTICIPATORY GRIEF

DEAR SIR,

Fulton and Gottesman's (*Journal*, July 1980, 137, 45-54) detailed analysis of the literature on Anticipatory Grief rightly raises doubts about the usefulness of this concept. Unfortunately they have failed to spot a very interesting and important finding which emerges from the large amount of research which they report.

- Despite all the differences in definition and methodology there are three studies which come up with clear evidence of a relationship between forewarning of bereavement and good outcome. The factor which distinguishes these three from the five other quantitative studies is young age. Natterson and Knudson's (1960) study of dying children and Glick's (1974) study of widows and widowers under age 45 both show that in the young age group, the opportunity to anticipate bereavement does reduce the intensity of subsequent distress. Ball's (1977) study allows direct comparison of widows over and under 45 years of age and confirms this finding.

Although Maddison and Walker's (1967) study is referred to by Fulton as being concerned with "young and middle aged" widows in fact widows under 45 were not included in their research.

One other study which was not included by Fulton is a comparison of relatives of 32 patients who had died in Sweden within a few hours of becoming ill and 55 controls most of whom had died from cancer (Lundin, 1979). Both groups were under the age of 65 and the change in morbidity was measured in days of sick-listing from Swedish social insurance claims during 2 years before and after the death. Lundin

found that the mean number of days sick increased from 41.2 to 54.0 in the unexpectedly bereaved group and that this was most marked in relatives of patients who had died in a violent accident ($P < .01$). No such increase was found in the control group although both before and after bereavement morbidity was higher than in the normal population (suggesting that relatives of patients with cancer may suffer stress-effects before as well as after bereavement).

It seems reasonable to conclude that deaths which are both unexpected and untimely are potentially pathogenic.

C. COLIN MURRAY PARKES

*The London Hospital Medical College
Turner Street, London E1 2AD*

References

- BALL, J. F. (1977) Widow's grief: the impact of age and mode of death. *Omega*, 7, 307-33.
GLICK, I. O., WEISS, R. S. & PARKES, C. M. (1974) *The First Year of Bereavement*. New York: John Wiley.
LUNDIN, T. (1979) *On Crisis Theory, Reactions to Sudden and Unexpected Death*. Proceedings of 10th International Conference for Suicide Prevention and Crisis Intervention. Ottawa, Canada.
MADDISON & WALKER, W. L. (1967) Factors affecting the outcome of conjugal bereavement. *British Journal of Psychiatry*, 113, 1057-67.
NATTERSON, J. M. & KNUDSON, A. G. JR. (1960) Observations concerning fear of death in fatally ill children and their mothers. *Psychosomatic Medicine*, 22, 465.

VULNERABILITY FACTORS AND DEPRESSION

DEAR SIR,

Roy (*Journal*, August 1978, 133, 106 and January 1981, 138, 75) purports to corroborate Brown *et al's* (1975, 1977) findings that unemployment, lack of a confiding relationship and early parental loss act as vulnerability factors in relation to depression. This conclusion, would not appear to follow from his data.

The features that constitute the essential defining characteristics of vulnerability factors have been hotly disputed (Tennant and Bebbington, 1978; Brown and Harris, 1978, 1980; Bebbington, 1980; Surtees, 1980). Nevertheless, a fundamental point of agreement is that their effects can only be demonstrated in conjunction with information pertaining to life events. Roy (1978) appears to concur with this view in that he defines vulnerability factors as "factors which increase the chances of developing a psychiatric disorder in the presence of an event or difficulty" (p 106).

Despite this definition, however, he failed to collect any information pertaining to the occurrence of life events or difficulties. Thus, in the absence of this information it is not possible to determine whether