

## OBESITY, SELF-ESTEEM AND SEXUALITY: A COMPARATIVE STUDY ABOUT 40 CASES

**J. Ben Thabet**<sup>1</sup>, **R. Sallemi**<sup>1</sup>, **M. Maâlej Bouali**<sup>1</sup>, **N. Charfi**<sup>2</sup>, **N. Charfi**<sup>1</sup>, **L. Zouari**<sup>1</sup>, **N. Zouari**<sup>1</sup>, **M. Abid**<sup>2</sup>, **M. Maâlej**<sup>1</sup>

<sup>1</sup>Psychiatry C, <sup>2</sup>Endocrinology, Hedi Chaker Hospital, Sfax, Tunisia

**Aim:** To assess the impact of self-esteem on the sexuality of obese patients compared to a sample from the general population.

**Patients and methods:** A cross-sectional case-control study. It involved 40 obese married patients (BMI  $\geq 30$  kg/m<sup>2</sup>) and 40 normal weight subjects from the general population. A questionnaire was presented with sociodemographic data, somatic and psychiatric personal histories, data on obesity and sexual life data. Self-esteem was assessed using the Rosenberg Self Esteem Scale (RSES).

**Results:** The sex-ratio (men/women) of obese patients was 0.53. The average BMI of obese patients was 33.48 kg/m<sup>2</sup> (SD = 3.44). The score at the RSES in the obese group was 30.15 (SD = 5.48) versus 32.2 (SD = 4.58) in the control group, with no significant difference ( $p = 0.808$ ). Among obese patients, the higher the score of self-esteem was, the more frequent sexual intercourses were, without significant difference ( $r = 0.254$ ,  $p = 0.119$ ). Sexual satisfaction was significantly correlated with the level of self-esteem ( $p = 0.022$ ). Obese people with low self-esteem were more likely to have a decrease in sexual desire (OR = 9.625, CI [1.378, 67.246]).

**Conclusion:** There is a link between low self-esteem of obese and an alteration of the quality of their sexual life. The therapist should check the level of self-esteem in obese patients and try to improve it. In this sense, the collaboration of the endocrinologist with the psychiatrist could be very useful.