

This is a “preproof” accepted article for Weed Science. This version may be subject to change in the production process, *and does not include access to supplementary material*.

DOI: 10.1017/wsc.2024.100

**Short title:** Cover crop in CGP

**Effect of fall-planted cover crops on weed suppression, grain sorghum yield, and profitability in the semi-arid Central Great Plains**

Sachin Dhand<sup>1</sup>, Vipin Kumar<sup>2</sup>, J. Anita Dille<sup>3</sup>, Augustine Obour<sup>4</sup>, Elizabeth A. Yeager<sup>5</sup>, and Johnathan Holman<sup>6</sup>

<sup>1</sup>Graduate Research Assistant, Kansas State University, Agricultural Research Center, Hays, KS, USA; <sup>2</sup>Associate Professor, Cornell University, School of Integrative Plant Science, Soil and Crop Sciences Section, Ithaca, NY, USA; <sup>3</sup>Professor, Kansas State University, Department of Agronomy, Manhattan, KS, USA; <sup>4</sup>Professor, Kansas State University, Agricultural Research Center, Hays, KS, USA; <sup>5</sup>Associate Professor, Kansas State University, Department of Agricultural Economics, Manhattan, KS, USA; <sup>6</sup>Professor, Kansas State University, Southwest Research and Extension Center, Garden City, KS, USA

**Author for correspondence:** Vipin Kumar, Associate Professor (ORCID: 0000-0002-8301-5878), Cornell University, School of Integrative Plant Science, Soil and Crop Sciences Section, 1115 Bradfield Hall, Ithaca, NY 14853. **Email:** [yk364@cornell.edu](mailto:yk364@cornell.edu)

## Abstract

Integrating cover crops (CC) in dryland crop rotations could help in controlling herbicide-resistant weeds. Field experiments were conducted at Kansas State University Agricultural Research Center near Hays, KS from 2020-2023 to determine the effect of fall-planted CC on weed suppression in grain sorghum, crop yield, and net returns in no-till dryland winter wheat (*Triticum aestivum* L.)-grain sorghum [*Sorghum bicolor* (L.) Moench]-fallow (W-S-F) rotation. The field site had a natural seedbank of glyphosate-resistant (GR) kochia [*Bassia scoparia* (L.) A. J. Scott] and Palmer amaranth (*Amaranthus palmeri* S. Watson). A CC mixture (triticale/winter peas/radish/rapeseed) was planted after wheat harvest and terminated at triticale heading stage before sorghum planting. Treatments included nontreated control, chemical fallow, CC terminated with glyphosate (GLY), and CC terminated with GLY+ acetochlor/atrazine (ACR/ATZ). Across three years, CC terminated with GLY+ACR/ATZ reduced total weed density by 34-81% and total weed biomass by 45-73% compared to chemical fallow during the sorghum growing season. Average grain sorghum yield was 786 to 1432 kg ha<sup>-1</sup> and did not differ between chemical fallow and CC terminated with GLY+ACR/ATZ. However, net returns were lower with both CC treatments (USD -\$275 to \$66) in all three years compared to chemical fallow (USD -\$111 to \$120). These results suggest that fallow replacement with fall-planted CC in the W-S-F rotation can help suppress GR *B. scoparia* and *A. palmeri* in the subsequent grain sorghum. However, the cost of integrating CC exceeded the benefits of improved weed control and lower net returns were recorded in all three years compared to chemical fallow.

**Keywords:** Cover crop; dryland; kochia; Palmer amaranth; residual herbicide

## Introduction

The Central Great Plains (CGP) is characterized by a semi-arid climate with relatively low annual precipitation (~300 to ~1200 mm) (Lenssen et al. 2007; NOAA 2024). To conserve soil moisture and to prevent soil erosion by wind, no-tillage (NT-) and fallow-based cropping systems are widely adopted in the region. Successful adoption of these soil conservation practices was achieved utilizing chemical-based weed control (Hansen et al. 2012; Kumar et al. 2020). However, the adoption of NT-based production systems has resulted in weed species representing smaller-seeded weeds like kochia [*Bassia scoparia* (L.) A.J. Scott], Palmer amaranth (*Amaranthus palmeri* S. Watson), horseweed [*Conyza canadensis* (L.) Cronquist; syn. *Erigeron canadensis* (L.) Cronquist], common lambsquarters (*Chenopodium album* L.), Russian thistle (*Salsola tragus* L.), downy brome (*Bromus tectorum* L.), wild oat (*Avena fatua* L.), foxtail species (*Setaria* spp.), and tumble windmill grass (*Chloris verticillata* Nutt.) (Jha et al. 2016; Nichols et al. 2015).

Winter wheat (*Triticum aestivum* L.)-grain sorghum [*Sorghum bicolor* (L.) Moench]-fallow (W-S-F) is a dominant crop rotation in the CGP region (Holman et al. 2022). This 3-year crop rotation includes a fallow period of approximately 10 months between winter wheat harvest and sorghum planting as well as 10 months of fallow period between sorghum harvest and the next winter wheat planting (Kumar et al. 2020). Continuous reliance on herbicides with the same site(s) of action for weed control has resulted in the evolution of herbicide-resistant weeds, including *B. scoparia*, *A. palmeri*, and *C. canadensis* (Heap 2024). For instance, glyphosate resistance is widespread among *B. scoparia* and *A. palmeri* populations in Kansas and other neighboring states in the CGP region (Heap 2024; Kumar et al. 2019a, 2019b, 2020; Westra et al. 2019). Evolution of glyphosate-resistant (GR) weed populations and limited availability of alternative effective herbicide options pose a serious production challenge for grain sorghum producers in the region. Previous researchers have reported that season-long weed interference can result in an average grain yield loss of 47% in sorghum, which is an estimated loss of around US \$953 million annually (Dille et al. 2020). Therefore, alternative integrated weed management strategies are needed to achieve effective control of herbicide-resistant weed populations in grain sorghum.

Integration of cover crop (CC) in crop rotations has been proven as one of the effective tools to suppress herbicide-resistant weeds in the CGP region (Kumar et al. 2020; Mesbah et al. 2019;

Obour et al. 2022a; Petrosino et al. 2015). Growing CC in the semi-arid CGP also provides several other benefits, including reduced soil erosion, enhanced nutrient cycling, increased microbial activity, improved soil health, and increased plant diversity and pollinator resources (Blanco-Canqui et al. 2011, 2013; Simon et al. 2022). Additionally, CC residue left on the soil surface after termination reduces soil temperature and soil moisture evaporation, thereby contributing to increased soil water storage (Holman et al. 2020, 2021). However, replacing the fallow period with CC in the semi-arid cropping systems sometimes reduces the yield of successive crops because of the reduced plant available water (Holman et al. 2018; Nielsen et al. 2016). However, the United States Department of Agriculture Natural Resources Conservation Service (USDA NRCS) provides some financial support to growers under the Environmental Quality Incentives Program (EQIP) to pay some of the cost of growing CC and improve net returns (Anonymous 2024). Previous studies have evaluated the effect of spring-planted CC on weed suppression and winter wheat yields when CCs replaced the fallow phase of W-S-F rotation in this region (Holman et al. 2022; Mesbah et al. 2019; Obour et al. 2022a). For instance, Obour et al. (2022a) reported that spring-planted CC (oats/triticale/spring peas) in W-S-F rotation can reduce weed biomass by 86 to 99% compared to weedy fallow. Holman et al. (2022) reported that spring-planted CC had no significant effect on wheat and grain sorghum yields when conditions were either extremely dry with poor yields or very wet with above-average yields, however, replacing fallow with CC increased the cost of production by 16 to 97% compared to fallow.

Farmers are currently relying on residual herbicides to manage GR weeds in the dryland W-S-F rotation (Kumar et al. 2020). Several researchers have previously documented the importance of residual herbicides in combination with CC to achieve season-long weed control (Perkins et al. 2021; Whalen et al. 2020). For instance, Whalen et al. (2020) reported that CC terminated with glyphosate plus 2,4-D in combination with residual herbicides (sulfentrazone plus chlorimuron) resulted in greater waterhemp [*Amaranthus tuberculatus* (Moq.) Sauer.] control (73 to 84%) compared to no residual herbicide (44 to 65%). Most CC weed suppression research studies were conducted in cotton (*Gossypium hirsutum* L.), corn (*Zea mays* L.), or soybean [*Glycine max* (L.) Merr.] in greater precipitation environments (Weisberger et al. 2023; Whalen et al. 2020). However, limited information exists regarding the integration of fall-planted CC in combination with soil residual herbicides at the termination of CC on weed suppression in

subsequent grain sorghum in the semi-arid CGP region. The main objectives of this study were to determine the effect of fall-planted CC in combination with soil residual herbicides on (1) weed suppression (density and biomass) in subsequent grain sorghum and grain yield, and (2) net returns with integrating CC in the no-till dryland W-S-F cropping system. We hypothesized that fall-planted CC, combined with residual herbicides, would provide adequate weed suppression in grain sorghum with minimal or no impact on sorghum yield, resulting in higher net returns.

## Materials and Methods

### Experiment design and treatments

A field experiment was conducted at Kansas State University Agricultural Research Center near Hays (KSU-ARCH), KS (38.85196°N, 99.34279°W) during 2020-2021, 2021-2022, and 2022-2023 growing seasons. The experiment was initiated in the fall of 2020. The soil type at the experimental site was a Roxbury silt loam with a pH of 6.9 and organic matter of 1.6%. The study site was under no-till W-S-F rotation for > 10 years prior to study initiation and had a natural uniform seedbank of GR *B. scoparia* and *A. palmeri* (Kumar, personal observations). All three phases of the W-S-F crop rotation were present each year. After wheat harvest, all plots were sprayed in late July with glyphosate (GLY) (Roundup PowerMax<sup>®</sup>, Bayer Crop Science, St. Louis, MO, USA) at 1260 g ae ha<sup>-1</sup> plus dicamba (Clarity<sup>®</sup>, BASF Corporation, Research Triangle Park, NC, USA) at 560 g ae ha<sup>-1</sup>. A CC mixture of winter triticale (× *Triticosecale* Wittm.) (60%)/winter peas (*Pisum sativum* L.) (30%)/rapeseed (*Brassica napus* L.) (5%)/radish (*Raphanus sativus* L.) (5%) was drill seeded into wheat stubble at a rate of 67 kg ha<sup>-1</sup> during each fall and terminated in the following spring at the triticale heading stage. The CC planting dates were September 28, October 7, and September 30 in 2020, 2021, and 2022, respectively. The CC was terminated on May 13, May 11, and May 22 in 2021, 2022, and 2023, respectively. During each spring, four treatments were established: (1) weedy fallow, (2) chemical fallow, (3) CC terminated with GLY alone, and (4) CC terminated with GLY + residual herbicide. In weedy fallow treatment, no CC was planted and no herbicides were applied to control weeds. In chemical fallow treatment, no CC was planted but the plot area was treated with GLY at 1260 g ae ha<sup>-1</sup> plus a premix of acetochlor/atrazine (ACR/ATZ) (Degree Xtra<sup>®</sup>, Bayer Crop Science, St. Louis, MO, USA) at 1665/826 g ai ha<sup>-1</sup> plus dicamba at 560 g ae ha<sup>-1</sup> at the same time as CC termination. For CC termination, GLY at 1260 g ae ha<sup>-1</sup> was used, and the residual herbicide was

a premix of ACR/ATZ at 1665/826 g ai ha<sup>-1</sup>. All treatments were established in a randomized complete block design with four replications. During 2020-2021, weedy fallow treatment was not present and there were only three treatments. The individual plot size was 45-m long and 13-m wide each year. During 2021-2022 and 2022-2023 experimental years, the initial chemical fallow plot was sub-divided into two to have both weedy fallow and chemical fallow treatments (45-m long and 6.5-m wide each) for comparison of weed suppression. A sorghum hybrid ‘DKS 38-16’ was planted at a seeding rate of 114,855 seeds ha<sup>-1</sup> in rows spaced 76-cm apart on June 9, June 2, and June 15 during 2021, 2022, and 2023, respectively. All local agronomic practices for grain sorghum production as recommended by Kansas State University were followed (Ciampitti et al. 2022a). No herbicides were applied in the grain sorghum growing season. Grain sorghum was harvested on November 4, October 26, and October 19 during 2021, 2022, and 2023, respectively. Data on monthly precipitation and air temperature over the 3-year study period were obtained from the Kansas State University Mesonet weather station (<https://mesonet.k-state.edu>) located approximately 400 m from the study site (38.8495°N, 99.3446°W) (Figure 1).

### **Cover crop biomass, weed density, and weed biomass**

Each year, the aboveground shoot biomass of CC was manually harvested from two 1-m<sup>2</sup> quadrats from each plot just before CC termination and oven-dried at 72 C for 4 days to obtain dry biomass. Weed density by species (number of emerged seedlings for each species) was recorded from two randomly placed 1-m<sup>2</sup> quadrats from each plot at CC termination and at monthly intervals until sorghum harvest (except 2021, where data at 90 days after CC termination was not collected) and aboveground weed biomass was manually harvested and oven-dried at 72 C for 4 days to obtain total weed dry biomass. The average of CC biomass, total weed density, and total weed dry biomass from the two quadrats in each plot at each time were used in the data analysis. The weed species composition was characterized by calculating the relative abundance of each species in each plot using the method described by Thomas (1985) and used by Obour et al. (2022a). Relative abundance was determined using equation 1. Relative density and relative frequency were calculated using equations 2 and 3, respectively.

$$\text{Relative abundance} = \frac{\text{Relative density} + \text{Relative frequency}}{2} \quad \text{Eqn. 1}$$

Relative density

$$= \left[ \frac{\text{Number of plants for each species within the quadrat per plot}}{\text{Total number of plants in that sampled quadrat}} \right] \times 100 \quad \text{Eqn. 2}$$

Relative frequency

$$= \left[ \frac{\text{Proportion of quadrats in which the species was present in a plot}}{\text{Frequency of all species in that sampled quadrat}} \right] \times 100 \quad \text{Eqn. 3}$$

### **Volumetric water content and grain sorghum yield**

The CC effect on soil water content at grain sorghum planting was determined gravimetrically in 30 cm increments up to 150 cm depth. Two soil cores were taken from each plot using a hydraulic probe (Giddings Machine Company) in June 2021 and 2023 before sorghum planting. During 2022, the soil samples were not collected. Soil sample portions from each 30-cm depth were weighed fresh and then dried at 105 C for 4 days to calculate bulk density by dividing the mass of oven-dry soil to the volume of the core. Gravimetric water content was calculated using equation 4.

$$\text{Gravimetric water content} = \left[ \frac{\text{Wet soil weight} - \text{Dry soil weight}}{\text{Dry soil weight}} \right] \quad \text{Eqn. 4}$$

Data from both soil cores were averaged to obtain a single soil water measurement that was converted to volumetric water content by multiplying with measured bulk density at each sampling depth. Data for volumetric water content were averaged for both CC treatments as both treatments were the same before termination. Grain sorghum yield was recorded by harvesting each whole plot using a Massey Ferguson 8XP small plot combine harvester (Massey Ferguson, Duluth, GA, USA) and was adjusted to 13.5% moisture content.

### **Economic analysis**

Gross returns were calculated by multiplying the grain sorghum yield and the price of sorghum grain. Net returns were calculated as gross returns minus total variable costs for each treatment for each year. Fixed costs were ignored in this analysis as they were assumed to be consistent

across treatments. Four-year average custom rate values published by Kansas State University Land Use Survey Program and the Kansas Department of Agriculture (AgManager 2022) were used for current field operations and input costs. Total variable cost was calculated by adding all the expenses for planting (CC and sorghum), inputs (fertilizer, herbicides, etc., and their application costs), and harvesting. Grain sorghum price for each experimental year was taken from the USDA Economic Research Services market reports (USDA ERS 2023). Prices for grain sorghum were calculated on a per kg basis and ranged from \$0.20 to \$0.24. All costs and revenue were calculated in U.S. dollars per hectare.

### **Statistical Analyses**

Data were tested for homogeneity of variance and normality of the residuals using the PROC UNIVARIATE procedure in SAS 9.3 (SAS Institute, Inc., SAS Campus Drive, Cary, NC). Data for total weed density and total dry biomass were log-transformed to improve the normality of the residuals and homogeneity of variance; however, back-transformed data were presented with mean separation based on the transformed data, whereas the rest of the data met both ANOVA assumptions. All data for CC biomass, total weed density, and total weed dry biomass at each time, volumetric water content, grain yield, and net returns were subjected to ANOVA using PROC MIXED procedure. For CC biomass data, year was considered as fixed effect and replication was considered as random effect. For total weed density and total weed dry biomass data, CC treatment, year, monthly timing, and their interactions were considered as fixed effects whereas replication and their interactions were considered as random effects. Repeated measures were accounted for monthly timing. For volumetric water content, CC treatment, year, soil depth, and their interactions were considered as fixed effects whereas replication and their interactions were considered as random effects. For data on grain sorghum yield and net returns, CC treatment, year, and their interactions were considered as fixed effects whereas replication and their interactions were considered as random effects. Data for total weed density, total weed dry biomass, volumetric water content, grain sorghum yield, and net returns were analyzed separately for each year because of significant year-by-treatment interaction ( $P < 0.01$ ). Treatment-by-monthly evaluation interaction for total weed density and total weed dry biomass was significant ( $P < 0.001$ ); therefore, data were sorted by monthly evaluation timings using PROC SORT with monthly evaluations treated as a repeated measure. Treatment means were



separated using Fisher's protected LSD test ( $P < 0.05$ ). The grain sorghum yields were low because of the drought conditions during the study period; therefore, a sensitivity analysis was conducted to obtain net returns to possible grain sorghum yield (700 to 7400 kg ha<sup>-1</sup>) and prices (\$0.09 to \$0.24 kg<sup>-1</sup>) in the region.

## Results and Discussion

Variable precipitation amount and frequency were observed at KSU-ARCH during the experimental periods 2020-2021, 2021-2022 and 2022-2023 (Figure 1). The total amount of precipitation received during the CC growing season (September to May) in 2020-2021, 2021-2022 and 2022-2023 was 217, 99, and 130 mm, respectively (Figure 1). The 30-year average precipitation from September to May in the region is 347 mm (Figure 1). No difference was recorded in aboveground CC dry biomass at the time of termination across the years and was 1520, 1130, and 1470 kg ha<sup>-1</sup> in 2021, 2022, and 2023, respectively, with an average of  $1370 \pm 123$  kg ha<sup>-1</sup>. During the sorghum growing season (June through October), the total precipitation amount was 256, 171, and 237 mm in 2021, 2022, and 2023, respectively (Figure 1). The 30-year average precipitation from June to October in the region is 341 mm (Figure 1).

### Total weed density and weed dry biomass

Across three years, four summer annual broadleaf weed species were observed at the study site, including *B. scoparia*, *A. palmeri*, Venice mallow (*Hibiscus trionum* L.), and puncturevine (*Tribulus terrestris* L.). Based on the relative abundance, *B. scoparia* and *A. palmeri* were the dominant weed species across three years (Tables 1, 3, and 4).

**2021 growing season.** *Amaranthus palmeri* was the most dominant weed species with a mean relative abundance of >40% across treatments at all monthly evaluation timings (Table 1). Prior to termination [0 days after termination (DAT)], CC reduced the total weed density by 86 to 95% compared to chemical fallow (no herbicide was applied in chemical fallow at this time) (Table 1). However, an application of GLY plus ACR/ATZ plus dicamba in chemical fallow at the time of CC termination reduced weed density at later evaluation timings. Total weed density at 60 DAT was dominated by *A. palmeri* (relative abundance = 90%) and was significantly greater (approximately 4 times) following the CC terminated with GLY only compared to the CC

terminated with GLY plus ACR/ATZ (Table 1). This would be due to a lack of residual herbicide and CC residue was not enough to suppress the emerging *A. palmeri* seedlings. At the time of grain sorghum harvest (120 DAT), CC terminated with GLY plus ACR/ATZ reduced the total weed density by 50% compared to chemical fallow (Table 1). Consistent with total weed density, CC at termination reduced the total weed dry biomass by 93% compared to chemical fallow (Table 2). The CC terminated with GLY only and with GLY plus ACR/ATZ reduced the total weed dry biomass by 50 to 65% and 42 to 68% at 60 to 120 DAT, respectively compared to chemical fallow. It is interesting to note that no differences were observed in total weed density between chemical fallow and CC terminated with GLY only at 120 DAT; however, the same CC treatment significantly reduced the total weed dry biomass by 42% compared to chemical fallow, indicating the suppressive effect of CC on weed growth that would ultimately reduce the weed seed production (Baraibar et al. 2018).

**2022 growing season.** Mean relative abundance was 46 to 56% for *B. scoparia*, 0 to 21% for *A. palmeri*, and 29 to 49% for *H. trionum* before CC termination (Table 3). Similar to 2021, the CC at termination reduced the total weed density by 90 to 93% compared to weedy fallow. At 30 DAT, the CC terminated with GLY plus ACR/ATZ reduced the total weed density by 92% compared to chemical fallow and 96% compared to weedy fallow. At 60 DAT, the mean relative abundance was 41 to 75% for *B. scoparia*, 11 to 53% for *A. palmeri*, and 12 to 31% for *H. trionum* among all treatments. The CC termination with GLY only treatment did not produce enough CC biomass to suppress emerging weed seedlings and resulted in 21 more weed seedlings m<sup>-2</sup> than the CC terminated with GLY plus ACR/ATZ at 60 DAT (Table 3). These results indicate the importance of residual herbicide with CC to achieve effective weed suppression. Our findings are consistent with Wiggins et al. (2016), who also concluded that CC alone was not enough for season-long control of GR *A. palmeri* and suggested integrating residual herbicides to complement the suppressive effect of CC. In the current study, total weed density did not differ between chemical fallow and CC terminated with GLY plus ACR/ATZ treatments at 120 DAT; however, weed density was nearly 95% lower in CC terminated with GLY plus ACR/ATZ compared to weedy fallow and CC terminated with GLY only (Table 3). These results are consistent with Obour et al. (2022a), who previously reported 82% reduction in total weed density with spring-planted CC mixture (oat/triticale/pea) compared to weedy fallow.

Petrosino et al. (2015) also reported a 78 to 94% reduction in *B. scoparia* density with fall-planted CC (triticale/triticale-hairy vetch mixture) compared to chemical fallow in winter wheat-fallow rotation. Similar to weed density reduction, CC at the time of termination provided >95% total weed dry biomass reduction compared to weedy fallow (Table 2). The presence of CC reduces the sunlight penetration to the ground for weed seed germination and also reduces their competitive ability for other resources, thereby resulting in lower weed biomass (Silva and Bagavathiannan 2023; Webster et al. 2016). The CC terminated with GLY only reduced total weed dry biomass by 94% at 30 DAT and 63% at 120 DAT compared to weedy fallow. In contrast, the CC terminated with GLY plus ACR/ATZ reduced total weed dry biomass by >95% as compared to weedy fallow throughout the sorghum growing season (Table 2). These results indicate the need of residual herbicide in combination with CC for a season-long weed control in grain sorghum. These results are consistent with Whalen et al. (2020), who previously reported that fall-planted CC, including Italian ryegrass [*Lolium perenne* L. ssp. *multiflorum* (Lam.) Husnot], oat (*Avena sativa* L.), and winter wheat provided 38 to 48% weed control without a residual herbicide; however, the control ranged from 72 to 85% under CC with a residual herbicide (sulfentrazone plus chlorimuron) application.

**2023 growing season.** No weed emergence was observed under both CC treatments at the time of termination (Table 4). In addition, no weed emergence was observed in the CC terminated with GLY plus ACR/ATZ treatment at 30 DAT. In contrast, a greater weed density of 91 plants m<sup>-2</sup> with a relative abundance of 70% for *A. palmeri* and 30% for *B. scoparia* was recorded in the CC terminated with GLY only treatment at 30 DAT (Table 4). This increase in *A. palmeri* and *B. scoparia* densities under CC terminated with GLY only was probably due to more precipitation within 30 DAT (78 mm, 33% of total precipitation received during the entire sorghum growing season), and lack of any residual herbicide applied at CC termination. Treatments, including the CC terminated with GLY plus ACR/ATZ and chemical fallow reduced the total weed density by 70% compared to weedy fallow at 90 DAT. Chemical fallow and CC terminated with GLY plus ACR/ATZ had much fewer weeds (7 to 8 plants m<sup>-2</sup>) compared to weedy fallow (54 plants m<sup>-2</sup> with 73% relative abundance of *A. palmeri*) at 120 DAT (Table 4). Similar to weed density reduction, CC terminated with GLY plus ACR/ATZ resulted in 98%, 46%, and 57% total weed dry biomass reduction compared to weedy fallow, chemical fallow, and CC terminated with

GLY only, respectively, at 60 DAT (Table 2). At 120 DAT, the CC terminated with GLY plus ACR/ATZ and chemical fallow reduced the total weed dry biomass by 83% and 65% compared to weedy fallow. Our results are consistent with Petrosino et al. (2015), who also reported that fall-planted triticale and a triticale-hairy vetch mixture reduced *B. scoparia* density by 78% and 94%, respectively, and biomass up to 98% compared to chemical fallow. Compared to chemical fallow, the CC terminated with GLY plus ACR/ATZ provided 5 to 18% more reductions in total weed dry biomass during the entire grain sorghum growing season (Table 2). Wiggins et al. (2016) reported that CC including cereal rye (*Secale cereale* L.), winter vetch, crimson clover (*Trifolium incarnatum* L.), or winter wheat had less than 65% control of *A. palmeri*, whereas the same CC in combination with PRE-applied acetochlor or fluometuron resulted in >87% control of *A. palmeri*.

### **Volumetric water content and grain sorghum yield**

The soil water content at sorghum planting is directly related to grain yields in the dryland region (Holman et al. 2023; Obour et al. 2022b). In 2021, fall-planted CC did not affect volumetric water content at grain sorghum planting compared to chemical fallow across all soil depths (Figure 2). This is likely due to greater precipitation at pre- and post-CC termination time (Figure 1). The precipitation received closer to termination or post-termination of CC likely recharged the soil profile thereby diminishing the effects of the growing CC on water availability. Furthermore, the CC residue likely decreased soil water evaporation, increasing moisture storage (Holman et al. 2020, 2021). In contrast, fall-planted CC did reduce the volumetric water content at grain sorghum planting compared to chemical fallow at 0-30 cm and 30-60 cm depths in the 2023 growing season (Figure 2). No differences in the water content were observed between CC and chemical fallow treatments at deeper soil depths. Similar to CC, the weedy fallow treatment also resulted in relatively low volumetric water content ( $0.16$  to  $0.20$   $\text{cm}^3 \text{cm}^{-3}$ ) up to 60 cm soil depth compared to chemical fallow ( $0.20$  to  $0.22$   $\text{cm}^3 \text{cm}^{-3}$ ) indicating soil water depletion by weeds. Holman et al. (2021) reported no difference in the available soil moisture at time of wheat planting between fallow and spring-planted CC left standing during the fallow phase of W-S-F rotation.

No differences in grain sorghum yields were observed between chemical fallow ( $1876$   $\text{kg ha}^{-1}$ ) and CC terminated with GLY plus ACR/ATZ ( $2072$   $\text{kg ha}^{-1}$ ) in 2021 and the least grain yield

(1456 kg ha<sup>-1</sup>) was recorded under CC terminated with GLY only (Figure 3). During the 2022 growing season, the overall grain sorghum yield was low due to lower season precipitation (171 mm) compared to 2021 (256 mm). Effective weed suppression (both density and total weed dry biomass) achieved with the CC terminated with GLY plus ACR/ATZ resulting in a greater grain sorghum yield (1319 kg ha<sup>-1</sup>) compared to chemical fallow (912 kg ha<sup>-1</sup>). The CC termination with GLY only suppressed weeds up to 30 DAT (Table 3) and *A. palmeri* emerging after 30 DAT reduced the grain sorghum yield (472 kg ha<sup>-1</sup>) (Figure 3). The total precipitation during the 2023 sorghum growing season was 237 mm, but the majority of this precipitation occurred in May and June. There was moisture stress at the boot stage of grain sorghum in September (only 17 mm of rainfall) that resulted in reduced grain yield (432 to 1323 kg ha<sup>-1</sup>) (Figures 1 and 3). No difference in grain sorghum yield was observed between chemical fallow and CC terminated with GLY plus ACR/ATZ. The precipitation events in May and June resulted in the emergence of several cohorts of *A. palmeri* following the CC terminated with GLY only that resulted in competition with grain sorghum and reduced yield (432 kg ha<sup>-1</sup>) compared to chemical fallow and CC terminated with GLY plus ACR/ATZ (Figure 3). Based on the precipitation amount and frequency, fall-planted CC had a variable impact on grain sorghum yield. Nielsen et al. (2016) reported a 10% reduction in winter wheat yield following a CC compared to fallow in the W-S-F rotation under the semi-arid CGP. Holman et al. (2022) reported that spring-planted CC (oat grain) after sorghum harvesting in the fallow phase of W-S-F rotation resulted in 29% less available soil moisture at sorghum planting and did not affect the wheat and sorghum yield compared to fallow. In south-central Kansas, Janke et al. (2002) reported no differences in grain sorghum yield following CC (hairy vetch/winter pea) in winter wheat-grain sorghum rotation compared to no CC in years with good rainfall; however, during years with low rainfall, CC establishment was poor, and grain sorghum yield was reduced because of water use by the CC compared to no CC treatment. Eash et al. (2021) also reported that CC had no impact on subsequent crop yields in a very low-yielding environment in Colorado, mainly due to low CC biomass. In this study, the chemical fallow treatment provided more available soil water in one of two years compared to CC treatments prior to sorghum planting. However, this did not translate into a higher grain yield as long as the residual herbicide ACR/ATZ was applied with GLY at CC termination.

## Economics analyses

During 2021, the CC terminated with GLY plus ACR/ATZ resulted in greater gross returns of \$497 ha<sup>-1</sup> compared to \$450 ha<sup>-1</sup> following chemical fallow (Table 5). This was because of better weed control and greater sorghum yield under CC terminated with GLY plus ACR/ATZ compared to chemical fallow. However, CC seed and planting costs decreased net returns. Chemical fallow had the highest net returns by \$55 ha<sup>-1</sup> compared to CC terminated with GLY plus ACR/ATZ. Net returns were negative under CC terminated with GLY only because of lower grain sorghum yield than chemical fallow and CC terminated with GLY plus ACR/ATZ.

Similar to 2021, CC terminated with GLY plus ACR/ATZ had greater gross returns (\$317 ha<sup>-1</sup>) followed by chemical fallow (\$219 ha<sup>-1</sup>) and CC terminated with GLY only (\$113 ha<sup>-1</sup>) in 2022 (Table 6). Weedy fallow had the lowest gross returns of \$24 ha<sup>-1</sup>. The net returns for 2022 were negative for all treatments because of low grain sorghum yield which suggests that gross return from grain sorghum was not enough to cover the variable input costs. However, net returns for CC terminated with GLY plus ACR/ATZ and chemical fallow were not significantly different (Table 6).

In 2023, the lower grain sorghum yield in CC terminated with GLY plus ACR/ATZ resulted in a lower gross return of \$181 ha<sup>-1</sup> compared to \$265 ha<sup>-1</sup> in chemical fallow (Table 7). Similar to 2022, the net returns in 2023 were also negative for all treatments. Chemical fallow had less negative net returns (-\$65 ha<sup>-1</sup>) compared to other treatments (-\$138 to -\$275 ha<sup>-1</sup>). The cost of CC seed and planting increased the variable cost for both CC treatments and thus resulted in greater negative net returns. Janke et al. (2002) also reported lower net returns with fall-planted CC (hairy vetch/winter pea) before grain sorghum because of lower grain sorghum yield in the years with low rainfall. Obour et al. (2022a) also reported that integrating CC in the dryland cropping system resulted in negative net returns compared to fallow. These results indicate that growing CC only for weed suppression was not profitable compared to chemical fallow.

Based on the sensitivity analysis, a minimum grain sorghum yield of 2000 kg ha<sup>-1</sup> was needed to obtain a positive net return (\$8 ha<sup>-1</sup>) following CC terminated with GLY plus ACR/ATZ at average grain sorghum price (\$0.22 kg<sup>-1</sup>) during the 3-year study period (Table 8). In the present study, the average grain sorghum yield was low (786 to 1432 kg ha<sup>-1</sup>). The estimated average grain sorghum yield in western Kansas ranged from 3800 to 5000 kg ha<sup>-1</sup>

(Ciampitti and Carcedo 2022) therefore, based on this yield scenario, the expected net returns would be \$404 to \$668 ha<sup>-1</sup> following CC terminated with GLY plus ACR/ATZ and \$506 to \$770 ha<sup>-1</sup> following chemical fallow at \$0.22 kg<sup>-1</sup> sorghum price (Table 8). At the lowest grain sorghum price (\$0.09 ha<sup>-1</sup>), a grain yield of 5000 kg ha<sup>-1</sup> was expected to cover the cost of CC seed and planting and obtain positive net returns (\$18 ha<sup>-1</sup>) under CC terminated with GLY plus ACR/ATZ, however, following chemical fallow, a lower yield of 3800 kg ha<sup>-1</sup> was sufficient to obtain positive net returns (\$12 ha<sup>-1</sup>) (Table 8). It is important to note that in the present study, greater weed control was observed following CC terminated with GLY plus ACR/ATZ over the grain sorghum growing season compared to chemical fallow. The POST herbicides were not applied in chemical fallow treatment but farmers generally apply POST herbicides to control later emerged weeds and this application would increase the cost of production and decrease the net returns following chemical fallow as compared to CC treatment. At the highest sorghum price (\$0.24 ha<sup>-1</sup>), CC terminated with GLY plus ACR/ATZ would result in \$48 ha<sup>-1</sup> net returns at only 2000 kg ha<sup>-1</sup> sorghum yield (Table 8). These results indicate the cost of integrating fall-planted CC in the W-S-F exceeded the benefits of improved weed control.

Results from this 3-year study indicate that integrating a fall-planted CC mixture after winter wheat harvest and terminating it with GLY in combination with residual herbicide before grain sorghum planting in the W-S-F rotation can provide an effective weed suppression in grain sorghum. Results showed that fall-planted CC terminated with GLY plus ACR/ATZ reduced total weed density by 34 to 81% and total weed biomass by 45 to 73% compared to chemical fallow over the grain sorghum growing season. No sorghum yield penalty was observed after CC terminated with GLY plus ACR/ATZ. However, due to the CC seed and planting costs, lower net returns were recorded in all three years compared to chemical fallow. These results suggest that growing CC for only weed suppression in the semi-arid CGP would not be profitable at current commodity prices. If the CC were used for hay or forage then net returns would be increased due to alternative use income (Holman et al. 2018, 2021, 2022, 2023; Obour et al. 2022a). However, weed control during the sorghum growing season would likely be affected after the CC residue removal from the field. Therefore, future studies should focus on understanding the timing for the removal of CC residue from the field and its interaction with weed control during the grain sorghum growing period.

**Acknowledgments.** We thank Mr. Taylor Lambert and Mr. Matthew Vredenburg for their assistance in conducting the field study.

**Funding.** The funding from the NC SARE Graduate Student Grant (GNC22-346) supported this work.

**Competing Interests.** The authors declare no competing interests.

## References

- AgManager (2022) Kansas Custom Rates 2022. Kansas Department of Agriculture and Kansas State University Land Use Survey Program. <https://www.agmanager.info/machinery/papers/custom-rates-survey>. Accessed: September 10, 2024
- Anonymous (2024) USDA Natural Resources Conservation Service. Environmental Quality Incentives Program. <https://www.nrcs.usda.gov/programs-initiatives/eqip-environmental-quality-incentives>. Accessed: September 10, 2024
- Baraibar B, Hunter MC, Schipanski ME, Hamilton A, Mortensen DA (2018) Weed suppression in cover crop monocultures and mixtures. *Weed Sci* 66:121-133
- Blanco-Canqui H, Holman JD, Schlegel AJ, Tatarko J, Shaver TM (2013) Replacing fallow with cover crops in a semiarid soil: Effects on soil properties. *Soil Sci Soc Am J* 77:1026-1034
- Blanco-Canqui H, Mikha MM, Presley DR, Claassen MM (2011) Addition of cover crops enhances no-till potential for improving soil physical properties. *Soil Sci Soc Am J* 75:1471-1482
- Ciampitti IA, Diaz DR, Onofre R, Lancaster S, Whitworth RJ, Aguilar J (2022a) Kansas sorghum management 2022 (Extension Bulletin MF3208). Kansas State University Agricultural Experiment Station and Cooperative Extension Service, 8 p
- Ciampitti IA, Carcedo A (2022b) Agronomy eUpdates. Sorghum management considerations: Planting practices, 908. [https://eupdate.agronomy.ksu.edu/article\\_new/sorghum-management-considerations-planting-practices-496](https://eupdate.agronomy.ksu.edu/article_new/sorghum-management-considerations-planting-practices-496). Accessed: September 10, 2024



- Dille JA, Stahlman PW, Thompson CR, Bean BW, Soltani N, Sikkema PH (2020) Potential yield loss in grain sorghum (*Sorghum bicolor*) with weed interference in the United States. *Weed Technol* 34:624-629
- Eash L, Berrada AF, Russell K, Fonte SJ (2021) Cover crop impacts on water dynamics and yields in dryland wheat systems on the Colorado Plateau. *Agron* 11:1102
- Hansen NC, Allen BL, Baumhardt RL, Lyon DJ (2012) Research achievements and adoption of no-till, dryland cropping in the semi-arid US Great Plains. *Field Crops Res* 132:196-203
- Heap I (2024) The International Herbicide-resistant weed database. <http://www.weedscience.org>. Accessed: September 10, 2024
- Holman JD, Arnet K, Dille J, Maxwell S, Obour A, Roberts T, Roozeboom K, Schlegel A (2018) Can cover or forage crops replace fallow in the semiarid central Great Plains? *Crop Sci* 58:932-944
- Holman JD, Assefa Y, Obour AK (2020) Cover-crop water use and productivity in the high plains wheat-fallow crop rotation. *Crop Sci* 61:1374-1385
- Holman JD, Obour AK, Assefa Y (2021) Fallow replacement cover crops in a semi-arid High Plains cropping system. *Crop Sci* 61:3799-3814
- Holman JD, Obour AK, Assefa Y (2022) Productivity and profitability with fallow replacement forage, grain, and cover crops in W-S-F rotation. *Crop Sci* 62:913-927
- Holman JD, Obour AK, Assefa Y (2023) Forage sorghum grown in a conventional wheat-grain sorghum-fallow rotation increased cropping system productivity and profitability. *Can J Plant Sci* 103:61-72
- Janke RR, Claassen MM, Heer WF, Jost J, Freyenberger S, Norman D (2002) The use of winter annual legume cover crops in a wheat-grain sorghum rotation in south central Kansas. *J Sus Agric* 20:69-88
- Jha P, Kumar V, Lim CA (2016) Herbicide resistance in cereal production systems of the US Great Plains: A review. *Field Crops Res* 183:56-68
- Kumar V, Liu R, Boyer G, Stahlman PW (2019a) Confirmation of 2, 4-D resistance and identification of multiple resistance in a Kansas Palmer amaranth (*Amaranthus palmeri*) population. *Pest Manag Sci* 75:2925-2933
- Kumar V, Jha P, Jugulam M, Yadav R, Stahlman PW (2019b) Herbicide-resistant kochia (*Bassia scoparia*) in North America: a review. *Weed Sci* 67:4-15

- Kumar V, Obour A, Jha P, Liu R, Manuchehri MR, Dille JA, Holman J, Stahlman PW (2020) Integrating cover crops for weed management in the semiarid US Great Plains: opportunities and challenges. *Weed Sci* 68:311-323
- Lenssen AW, Johnson GD, Carlson GR (2007) Cropping sequence and tillage system influences annual crop production and water use in semiarid Montana, USA. *Field Crops Res* 100:32-43
- Mesbah A, Nilahyane A, Ghimire B, Beck L, Ghimire R (2019) Efficacy of cover crops on weed suppression, wheat yield, and water conservation in winter wheat–sorghum–fallow. *Crop Sci* 59:1745-1752
- Nichols V, Verhulst N, Cox R, Govaerts B (2015) Weed dynamics and conservation agriculture principles: A review. *Field Crops Res* 183:56-68
- Nielsen DC, Lyon DJ, Higgins RK, Hergert GW, Holman JD, Vigil MF (2016) Cover crop effect on subsequent wheat yield in the central Great Plains. *Agron J* 108:243-256
- NOAA (2024). National Oceanic and Atmospheric Administration. <https://www.noaa.gov>. Accessed: March 10, 2024
- Obour AK, Dille J, Holman J, Simon LM, Sancewich B, Kumar V (2022a) Spring-planted cover crop effects on weed suppression, crop yield, and net returns in no-tillage dryland crop production. *Crop Sci* 62:1981-1996
- Obour AK, Holman JD, Assefa Y (2022b) Grain sorghum productivity as affected by nitrogen rates and available soil water. *Crop Sci* 62:360-1372
- Perkins CM, Gage KL, Norsworthy JK, Young BG, Bradley KW, Bish MD, Hager A, Steckel LE (2021) Efficacy of residual herbicides influenced by cover-crop residue for control of *Amaranthus palmeri* and *A. tuberculatus* in soybean. *Weed Technol* 35:77-81
- Petrosino JS, Dille JA, Holman JD, Roozeboom KL (2015) Kochia suppression with cover crops in southwestern Kansas. *Crop Forage Turfgrass Manag* 1:1-8
- Silva GC, Bagavathiannan M (2023) Mechanisms of weed suppression by cereal rye cover crop: A review. *Agron J* 115:1571-1585
- Simon LM, Obour AK, Holman JD, Roozeboom KL (2022) Long-term cover crop management effects on soil properties in dryland cropping systems. *Agric Ecosyst Environ* 328:107852

- Teasdale JR, Pillai P, Collins RT (2005) Synergism between cover crop residue and herbicide activity on emergence and early growth of weeds. *Weed Sci* 53:521-527
- Thomas AG (1985) Weed survey system used in Saskatchewan for cereal and oilseed crops. *Weed Sci* 33:34-43
- USDA ERS (2023) Commodity cost and returns. Economic Research Service. <https://www.ers.usda.gov/data-products/>. Accessed: September 11, 2024
- Webster TM, Simmons DB, Culpepper AS, Grey TL, Bridges DC, Scully BT (2016) Factors affecting potential for Palmer amaranth (*Amaranthus palmeri*) suppression by winter rye in Georgia, USA. *Field Crops Res* 192:103–109
- Weisberger DA, Bastos LM, Sykes VR, Basinger NT (2023) Do cover crops suppress weeds in the US Southeast? A meta-analysis. *Weed Sci* 71:244-254
- Westra EP, Nissen SJ, Getts TJ, Westra P, Gaines TA (2019) Survey reveals frequency of multiple resistance to glyphosate and dicamba in kochia (*Bassia scoparia*). *Weed Technol* 33:664-672
- Whalen DM, Shergill LS, Kinne LP, Bish MD, Bradley KW (2020) Integration of residual herbicides with cover crop termination in soybean. *Weed Technol* 34:11-18
- Wiggins MS, Hayes RM, Steckel LE (2016) Evaluating cover crops and herbicides for glyphosate-resistant Palmer amaranth (*Amaranthus palmeri*) control in cotton. *Weed Technol* 30:415-422

**Table 1.** Total weed density and mean relative abundance of weed species observed in the cover crop (CC) treatments at 0 to 120 days after CC termination (DAT) in 2021<sup>a, b</sup>

Treatments	Total weed density		Mean relative abundance			
			<i>Bassia scoparia</i>	<i>Amaranthus palmeri</i>	<i>Hibiscus trionum</i>	<i>Tribulus terrestris</i>
	plants m <sup>-2</sup>		-----%			
<b>At 0 DAT</b>						
Chemical fallow	43	a	43	41	16	0
CC + GLY	6	b	0	100	0	0
CC + GLY + ACR/ATZ	4	b	0	100	0	0
<b>At 30 DAT</b>						
Chemical fallow	10	a	38	50	4	8
CC + GLY	10	a	12	69	0	19
CC + GLY + ACR/ATZ	6	a	11	53	8	28
<b>At 60 DAT</b>						
Chemical fallow	9	a	24	70	0	6
CC + GLY	12	a	10	90	0	0
CC + GLY + ACR/ATZ	3	b	10	90	0	0
<b>At 120 DAT</b>						
Chemical fallow	4	a	50	50	0	0
CC + GLY	5	a	29	71	0	0
CC + GLY + ACR/ATZ	2	b	39	61	0	0

<sup>a</sup>CC + GLY indicates cover crop terminated with glyphosate only and CC + GLY + ACR/ATZ indicates cover crop terminated with glyphosate plus a premix of acetochlor/atrazine. Means followed by the same letter within a column at each timing are not different according to Fisher's protected LSD at P < 0.05

<sup>b</sup>Cover crop was terminated on May 13, 2021

**Table 2.** Total weed dry biomass in the cover crop (CC) treatments at 0, 30, 60, 90, and 120 days after CC termination during 2021 to 2023 growing seasons<sup>a</sup>

Treatments	Total weed dry biomass																											
	2021				2022					2023																		
	0	30	60	120	0	30	60	90	120	0	30	60	90	120														
	-----g m <sup>-2</sup> -----																											
Weedy fallow	-	-	-	-	34	a	10	a	11	a	193	a	26	a	12	a	11	a	38	a	143	a	162	a				
Chemical fallow	15	a	72	a	13	a	14	a	30	a	2	c	3	bc	6	c	12	c	9	a	5	c	13	c	345	b	560	b
CC + GLY	1	b	41	a	69	b	85	b	1	b	6	b	15	b	41	b	97	b	0	b	44	b	16	b	513	b	125	a
CC + GLY + ACR/ATZ	1	b	24	a	48	b	47	b	1	b	1	c	1	c	3	c	10	c	0	b	0	d	7	d	236	c	278	c

<sup>a</sup>CC + GLY indicates cover crop terminated with glyphosate only and CC + GLY + ACR/ATZ indicates cover crop terminated with glyphosate plus a premix of acetochlor/atrazine. Means followed by the same letter within a column are not different according to Fisher's protected LSD at P < 0.05.

**Table 3.** Total weed density and mean relative abundance of weed species observed in the cover crop (CC) treatments at 0 to 120 days after CC termination (DAT) in 2022<sup>a, b</sup>

Treatments	Total weed density plants m <sup>-2</sup>		Mean relative abundance			
			<i>Bassia scoparia</i>	<i>Amaranthus palmeri</i>	<i>Hibiscus trionum</i>	<i>Tribulus terrestris</i>
<b>At 0 DAT</b>						
Weedy fallow	58	a	46	21	33	0
Chemical fallow	47	a	53	18	29	0
CC + GLY	4	b	51	0	49	0
CC + GLY + ACR/ATZ	6	b	56	6	38	0
<b>At 30 DAT</b>						
Weedy fallow	27	ab	56	36	6	2
Chemical fallow	13	b	62	32	6	0
CC + GLY	37	a	41	54	5	0
CC + GLY + ACR/ATZ	1	c	62	38	0	0
<b>At 60 DAT</b>						
Weedy fallow	28	a	58	11	31	0
Chemical fallow	6	bc	47	53	0	0
CC + GLY	22	ab	41	47	12	0
CC + GLY + ACR/ATZ	1	c	75	25	0	0
<b>At 90 DAT</b>						
Weedy fallow	27	a	80	20	0	0
Chemical fallow	1	b	100	0	0	0
CC + GLY	13	a	43	57	0	0
CC + GLY + ACR/ATZ	1	b	87	13	0	0
<b>At 120 DAT</b>						
Weedy fallow	19	a	93	7	0	0
Chemical fallow	2	b	91	9	0	0
CC + GLY	21	a	46	54	0	0
CC + GLY + ACR/ATZ	1	b	89	11	0	0

<sup>a</sup>CC + GLY indicates cover crop terminated with glyphosate only and CC + GLY + ACR/ATZ indicates cover crop terminated with glyphosate plus a premix of acetochlor/atrazine. Means followed by the same letter within a column at each timing are not different according to Fisher's protected LSD at P < 0.05.

<sup>b</sup>Cover crop was terminated on May 11, 2022

**Table 4.** Total weed density and mean relative abundance of weed species observed in the cover crop (CC) treatments at 0 to 120 days after CC termination (DAT) in 2023<sup>a, b</sup>

Treatments	Total weed density plants m <sup>-2</sup>		Mean relative abundance			
			<i>Bassia scoparia</i>	<i>Amaranthus palmeri</i>	<i>Hibiscus trionum</i>	<i>Tribulus terrestris</i>
<b>At 0 DAT</b>						
Weedy fallow	32	a	100	0	0	0
Chemical fallow	28	a	100	0	0	0
CC + GLY	0	b	0	0	0	0
CC + GLY + ACR/ATZ	0	b	0	0	0	0
<b>At 30 DAT</b>						
Weedy fallow	142	a	37	61	2	1
Chemical fallow	12	b	0	100	0	0
CC + GLY	91	a	30	70	0	0
CC + GLY + ACR/ATZ	0	b	0	0	0	0
<b>At 60 DAT</b>						
Weedy fallow	22	a	13	87	0	0
Chemical fallow	7	b	0	100	0	0
CC + GLY	15	ab	25	75	0	0
CC + GLY + ACR/ATZ	1	b	0	100	0	0
<b>At 90 DAT</b>						
Weedy fallow	30	a	16	84	0	0
Chemical fallow	9	b	10	90	0	0
CC + GLY	20	ab	15	85	0	0
CC + GLY + ACR/ATZ	9	b	0	100	0	0
<b>At 120 DAT</b>						
Weedy fallow	54	a	27	73	0	0
Chemical fallow	8	b	19	81	0	0
CC + GLY	22	ab	27	73	0	0
CC + GLY + ACR/ATZ	7	b	0	100	0	0

<sup>a</sup>CC + GLY indicates cover crop terminated with glyphosate only and CC + GLY + ACR/ATZ indicates cover crop terminated with glyphosate plus a premix of acetochlor/atrazine. Means followed by the same letter within a column at each timing are not different according to Fisher's protected LSD at P < 0.05.

<sup>b</sup> Cover crop was terminated on May 22, 2023

**Table 5.** Economic analyses of grain sorghum after fall-planted cover crop in 2021 growing season<sup>a</sup>

Variables	Treatments		
	Chemical fallow	CC + GLY	CC + GLY + ACR/ATZ
	-----\$ ha <sup>-1</sup> -----		
Grain sorghum yield <sup>b</sup>	1877.0	1456.0	2072.0
Grain sorghum price <sup>c</sup>	0.24	0.24	0.24
Revenue from grain sorghum	450.5	349.4	497.3
Gross returns	450.5	349.4	497.3
Variable input costs			
Cover crop seed	0	95.7	95.7
Cover crop planting	0	37.1	37.1
Grain sorghum seed	30.4	30.4	30.4
Grain sorghum planting	32.1	32.1	32.1
Fertilizer with application cost	66.7	66.7	66.7
Herbicide	126.9	25.6	95.7
Herbicide application cost	13.6	13.6	13.6
Grain sorghum harvesting	60.3	60.3	60.3
Total variable cost	330.0	361.5	431.6
Net returns	120.5 a	-12.1 b	65.7 b

<sup>a</sup>CC + GLY indicates cover crop terminated with glyphosate only and CC + GLY + ACR/ATZ indicates cover crop terminated with glyphosate plus a premix of acetochlor/atrazine. Means followed by the same letter among treatments are not different according to Fisher's protected LSD at  $P < 0.05$

<sup>b</sup>Grain sorghum yield is in  $\text{kg ha}^{-1}$

<sup>c</sup>Grain sorghum price is in  $\text{\$ kg}^{-1}$



**Table 6.** Economic analyses of grain sorghum after fall-planted cover crop in 2022 growing season<sup>a</sup>

Variables	Treatments			
	Weedy fallow	Chemical fallow	CC + GLY	CC + GLY + ACR/ATZ
	-----\$ ha <sup>-1</sup> -----			
Grain sorghum yield <sup>b</sup>	101.0	912.0	472.0	1319.0
Grain sorghum price <sup>c</sup>	0.24	0.24	0.24	0.24
Revenue from grain sorghum	24.2	218.9	113.3	316.6
Gross returns	24.2	218.9	113.3	316.6
Variable input costs				
Cover crop seed	0	0	95.7	95.7
Cover crop planting	0	0	37.1	37.1
Grain sorghum seed	30.4	30.4	30.4	30.4
Grain sorghum planting	32.1	32.1	32.1	32.1
Fertilizer with application cost	66.7	66.7	66.7	66.7
Herbicide	0	126.9	25.6	95.7
Herbicide application cost	0	13.6	13.6	13.6
Grain sorghum harvesting	60.3	60.3	60.3	60.3
Total variable cost	189.5	330.0	361.5	431.6
Net returns	-165.3 b	-111.1 a	-248.2 b	-115.0 a

<sup>a</sup>CC + GLY indicates cover crop terminated with glyphosate only and CC + GLY + ACR/ATZ indicates cover crop terminated with glyphosate plus a premix of acetochlor/atrazine. Means followed by the same letter among treatments are not different according to Fisher's protected LSD at  $P < 0.05$ .

<sup>b</sup>Grain sorghum yield is in  $\text{kg ha}^{-1}$

<sup>c</sup>Grain sorghum price is in  $\text{\$ kg}^{-1}$

**Table 7.** Economic analyses of grain sorghum after fall-planted cover crop in 2023 growing season<sup>a</sup>

Variables	Treatments			
	Weedy fallow	Chemical fallow	CC + GLY	+ CC + GLY + ACR/ATZ
	-----\$ ha <sup>-1</sup> -----			
Grain sorghum yield <sup>b</sup>	257.0	1323.0	432.0	905.0
Grain sorghum price <sup>c</sup>	0.20	0.20	0.20	0.20
Revenue from grain sorghum	51.4	264.6	86.4	181.0
Gross returns	51.4	264.6	86.4	181.0
Variable input costs				
Cover crop seed	0	0	95.7	95.7
Cover crop planting	0	0	37.1	37.1
Grain sorghum seed	30.4	30.4	30.4	30.4
Grain sorghum planting	32.1	32.1	32.1	32.1
Fertilizer with application cost	66.7	66.7	66.7	66.7
Herbicide	0	126.9	25.6	95.7
Herbicide application cost	0	13.6	13.6	13.6
Grain sorghum harvesting	60.3	60.3	60.3	60.3
Total variable cost	189.5	330.0	361.5	431.6
Net returns	-138.1 b	-65.4 a	-275.1 c	-250.6 c

<sup>a</sup>CC + GLY indicates cover crop terminated with glyphosate only and CC + GLY + ACR/ATZ indicates cover crop terminated with glyphosate plus a premix of acetochlor/atrazine. Means followed by the same letter among treatments are not different according to Fisher's protected LSD at P < 0.05


<sup>b</sup>Grain sorghum yield is in kg ha<sup>-1</sup>

<sup>c</sup>Grain sorghum price is in \$ kg<sup>-1</sup>


**Table 8.** Net returns to possible grain sorghum yield (700 to 7400 kg ha<sup>-1</sup>) and prices (\$0.09 to \$0.24 kg<sup>-1</sup>) in the region<sup>a</sup>

Grain sorghum yield kg ha <sup>-1</sup>	Net returns													
	Chemical fallow							CC + GLY + ACR/ATZ						
	0.09	0.12	0.16	0.18	0.20	0.22	0.24	0.09	0.12	0.16	0.18	0.20	0.22	0.24
	-----\$ ha <sup>-1</sup> -----													
700	-267	-246	-218	-204	-190	-176	-162	-369	-348	-320	-306	-292	-278	-264
1400	-204	-162	-106	-78	-50	-22	6	-306	-264	-208	-180	-152	-124	-96
2000	-150	-90	-10	30	70	110	150	-252	-192	-112	-72	-32	8	48
2600	-96	-18	86	138	190	242	294	-198	-120	-16	36	88	140	192
3200	-42	54	182	246	310	374	438	-144	-48	80	144	208	272	336
3800	12	126	278	354	430	506	582	-90	24	176	252	328	404	480
4400	66	198	374	462	550	638	726	-36	96	272	360	448	536	624
5000	120	270	470	570	670	770	870	18	168	368	468	568	668	768
5600	174	342	566	678	790	902	1014	72	240	464	576	688	800	912
6200	228	414	662	786	910	1034	1158	126	312	560	684	808	932	1056
6800	282	486	758	894	1030	1166	1302	180	384	656	792	928	1064	1200
7400	336	558	854	1002	1150	1298	1446	234	456	752	900	1048	1196	1344


<sup>a</sup>CC + GLY + ACR/ATZ indicates cover crop terminated with glyphosate plus a premix of acetochlor/atrazine



**Figure 1.** Total monthly precipitation (mm) and average monthly air temperature (C) from 2020 to 2023 growing seasons.



**Figure 2.** Fall-planted cover crop effect on volumetric water content at grain sorghum planting in 2021 (A) and 2023 (B) growing seasons. Error bars represent standard error of the mean.



**Figure 3.** Fall-planted cover crop effect on grain sorghum yield over three growing seasons. CC + GLY indicates cover crop terminated with glyphosate only and CC + GLY + ACR/ATZ indicates cover crop terminated with glyphosate plus a premix of acetochlor/atrazine. Means followed by the same letter within the year are not significantly different according to Fisher's protected LSD test at  $P < 0.05$ . Error bars represent standard error of the mean.