

Research Paper

*Joint corresponding authors.

Cite this article: Hu L *et al.* (2024). Alteration in microbes changed the contents of oviposition-deterrent pheromones on the *Spodoptera litura* egg surface. *Bulletin of Entomological Research* 1–8. <https://doi.org/10.1017/S000748532400066X>

Received: 26 June 2024

Revised: 18 August 2024

Accepted: 20 October 2024

Keywords:


16S rDNA sequencing; egg surface; microorganisms symbiotic; oviposition-deterrent pheromones; *Spodoptera litura*

Corresponding author:

Jianmei Shen;

Email: shenjianmei@zhku.edu.cn

Alteration in microbes changed the contents of oviposition-deterrent pheromones on the *Spodoptera litura* egg surface

Liming Hu¹, Yirui Chen¹, Qingjun Wu², Qiumei Zeng¹, Taoli Zhang¹, Guohui Yu¹, Muyang He¹, Dasong Chen³, Xiangning Su⁴, Yuping Zhang⁴, Zhenfei Zhang^{4,*} and Jianmei Shen^{1,*} 

¹Key Laboratory of Green Prevention and Control on Fruits and Vegetables in South China, Ministry of Agriculture and Rural Affairs, Zhongkai University of Agriculture and Engineering, Guangzhou, China; ²State Key Laboratory of Vegetable Biobreeding, Department of Plant Protection, Institute of Vegetables and Flowers, Chinese Academy of Agricultural Sciences, Beijing, China; ³Guangdong Key Laboratory of Animal Conservation and Resource Utilization, Guangdong Public Laboratory of Wild Animal Conservation and Utilization, Institute of Zoology, Guangdong Academy of Sciences, Guangzhou, China and ⁴Plant Protection Research Institute, Guangdong Academy of Agricultural Sciences & Key Laboratory of Green Prevention and Control of Fruits and Vegetables in South China, Ministry of Agriculture and Rural Affairs & Key Laboratory of High Technology for Plant Protection of Guangdong Province, Guangzhou, China

Abstract

Microorganisms symbiotic with insects, whether permanently or temporarily, play a crucial role in the nutrition, development, reproduction, defence, and metamorphosis regulation. In some Lepidoptera, oviposition-deterrent pheromones (ODPs) on egg surface were used by pregnant females to modify the behaviour of conspecifics to avoid excessive competition for limited resources. In this study, we constructed four different *Spodoptera litura* groups, including OH, OA, SH, and OA, which either feed on different hosts or grow in different environments. The 16S rDNA libraries of microbes from the egg surface of the four groups were constructed and sequenced. According to *alpha* and *beta* diversity indices, the microbes in environments and diets considerably influenced the richness, diversity, and community compositions of the microbiota on egg surfaces. The quantity of the main ODP components and the corresponding oviposition-deterrent activity among four groups were significantly differed among the four groups. The result of this study revealed that altering of microbes in environments or diets considerably changed the contents of ODP and oviposition-deterrent activity. As ODPs impart oviposition-deterrent activity towards closely related species, the findings of this study suggest that we should pay more attention to the role of symbiotic microorganisms in changing the ability of insects, especially sympatric species, to occupy the optimal niche when developing novel pest-control strategies.

Introduction

Pregnant females use oviposition-deterrent pheromones (ODPs) to modify the behaviour of conspecifics to avoid excessive competition for limited resources (Thompson and Pellmyr, 1991; Nufio and Papaj, 2001; Honda, 2010). Extracts from the egg masses of several Noctuid insects, including *Spodoptera littoralis*, *Mamestra configurata*, and *Helicoverpa armigera*, have been proved to have oviposition-deterrent effects on conspecific females (Ulmer *et al.*, 2003; Liu *et al.*, 2008; Gomaa, 2010). *Spodoptera litura* (Lepidoptera: Noctuidae) is a polyphagous pest that seriously threatens world agricultural production and food security (Ahmad *et al.*, 2013), whose preference for good-quality plants is weaker than that of oligophagous insects (Gripenberg, *et al.*, 2010). Pregnant *S. litura* presents a keen sensing ability to select optimal oviposition sites (Calumpang, 2013). However, whether the microbes in the environment and diet influenced the main ODP components and corresponding deterrent activity remain unclear.

Microorganisms symbiotic with insects, whether permanently or temporarily, play a crucial role in the nutrition, development, reproduction, defence, metamorphosis regulation, and numerous other functions in insects (Douglas, 2009; Provorov and Onishchuk, 2018; Malacrino, 2022). Different insects require unique microbial populations in different environments, and sometimes even in different parts of the same insect, such as the salivary gland, midgut, and ovary, to perform unique physiological metabolism (Gimonneau *et al.*, 2014; Qiao *et al.*, 2019; Zhang *et al.*, 2023). For example, symbiotic microbes present in the gut of termites are essential for digesting cellulose and nitrogen required for nutrition throughout their life (Zhou *et al.*, 2019). Female *Costelytra zealandica* use phenol as their sex pheromone,

which is produced by symbiotic bacteria in the accessory or col-
 lerial gland (Marshall *et al.*, 2016). The structure of insect bac-
 terial communities is always shaped by various endogenous and
 exogenous factors, including parental transmission, environment,
 and host-associated bacteria (Wierz *et al.*, 2021).

Many insects use vertical transmission via the egg as a mech-
 anism to ensure the transfer of microorganisms that are added to
 the egg surface by the females in secretions or faeces (Paniagua
 Voiro *et al.*, 2018). In particular, some neonate lepidopteran lar-
 vae bite through their egg shell while hatching and often fully
 ingest it after hatching, which will accelerate the spread of egg-
 surface microorganisms to the next generation (Brinkmann
et al., 2008; Mason and Raffa, 2014).

Certain microorganisms associated with insects have been
 identified as the source of chemicals altering the behaviour of
 conspecifics or other organisms (Xu *et al.*, 2016; Engl and
 Kaltenpoth, 2018; Mazorra-Alonso *et al.*, 2021). *Drosophila* pre-
 fers to mate with conspecifics with a similar gut microbiota,
 which is probably linked to microbiota-dependent variation in
 the cuticle hydrocarbon profile (Sharon *et al.*, 2011).
 Microorganisms play an important role in regulating metabolism,
 reproduction pheromone synthesis, and nutrition (Douglas, 2009;
 Provorov and Onishchuk, 2018; Xu *et al.*, 2019). As an important
 kind of pheromone in insects, ODPs can avoid the fierce competi-
 tion of offspring among the inter- or intra-specifics through
 altering the oviposition behaviour (Anderson and Löfqvist,
 1996; Růžička, 1996), and microorganisms play an important
 role in regulating pheromone synthesis (Engl and Kaltenpoth,
 2018; Mazorra-Alonso *et al.*, 2021; Moyano *et al.*, 2023).
 Therefore, exploring whether microorganisms in the host and
 environment can cause substantial variations in ODP compo-
 nents would provide important insights into the function of
 microorganisms in the competition among sympatric insects for
 host and ecological niches.

Materials and methods

Insect rearing and sample processing

The pupae of *S. litura* were originally purchased from Ke Yun
 Biological Liability Co., Ltd. The larvae were reared with the
 leaves of sweet potato and artificial diet, and the adults were
 offered 10% honey water. Composition (g) of the artificial
 diet was as follows: agar (3), soybean flour (15), wheatgerm
 (15), yeast powder (6.5), casein (3) methyl sorbate (0.375), mul-
 tivitamin (0.15), ascorbic acid (0.6), and sodium benzoate
 (0.375), mixed in 150 ml of distilled water. The insectarium
 was kept at 27°C ± 2°C under 60–70% RH and an L12 (1200
 lux):D12 h photoperiod (Gupta *et al.*, 2005). To better under-
 stand the impact of the microbes on the host and environment
 in relation to the number of ODPs, the pupae were soaked in
 75% alcohol for 55 s, 5–6 days after eclosion. The pupae were
 then rinsed three times with sterilised water in a super-clean
 environment. The water obtained from the third rinse was tested
 for sterility to ensure that the surface of the pupa was sterile.
 The surface-sterilised pupae were divided into four groups:
 OH, SH, SA, and OA. In the SH and SA groups, the
 surface-sterilised pupae, subsequent adults, eggs, and till to
 eggs of next generation were all continuously kept in sterile
 environment. In the SH and SA groups, the adults were both
 fed with sterilised honey water (10%), and the larval were reared
 with sterilised sweet potato leaves (75% alcohol 15 s, then 1%

NaClO₃ 10 s, sterilised water three times) and sterilised artificial
 diet (103.4 kPa, 121.3°C, 15 min), respectively. Differently, the
 larvae in the OH group were kept in an open environment
 and fed with leaves of sweet potato collected directly from the
 outdoors; in the OA group, though the artificial diets were
 also sterilised, the whole insect breeding period was in an
 open environment.

Library construction and high-throughput sequencing

Five hundred eggs from at least three different egg masses were
 collected and washed using 1 ml (1×) sterile phosphate buffered
 saline per sample using a sterile centrifuge tube. Each treatment
 had three replicates. The total genome DNA was extracted from
 the samples using the cetyltrimethylammonium bromide/sodium
 dodecyl sulphate method. The 16S rDNA genes were amplified in
 the hypervariable region of the bacterial 16S rDNA gene V4
 region using the universal primer (F341: 5-CCTAYGGGRBG
 CASCAG-3; 806 R: 5-GGACTACNNGGTATCTAAT-3) with a
 special barcode for each sample. Libraries were generated using
 Illumina TruSeq DNA PCR-Free Library Preparation Kit
 (Illumina, USA) following purification with GeneJET Gel
 Extraction Kit (Thermo Scientific). The library quality was
 assessed using the Qubit@ 2.0 Fluorometer (Thermo Scientific)
 and Agilent Bioanalyzer 2100 system. The library sequencing
 was supported by Novogene Co., Ltd, China, on the Illumina
 NovaSeq PE250 platform. Paired-end reads from the original
 DNA fragments are merged using FLASH (Magoc and Salzberg,
 2011) and filtered via the FASTP software package (Caporaso
et al., 2010). The obtained effective tags, those with ≥97% similar-
 ity, were assigned to the same operational taxonomic units
 (OTUs) based on Uparse (Edgar, 2013). Then, a representative
 sequence was picked for each OTU, and RDP CLASSifier was
 used to annotate taxonomic information based on the
 SILVA138 library via the Mothur method (Wang *et al.*, 2007).

Statistical and bioinformatics analysis

Alpha diversity indices, including PD whole tree, Ace, Shannon,
 and Simpson, were estimated using QIIME software (Version
 1.9.1) and compared using R software based on Wilcox test to
 explore group variations (Dixon, 2003). Principal coordinates
 analysis (PCoA) based on weighted unifracs distance was con-
 ducted to illustrate the variability in the microbiota community
 composition between the groups. Furthermore, the significance
 of the four groups, including OH, SH, OA, and SH, was examined
 using the analysis of molecular variance (AMOVA) method based
 on weighted unifracs distance (Roewer, 1996). Based on the taxo-
 nomic information, the relative abundance of the top ten bacterial
 compositions of *S. litura* egg surface of four groups, including
 OH, SH, OA, and SH, at the phylum and genus levels.
 Furthermore, biomarkers with significant differences among
 groups at different taxonomic levels were identified considering
 four as the threshold of the linear discriminant analysis (LDA)
 score (Segata *et al.*, 2011).

Quantitative analysis of compounds eluted from the egg surface

Five hundred eggs of *S. litura* were collected 6–8 h after oviposi-
 tion in a sterile environment. The eggs were subsequently placed
 in a 2 ml autosampler vial containing 0.5 ml *n*-hexane. After

gentle shaking, the samples were stored at 4°C for 24 h. Next, 0.4 ml supernatant was collected into a new vial and stored at -20°C. Three biological replicates were performed. The extracts were analysed using Shimadzu GC-MS (QP2010 plus) equipped with a DB-WAXETR column (30 m × 0.25 mm ID, 0.25 µm film thickness). Helium was used as the carrier gas (1.0 ml min⁻¹). The temperature programme maintained an initial temperature of 110°C for 4 min, which was increased to 160°C at 10°C min⁻¹ and maintained for 1 min, and subsequently increased to 210°C at 4°C min⁻¹ and maintained for 3 min, and then increased to 240°C and maintained for 8 min. Using ethyl myristate as an internal standard compound, the quantity of six compounds that have been proved as the main ODP components, including palmitic, palmitoleic, stearic, oleic, linoleic, and α-linolenic acids, among the four groups was examined via one-way analysis of variance (ANOVA).

Comparison of oviposition-deterrent activity

Newly emerged adults are placed in a mating cage, allowed to mate freely, and transferred to a new cage after mating to ensure each female has mated, who was placed in a cylinder (diameter: 9 cm; axial length: 16 cm) composed of metal mesh covered with filter paper (diameter: 12 cm) at each end. Compounds were mixed according to the ratio of the quantitative test and diluted in 0.3 ml hexane. The mixture was used as the control and coated evenly on the filter paper on one side of the cylinder cage. As the control, the filter paper on the opposite side was treated only with the same volume of hexane. During the bioassay, ventilation of the room should be turned on to vent odour continuously. Both ends of the test cage maintain the same light intensity. Ten replicates were carried out, and the number of eggs in each side were counted after 48 h. Oviposition-deterrent effect was indicated with oviposition-deterrent indices (ODI) (Lundgren, 1975). The ODI was calculated using the following equation: $ODI = (C - T) / (C + T)$, where *C* is the number of eggs laid on the control, and *T* represents the number of eggs on the treatment. The ODI was compared between the four groups with ANOVA.

Results

rRNA gene sequencing and microbiota comparison of the egg surface among the four groups based on α-diversity index

We obtained effective tags from each sample after merging and filtering the raw tags. The average number of clean tags obtained from the OH, SH, OA, and SA groups were 82,348, 84,204, 79,031, and 81,793, which represented 98.51, 98.37, 98.07 and 98.45% of the total number of raw tags, respectively. The index goods coverage was used to measure sequencing integrity. The coverage ranged from 99.80 to 100%, revealing that the majority of species present in the samples were well sequenced. Detailed information regarding library sequencing is provided in table 1.

The alpha diversity indices, including PD whole tree (fig. 1A), Ace (fig. 1B), Shannon (fig. 1C), and Simpson (fig. 1D) values, demonstrate the bacterial richness and diversity of the egg surface bacterial. Compared with the SH group, the OH group reveals higher PD whole tree, Shannon, and Simpson index values (Wilcox test, $P < 0.05$), which suggest that microorganisms in the environments considerably influence the richness and diversity of bacterial on the egg surface (fig. 1A, C, D). Although there is no significant difference in the Ace index between any two groups, group OH still shows higher bacterial diversity than the SH group (fig. 1B). The impact of environmental microorganisms on bacterial richness and diversity is also revealed by the extremely significant difference in the Simpson index between group OA and group SA (Wilcox test, $P < 0.01$) (fig. 1D).

When supplied with a sterile environment, the diet significantly influences the bacterial richness and diversity between the SH and SA groups, as per the Simpson index (Wilcox test, $P < 0.05$; fig. 1D). This suggests that the endophytic bacteria of the host play a decisive role in the structurally shaping egg surface microorganisms only in a sterile environment.

Microbial community comparison of the egg surface among the four groups based on β-diversity index

A scatter plot was used to show the variability of microbiota communities among different groups based on weighted unifrac

Table 1. Quality summary of libraries construction and sequencing

Sample name	Raw tags	Clean tags	Effective tags	Avg. length (nt)	Q30	Goods_coverage
OH1	90,270	88,839	74,819	426	93.78	0.998
OH1	83,591	82,368	72,098	428	93.3	0.998
OH1	76,922	75,836	70,616	429	93.69	0.999
SH1	87,370	86,200	80,070	429	93.65	0.999
SH1	88,661	87,123	79,538	428	93.04	0.999
SH1	80,776	79,289	73,095	428	93.23	0.998
OA1	77,959	76,262	69,352	428	92.96	0.999
OA1	78,967	77,526	70,680	428	93.5	0.999
OA1	84,833	83,304	75,308	427	93.25	0.999
SA1	82,370	81,133	71,012	427	93.42	1
SA2	85,526	84,061	75,421	428	93.2	0.999
SA3	81,342	80,185	70,981	428	93.88	0.999

OH, group provided with host in open environment; SH, group provided with sterile host and environment; OA, group provided with artificial diets in open environment; SA, group provided with artificial diets in open environment.

Figure 1. Comparison of alpha diversity index, including (A) PD whole tree, (B) Ace (C) Shannon, and (D) Simpson values, of microbial communities on the egg surface of *S. litura* supplied with four diet or environment. Alpha diversity index of each group was compared with Wilcoxon test. * $P < 0.05$; ** $P < 0.01$.





Figure 2. PCoA for microbiota variability based on weighted unifracs distance. Each symbol represents a sample. The variance explained by the PCoA is indicated on the axes.

distance (fig. 2). The first two principal coordinates of the plot, PCoA1 and PCoA2, accounted for 68.19 and 22.08% of the data variation, respectively. The AMOVA analysis revealed a significant difference among the four groups ($P < 0.001$).

The taxonomic information of each OTU at different levels was obtained according to the annotation result in the SILVA138 library. Figure 3A, B show the relative abundance of the top ten bacterial compositions found on the surface of *S. litura* eggs of the OH, SH, OA, and SA groups at the phylum and genus levels, respectively. OTUs that were not assigned to known microbial phyla were designated as ‘Unidentified Bacteria’, which, on average, accounted for 0.69% of the total data (fig. 3A). Microbial phyla with low abundance were grouped together as ‘others’. The Proteobacteria (mean \pm SD = $82.9649 \pm 15.32\%$) was the main microbial phyla of all the samples. However, the relative abundance of Firmicutes phylum in the groups fed with artificial diet, i.e. OA ($37.82 \pm 5.33\%$) and SA

($11.38 \pm 1.32\%$), was significantly higher than that in the host-supplied groups OH ($2.59 \pm 0.66\%$) and SH ($1.73 \pm 1.02\%$); ANOVA, Duncan, $P < 0.05$; fig. 3A). At the genus level (fig. 3B), the major genera found in the samples changed significantly depending on the environment or diet. Acinetobacter ($54.96 \pm 13.48\%$) in the OH groups, Enterobacter ($74.55 \pm 1.97\%$) in the SH groups, Enterococcus ($34.60 \pm 5.24\%$) in the OA groups, and Providencia ($58.56 \pm 2.68\%$) in the SA groups were significantly higher than that in the other groups (ANOVA, Duncan, $P < 0.05$).

To identify the biomarker that revealed significant differences among the groups at different taxonomic levels, LDA was conducted using four as the threshold (fig. 4). At the genus level, some new genera were identified as the biomarker. For example, Enterococcus and Serratia genera were identified as the dominant genus and important biomarker of the OA group, respectively. Similarly, except for Providencia, Tazewell was also identified as the biomarker of the SA group.

Comparison of ODP components among the four groups

Figure 5 shows a decreasing trend in the average levels of six compounds from the OH, SH, and OA groups to the SA group. The OH and SH groups were fed on the host; however, the OH group exhibited significantly higher levels of palmitoleic, stearic, oleic, and linoleic acids compared with the SH group. When supplied with an artificial diet, the levels of palmitic and α -linolenic acids in the SA group were significantly lower than those in the OA group. The comparison between the OH and OA groups, both of which were supplied in open environments, also revealed significant differences in the levels of four components, including palmitoleic, stearic, oleic, and linoleic acids. In a sterile environment, replacing sweet potato leaves with an artificial diet significantly decreased the levels of stearic, oleic, and α -linolenic acids.

Oviposition-deterrent activity assay and comparison

Compounds were mixed according to the ratio of the quantitative test and the ODI among the four groups was compared, and the result revealed that the ODI of the OH group was significantly




Figure 3. Relative abundance of top ten bacterial composition of *S. litura* egg surface supplied with four different diets, including OH, SH, OA, and SH at the phylum at (A) the phylum and (B) genus levels, respectively.




Figure 4. Identification of the bacterial biomarker among the groups at different taxonomic levels; LDA was conducted using four as the threshold. The means of capital letter before the name of biomarkers is phylum (P), class (C), order (O), family (F), genus (G), and species (S).

higher than that of the SH, OA, and SA groups. Comparably, the ODI of the SA group was significantly lower than that of other groups. There was no obvious difference between the SH and

OA groups (fig. 6). These results suggest that changes in the microbial population in environments or diet considerably affect the levels of compounds on the *S. litura* egg surface.




Figure 5. Comparison of ODP contents from the egg surface of *S. litura* provides different diet or environment. Different letters indicate a significant difference in the relative expression levels (one-way ANOVA, $P < 0.05$). OH: group provided with natural host in open environment; SH: group provided with sterile host and environment; OA: group provided with artificial diets in open environment; SA, group provided with artificial diets in open environment.




Figure 6. Comparison of oviposition deterrent activity of ODPs composed of authentic compounds equivalent to 500 eggs among the OH, OA, SH, and SA groups. Different letters indicate a significant difference in the relative expression levels (one-way ANOVA, $P < 0.01$).

Discussion

Microorganisms supply the necessary assistance to insects in different habitats to complete their unique biology, development, fitness, and lifestyle processes (Kellner, 2003; Douglas, 2009; Colman *et al.*, 2012). Phytophagous insects harbour symbiotic bacteria that can be transmitted vertically from parents to offspring, in which the egg is a key link between stages of completely different morphologies.

The specialised structure of the terminal digestive tracts and genital ensures that symbionts are transmitted from the gut to the outside of the eggs (Kellner, 2003). A recent study using 4000 publicly available sequencing data found that besides taxonomy, the origin of the sample (whether from a laboratory or fields) (Shannon $R^2 = 0.087$; $P < 0.001$) and the diet (Shannon $R^2 = 0.036$; $P < 0.001$) were important factors that affected variation in the bacterial diversity of the gut lumen (Malacrino, 2022). Herein, we compared the richness and diversity of microbiota at the egg surface of *S. litura* supplied with different

environments and diets. Compared to diets, environmental factors had a greater influence on bacterial richness and diversity, which agrees with the results of large-scale data analysis of guts.

In the guts of insects from different orders, the dominant microbial phylum was identified as the Proteobacteria or/and Firmicutes phyla (Dillon and Charnley, 2002; Haynes *et al.*, 2003; Schloss *et al.*, 2006; Behar *et al.*, 2008; Li *et al.*, 2022). At the genus level, environment and diet greatly influenced the microbial structure of the *S. litura* guts (Xia *et al.*, 2020). Similar to that in the guts, the result of this study showed that the microbial structure of the egg surface could be influenced by microbes in the environments and diets.

Microorganisms associated with eggs may produce antibiotics that prevent the new hatching larvae from pathogen infection, implying that microbe variations on the egg surface may play an important role in environment adaptability. For example, bacteria on the egg surface of *Holotrichia obliqua* inhibit the multiplication of the entomopathogens *Bacillus thuringiensis* and *Beauveria bassiana* (Wang *et al.*, 2021). ODPs were used by pregnant females to avoid excessive competition for limited resources. The same ODP components shared among closely related species can produce oviposition-deterrent activity to closely related species. For example, a mixture of the five fatty acids found in *Ostrinia zealis* exhibited significant oviposition-deterrent effects on all the three other *Ostrinia* species (Li and Ishikawa, 2004). Herein, *S. litura* ODPs were identified, and the main components were fatty acids, some of which were shared with other lepidopteran insects, such as *Spodoptera frugiperda*, *H. armigera* (Li *et al.*, 2001), *Ostrinia furnacalis* (Li and Ishikawa, 2004), and *Phthorimaea operculella* (Zhang *et al.*, 2018). The relationship between the relative levels of unsaturated fatty acids in eggs and microorganisms was reported in an early study (Jackson *et al.*, 1968). The results of this study revealed that the quantity of ODPs and the deterrent activity will be influenced by the microorganisms depending on environments and diets. ODP component changes may cause alteration in the abilities of intraspecific and interspecific insects to occupy the most suitable niches. Chemical substances present

on the egg surface and egg-associated microorganisms could trigger downstream defence responses in the host (Hilfiker *et al.*, 2014; Berteza *et al.*, 2020; Li *et al.*, 2023), and the induced plant volatiles could be used as synomones in attracting natural enemies (Fatouros *et al.*, 2008, 2005; Conti *et al.*, 2010). The findings of this study suggest that we should pay more attention to the role of microorganisms in changing the ability of insects, especially sympatric species, to occupy the optimal niche when developing novel pest-control strategies.

Acknowledgements. This work was funded by the national key R&D programme of China (2022YFD1401200); The special projects in key areas of ordinary universities in Guangdong province (2021|ZDZX4020; 2023ZDZX4019); The Science and Technology Planning Project of Guangzhou, China (2023B03J1268); The Zhongkai University of Agriculture and Engineering – Plant Protection Research Institute, Guangdong Academy of Agricultural Sciences joint graduate training demonstration base; Guangdong teaching and research Letter (2021, No. 2); the Zhongkai University of Agriculture and Engineering – Institute of Nanfan & Seed Industry, Guangdong Academy of Sciences joint graduate training demonstration base; Guangdong teaching and research letter (2022, No. 1).

Author contributions. Jianmei Shen and Zhenfei Zhang designed the study; Liming Hu wrote the manuscript; Yirui Chen mainly performed quantitative analysis of compounds; and all authors have equally important contributions in other instrument operation and data analysis.

Competing interests. None.

References

- Ahmad M, Ghaffar A and Rafiq M (2013) Host plants of leaf worm, *Spodoptera litura* (Fabricius) (Lepidoptera: Noctuidae) in Pakistan. *Asian Journal of Agriculture and Biology* **2013**, 23–28.
- Anderson P and Löfqvist J (1996) Asymmetric oviposition behaviour and the influence of larval competition in the two pyralid moths *Ephestia kuehniella* and *Plodia interpunctella*. *Oikos* **76**, 47–56.
- Behar A, Yuval B and Jurkevitch E (2008) Gut bacterial communities in the Mediterranean fruit fly (*Ceratitidis capitata*) and their impact on host longevity. *Journal of Insect Physiology* **54**, 1377–1383.
- Berteza CM, Casacci LP, Bonelli S, Zampollo A and Barbero F (2020) Chemical, physiological and molecular responses of host plants to lepidopteran egg-laying. *Frontiers in Plant Science* **10**, 1768.
- Brinkmann N, Martens R and Tebbe CC (2008) Origin and diversity of metabolically active gut bacteria from laboratory-bred larvae of *Manduca sexta* (Sphingidae, Lepidoptera, Insecta). *Applied and Environmental Microbiology* **74**, 7189–7196.
- Calumpang SMF (2013) Behavioral response of *Spodoptera litura* (F) (Lepidoptera: Noctuidae) to selected herbs and eggplant. *Journal of the International Society for Southeast Asian Agricultural Sciences* **19**, 95–103.
- Caporaso JG, Kuczynski J, Stombaugh J, Bittinger K, Bushman FD, Costello EK, Fierer N, Gonzalez Peña A, Goodrich JK, Gordon JI, Hutley GA, Kelley ST, Knights D, Koenig JE, Ley RE, Lozupone CA, McDonald D, Muegge BD, Pirrung M, Reeder J, Sevinsky JR, Turnbaugh PJ, Walters W, Widmann J, Yatsunenko T, Zaneveld JR and Knight R (2010) QIIME allows analysis of high-throughput community sequencing data. *Nature Methods* **7**, 335–336.
- Colman DR, Toolson EC and Takacs-Vesbach CD (2012) Do diet and taxonomy influence insect gut bacterial communities? *Molecular Ecology* **21**, 5124–5137.
- Conti E, Salerno G, Leombruni B, Frati F and Bin F (2010) Short-range allelochemicals from a plant–herbivore association: a singular case of oviposition-induced synomone for an egg parasitoid. *Journal of Experimental Biology* **213**, 3911–3919.
- Dillon R and Charnley K (2002) Mutualism between the desert locust *Schistocerca gregaria* and its gut microbiota. *Research in Microbiology* **153**, 503–509.
- Dixon P (2003) A package of R functions for community ecology. *Journal of Vegetation Science* **14**, 926–937.
- Douglas AE (2009) The microbial dimension in insect nutritional ecology. *Functional Ecology* **23**, 38–47.
- Edgar RC (2013) UPARSE: highly accurate OTU sequences from microbial amplicon reads. *Nature Methods* **10**, 996–998.
- Engl T and Kaltenpoth M (2018) Influence of microbial symbionts on insect pheromones. *Natural Product Reports* **35**, 386–397.
- Fatouros NE, Bukovinszkine-Kiss G, Kalkers LA, Soler Gamborena R, Dicke M and Hilker M (2005) Oviposition-induced plant cues: do they arrest trichogramma wasps during host location? *Entomologia Experimentalis et Applicata* **115**, 207–215.
- Fatouros NE, Broekgaarden C, Bukovinszkine-Kiss G, van Loon JJA, Mumm R, Huigens ME, Dicke M and Hilker M (2008) Male-derived butterfly anti-aphrodisiac mediates induced indirect plant defense. *Proceedings of the National Academy of Sciences of the United States of America* **105**, 10033–10038.
- Gimonneau G, Tchioffo MT, Abate L, Boissière A, Awono-Ambene PH, Nsango SE, Christen R and Morlais I (2014) Composition of *Anopheles coluzzii* and *Anopheles gambiae* microbiota from larval to adult stages. *Infection Genetics and Evolution* **28**, 715–724.
- Gomaa HA (2010) Secretion of *Spodoptera littoralis* female reproductive system on deposited egg masses that possibly acts as oviposition deterring substance for conspecific females. *Archives of Phytopathology and Plant Protection* **43**, 1081–1087.
- Gripenberg S, Mayhew PJ, Parnell M and Roslin T (2010) A meta-analysis of preference-performance relationships in phytophagous insects. *Ecology Letters* **13**, 383–393.
- Gupta GP, Rani S, Birah A and Raghuraman M (2005) Improved artificial diet for mass rearing of the tobacco caterpillar, *Spodoptera litura* (Lepidoptera: Noctuidae). *International Journal of Tropical Insect Science* **25**, 55–58.
- Haynes S, Darby AC, Daniell TJ, Webster G, Van Veen FJF, Godfray HCJ, Prosser JI and Douglas AE (2003) Diversity of bacteria associated with natural aphid populations. *Applied and Environmental Microbiology* **69**, 7216–7223.
- Hilfiker O, Groux R, Bruessow F, Kiefer K, Zeier J and Reymond P (2014) Insect eggs induce a systemic acquired resistance in Arabidopsis. *The Plant Journal: for Cell and Molecular Biology* **80**, 1085–1094.
- Honda K (2010) Chemical basis of differential oviposition by lepidopterous insects. *Archives of Insect Biochemistry and Physiology* **30**, 1–23.
- Jackson LL, Baker GL and Henry JE (1968) Effect of malamoeba locustae infection on the egg lipids of the grasshopper *Melanoplus bivittatus*. *Journal of Insect Physiology* **14**, 1773–1778.
- Kellner RLL (2003) The role of microorganisms for eggs and progeny. In Hilker M and Meiners T (eds), *Chemoecology of Insect Eggs and Egg Deposition*. Berlin, Vienna: Blackwell Publishers, pp. 149–164.
- Li GQ and Ishikawa Y (2004) Oviposition deterrents in larval frass of four *Ostrinia* species fed on an artificial diet. *Journal of Chemical Ecology* **30**, 1445–1456.
- Li GQ, Han ZJ, Mu LJ, Qin XR, Chen CK and Wang YC (2001) Natural oviposition-deterrent chemicals in female cotton bollworm, *Helicoverpa armigera* (Hubner). *Journal of Insect Physiology* **47**, 951–956.
- Li DD, Li JY, Hu ZQ, Liu TX and Zhang SZ (2022) Fall armyworm gut bacterial diversity associated with different developmental stages, environmental habitats, and diets. *Insects* **13**, 762.
- Li B, Liu FJ, He X, Liu XL and Lu M (2023) Temporal transcriptomic changes in willow leaves oviposited by *Plagioderma versicolora*. *Integrative Zoology* **19**, 784–787.
- Liu ML, Yu HJ and Li GQ (2008) Oviposition deterrents from eggs of the cotton bollworm, *Helicoverpa armigera* (Lepidoptera: Noctuidae): chemical identification and analysis by electroantennogram. *Journal of Insect Physiology* **54**, 656–662.
- Lundgren LN (1975) Natural plant chemicals acting as oviposition deterrents on cabbage butterflies (*Pieris brassicae* (L.), *P. rapae* (L.) and *P. napi* (L.)). *Zoologica Scripta* **4**, 253–258.
- Magoc T and Salzberg SL (2011) FLASH: fast length adjustment of short reads to improve genome assemblies. *Bioinformatics (Oxford, England)* **27**, 2957–2963.

- Malacrinò A** (2022) Host species identity shapes the diversity and structure of insect microbiota. *Molecular Ecology* **31**, 723–735.
- Marshall DG, Jackson TA, Unelius CR, Wee SL, Young SD, Townsend RJ and Suckling DM** (2016) *Morganella morgani* bacteria produces phenol as the sex pheromone of the New Zealand grass grub from tyrosine in the colleterial gland. *Die Naturwissenschaften* **103**, 59.
- Mason CJ and Raffa KF** (2014) Acquisition and structuring of midgut bacterial communities in gypsy moth (Lepidoptera: Erebidæ) larvae. *Environmental Entomology* **43**, 595–604.
- Mazorra-Alonso M, Tomás G and Soler JJ** (2021) Microbially mediated chemical ecology of animals: a review of its role in conspecific communication, parasitism and predation. *Biology* **10**, 274.
- Moyano A, Croce AC and Scolari F** (2023) Pathogen-mediated alterations of insect chemical communication: from pheromones to behavior. *Pathogens (Basel, Switzerland)* **12**, 1350.
- Núño CR and Papaj DR** (2001) Host marking behavior in phytophagous insect and parasitoids. *Entomologia Experimentalis et Applicata* **99**, 273–293.
- Paniagua Voirol LR, Frago E, Kaltenpoth M, Hilker M and Fatouros NE** (2018) Bacterial symbionts in Lepidoptera: their diversity, transmission, and impact on the host. *Frontiers in Microbiology* **9**, 556.
- Provorov NA and Onishchuk OP** (2018) Microbial symbionts of insects: genetic organization, adaptive role, and evolution. *Microbiology (Reading, England)* **87**, 151–163.
- Qiao HL, Keesey IW, Hansson BS and Knaden M** (2019) Gut microbiota affects development and olfactory behavior in *Drosophila melanogaster*. *Journal of Experimental Biology* **222**, jeb192500.
- Roewer L** (1996) Analysis of molecular variance (AMOVA) of Y-chromosome-specific microsatellites in two closely related human populations. *Human Molecular Genetics* **5**, 1029–1033.
- Růžička Z** (1996) Oviposition-deterrent pheromone in Chrysopidae (Neuroptera): intra- and interspecific effects. *European Journal of Entomology* **93**, 161–166.
- Schloss PD, Delalibera IT, Handelsman J and Raffa KF** (2006) Bacteria associated with the guts of two wood-boring beetles: *Anoplophora glabripennis* and *Saperda vestita* (Cerambycidae). *Environmental Entomology* **35**, 625–629.
- Segata N, Izard J, Waldron L, Gevers D, Miropolsky L, Garrett WS and Huttenhower C** (2011) Metagenomic biomarker discovery and explanation. *Genome Biology* **12**, R60.
- Sharon G, Segal D, Ringo JM, Hefetz A, Zilber-Rosenberg I and Rosenberg E** (2011) Commensal bacteria play a role in mating preference of *Drosophila melanogaster*. *Proceedings of the National Academy of Sciences of the United States of America* **107**, 20051–20056.
- Thompson JN and Pellmyr O** (1991) Evolution of oviposition behavior and host preference in Lepidoptera. *Annual Review of Entomology* **36**, 65–89.
- Ulmer B, Gillott C and Erlandson M** (2003) Conspecific eggs and bertha armyworm, *Mamestra configurata* (Lepidoptera: Noctuidæ), oviposition site selection. *Environmental Entomology* **32**, 529–534.
- Wang Q, Garrity GM, Tiedje JM and Cole JR** (2007) Naive Bayesian classifier for rapid assignment of rRNA sequences into the new bacterial taxonomy. *Applied and Environmental Microbiology* **73**, 5261–5267.
- Wang K, Liu Q, Liu CQ, Geng LL, Wang GR, Zhang J and Shu CL** (2021) Dominant egg surface bacteria of *Holotrichia oblita* (Coleoptera: Scarabaeidae) inhibit the multiplication of *Bacillus thuringiensis* and *Beauveria bassiana*. *Scientific Reports* **11**, 9499.
- Wierz JC, Gaube P, Klebsch D, Kaltenpoth M and Flórez LV** (2021) Transmission of bacterial symbionts with and without genome erosion between a beetle host and the plant environment. *Frontiers in Microbiology* **12**, 715601.
- Xia X, Lan B, Tao X, Lin J and You M** (2020) Characterization of *Spodoptera litura* gut bacteria and their role in feeding and growth of the host. *Frontiers in Microbiology* **11**, 1492.
- Xu L, Lu M, Xu D, Chen L and Sun J** (2016) Sexual variation of bacterial microbiota of *Dendroctonus valens* guts and frass in relation to verbenone production. *Journal of Insect Physiology* **95**, 110–117.
- Xu L, Deng J, Zhou F, Cheng C, Zhang L, Zhang J and Lu M** (2019) Gut microbiota in an invasive bark beetle infected by a pathogenic fungus accelerates beetle mortality. *Journal of Pest Science* **92**, 343–351.
- Zhang XG, Li X, Gao YL, Liu Y, Dong WX and Xiao C** (2018) Oviposition deterrents in larval frass of potato tuber worm moth, *Phthorimaea operculella* (Lepidoptera: Gelechiidae). *Neotropical Entomology* **48**, 496–502.
- Zhang YX, Zhang SK and Xu LT** (2023) The pivotal roles of gut microbiota in insect plant interactions for sustainable pest management. *Npj Biofilms and Microbiomes* **9**, 66.
- Zhou J, Duan J, Gao M, Wang Y, Wang X and Zhao K** (2019) Diversity, roles, and biotechnological applications of symbiotic microorganisms in the gut of termite. *Current Microbiology* **76**, 755–761.