

restricted interests and behaviors. ASD is a lifelong neurodevelopmental disorder, however there is a lack of answers and research for adults with ASD. There are shared aspects of odd thinking, rigid behaviors and impaired socialization in schizophrenia and ASD and COS seems to have a strong relationship with ASD, being comorbid in up to 50% of cases.

**Conclusions:** Usually the evaluation of the developmental history of the person, prodrome and onset, its course and the presence of positive symptoms of schizophrenia is enough to help us find a diagnosis. Unfortunately, in some ages the conclusion is not so easy to find. However is essential to determine whether the clinical manifestations belong to the autistic spectrum, the schizophrenic or result from comorbidity.

**Keywords:** autism; childhood onset schizophrenia; childhood psychosis; Early-onset schizophrenia

## EPP1240

### What is the best approach for patients with prolonged duration of untreated psychosis (DUP) - about 2 clinical cases with dup longer than 10 years

R. Gomes<sup>1\*</sup>, C. Santos<sup>2</sup> and F. Moutinho<sup>1</sup>

<sup>1</sup>Psychiatry And Mental Health Department, Hospital Garcia de Orta, E.P.E., Almada, Portugal, Portugal and <sup>2</sup>Psychiatry, Hospital Garcia de Orta, Almada, Portugal

\*Corresponding author.

doi: 10.1192/j.eurpsy.2021.1443

**Introduction:** Studies have consistently found that many individuals with psychosis experience significant delays before receiving treatment. DUP refers to the period between the emergence of psychotic symptoms and the initiation of appropriate clinical treatment.

**Objectives:** To review current knowledge on the best approach for patients with schizophrenia (SCZ) and prolonged DUP.

**Methods:** Non-systematic review of literature through search on PubMed database, following the terms “DUP and treatment” and “impact of longer DUP”. Two clinical cases are described.

**Results:** The clinical cases describe patients with SCZ with DUPs older than 10 years, in whom we could not achieve complete clinical remission after several therapeutic trials and whose prognosis was admitted as reserved. Longer DUP is an independent predictor of poorer outcome in SCZ, including the poor response to treatment and difficulty in achieving remission, predicting treatment resistance. Identifying treatment-resistant patients is crucial due to the importance of initiating clozapine as early as possible since the chances of responding are higher.

**Conclusions:** DUP is a key prognostic variable in psychosis, revealing the significance of early treatment. Patients with long DUP should be regarded as at high risk of poor recovery. The detection of these patients enables clinicians to avoid unnecessary exposure to ineffective treatments while effective interventions are delayed. However, in view of adverse side effects of clozapine, future studies need to examine relevant predictors to detect accurately non-responders. We also suggest further studies to understand if there is correspondence between DUP and different stages of the disease that justify these results.

**Keywords:** schizophrénia; duration of untreated psychosis; psychosis; clozapine

## EPP1242

### Adherence to treatment in patients with delusional disorder - study of acute inpatient population in psychiatry ward between 2007-2017

A. Delgado<sup>1\*</sup>, D. Barbosa<sup>2</sup> and R. Curreal<sup>1</sup>

<sup>1</sup>Psychiatry, Centro Hospitalar Universitário de São João, Porto, Portugal and <sup>2</sup>Psychiatry, Sao Joao Hospital and University Centre, Porto, Portugal

\*Corresponding author.

doi: 10.1192/j.eurpsy.2021.1444

**Introduction:** Delusional disorder is a mental illness in which delusions are the dominant symptom. Delusional disorder is not well studied relative to other psychotic disorders - it is poorly understood in practically every aspect of its nature, including cause, phenomenology, prevalence, comorbidity, course, treatment, and prognosis.

**Objectives:** To study the clinical and sociodemographic characteristics of individuals admitted for inpatient treatment with the diagnosis of delusional disorder, in particular the adherence to treatment.

**Methods:** Retrospective observational study of inpatient treatment of patients with delusional disorder diagnosis between January 1<sup>st</sup> 2007 and 31<sup>th</sup> December of 2017 in the Psychiatry Service of CHUSJ. Follow up of 2 years from discharge. Data collected included sociodemographic characteristics and clinical features. Descriptive analysis of the results was performed using SPSS (v.26).

**Results:** In the period of time analyzed, 152 hospitalizations were identified, corresponding to 114 patients: 38.2% male and 62.8% female. The average age was 58 years. 3 months after discharge: 65% of patients were going to the medical appointments, which dropped to 60% in 6 months, 55% in 12 months, 53% in 12 and 24 months. Regarding adherence to the treatment: 65% of patients were still adherent to medication in 3 months time, 55% in 6 months, dropping to 50% in a year and to 48% in 2 years. There is a relation between involuntary discharge and adherence to consultations and medication.

**Conclusions:** A cardinal characteristic of delusional disorder, conviction that one is not mentally ill, contributes complexity to the treatment challenges and profoundly affects the therapeutic relationship.

**Keywords:** Delusional disorder; treatment adhesion

## EPP1245

### The impact of social cognition deficits on real life functioning in 22q11.2 deletion syndrome: A comparative study with a large population of patients with schizophrenia.

T. Accinni\*, M. Frascarelli, A. Buzzanca, F. Ghezzi, L. Carlone, A. Panzera, A. Moschillo, N. Girardi and F. Di Fabio

Department Of Human Neurosciences Sapienza University Of Rome, Viale dell'Università 30, Rome, Italy

\*Corresponding author.

doi: 10.1192/j.eurpsy.2021.1445

**Introduction:** 22q11.2 Deletion Syndrome (22q11.2DS) represents a congenital syndrome with several clinical features. It entails a 25% risk of psychotic onset in lifespan. 22q11.2DS is a reliable model for biological vulnerability to schizophrenia.