

therapies into doctor-patient interactions to allow increased communication and understanding.

### Why Should ACT Work When CBT Has Failed? a Study Assessing Acceptability and Feasibility of Acceptance and Commitment Therapy (ACT) for Paediatric Patients With Chronic Fatigue Syndrome/myalgic Encephalomyelitis (CFS/ME)

Dr Jamie Leveret<sup>1\*</sup>, Dr Jen Starbuck<sup>2</sup>, Dr Kate Chapple<sup>2</sup>, Dr Sam Gubb<sup>1</sup>, Ms Hannah Kwuo<sup>1</sup>, Dr Sarah Burge<sup>1</sup>, Mr Morain Li<sup>1</sup>, Dr Philippa Clery<sup>1</sup> and Professor Esther Crawley<sup>1,2</sup>

<sup>1</sup>University of Bristol, Bristol, United Kingdom and <sup>2</sup>Royal United Hospital Bath, Bath, United Kingdom

\*Presenting author.

doi: 10.1192/bjo.2022.209

**Aims.** Paediatric chronic fatigue syndrome/myalgic encephalomyelitis (CFS/ME) affects 0.5–3.28% of children. NICE guidance recommends Activity Management, Graded Exercise Therapy or Cognitive Behavioural Therapy for fatigue (CBT-f). Approximately 15% of patients do not achieve full recovery within one year with current treatments. Acceptance and Commitment Therapy (ACT) is an effective treatment in many chronic illnesses. There are no studies investigating ACT for paediatric CFS/ME. This feasibility study aimed to assess if ACT is a feasible and acceptable alternative treatment when current treatment has not led to recovery.

**Methods.** This feasibility cohort study aimed to enrol a minimum of 12 participants aged 11–18 years with CFS/ME attending the Royal United Hospitals Bath NHS Foundation Trust Specialist Paediatric CFS/ME Service, who were still symptomatic after 12 months or 12 sessions of standard treatment and were offered six to 12 sessions of ACT. Retention and recruitment data were analysed. Participants were asked to complete questionnaires before, during and after treatment. A selection of participants and their parents were interviewed about their experience of the study. Interviews were analysed using thematic analysis.

**Results.** 19 participants (95% of those approached) were recruited. Only 4 participants of this hard-to-reach group did not complete treatment.

In almost all sessions participants reported that they felt ‘totally listened to in post session questionnaires (31/33 sessions).

Preliminary interviews (n = 12) indicate acceptability of ACT, with all young people and their parents stating that they thought ACT should be offered to this population. Participants particularly commented that the absence of thought challenging (used in CBT-f) was a positive element of ACT. Participant’s openness to try new approaches and altruistic desire to be in a study was noted. **Conclusion.** Recruitment data indicate that it is feasible to recruit and retain 11–18-year-olds with CFS/ME to a study offering ACT. Interviews with participants and parents were broadly positive suggesting ACT is an acceptable treatment in this population.

Results indicated that it is both feasible and acceptable to offer ACT to 11–18-year-olds with CFS/ME using this protocol, supporting the prospect of an RCT in this area.

### Examining Grey Matter Structural Abnormalities in Young People Exposed to Childhood Maltreatment and Peer Victimization

Dr Lena Lim\*

SICS, ASTAR, Singapore, Singapore. IoPPN, King’s College London, London, United Kingdom

\*Presenting author.

doi: 10.1192/bjo.2022.210

**Aims.** Early-life interpersonal stress, particularly childhood maltreatment (CM), is associated with neurobiological abnormalities. However, few studies have investigated the neural effects of peer victimisation (PV). This study examines the common and specific associations between CM, PV and brain structural alterations in healthy youths.

**Methods.** Grey matter volume (GMV) and cortical thickness (CT) data were collected from 105 age- and gender-matched healthy youths (34 CM, 35 PV and 36 controls). Region-of-interest (ROI) and whole-brain analyses were conducted.

**Results.** For the ROI, the CM group had smaller GMV than controls in left IFG, bilateral anterior insula, postcentral and lingual regions, which were associated with higher emotional abuse, along with smaller insular GMV than the PV group. The PV group had smaller left lingual GMV than controls, which was positively associated with age of bully onset. At the whole-brain level, both CM and PV groups had smaller GMV than controls in a cluster comprising left post/pre-central, inferior frontal, insula, superior parietal and supramarginal gyri. The PV group alone had increased CT in a cluster comprising left superior frontal, anterior cingulate and medial orbitofrontal gyri, which was related to greater cyberbullying.

**Conclusion.** Early-life interpersonal stress from carers and peers is associated with common structural alterations of the inferior frontal-limbic, sensory and lingual regions involved in cognitive control, emotion and sensory processing. The findings of a CM-specific reduced anterior insular GMV and a PV-specific increased CT in the left medial prefrontal cluster is intriguing and underscores the unique negative effects of CM and PV, particularly cyberbullying.

### A Case-Control Study Measuring Mentalization in Individuals With PTSD Compared to Controls Using the STOMP Task

Dr Poppy MacInnes<sup>1\*</sup> and Dr Chantelle Wiseman<sup>2,3</sup>

<sup>1</sup>Royal United Hospital, Bath, United Kingdom; <sup>2</sup>University of Bristol, Bristol, United Kingdom. and <sup>3</sup>University of Cardiff, Cardiff, United Kingdom

\*Presenting author.

doi: 10.1192/bjo.2022.211

**Aims.** Social cognition is impaired in a variety of psychiatric conditions; evidence for impairment in individuals with PTSD is increasing. Mentalization is one domain of social cognition that refers to the capacity to understand other people by ascribing mental states to them. The STOMP task (Spontaneous Theory of Mind Protocol) involves an individual watching two minutes of a silent video and describing what they see. As part of a wider project examining social cognition in PTSD, we aimed to find out whether mentalization in the STOMP task differs between patients with PTSD compared to controls.

**Methods.** 171 individuals undertook the task: 30 patients were recruited from centres in Cardiff and Bristol at the start of their psychological therapy; 141 controls were recruited through Prolific website. Participants watched a 2-minute silent video and were asked to write 7–10 sentences about the clip. Qualtrics software selected the video and collected the texts. The verbs of the texts were coded and given a score by PM